

## **OPPOSITE THRUST-VERGENCIES IN THE PRECORDILLERA AND WESTERN CORDILLERA IN NORTHERN CHILE AND STRUCTURALLY LINKED CENOZOIC PALEOENVIRONMENTAL EVOLUTION**

Reynaldo CHARRIER(1-7), Gérard HERAIL(2-7), John FLYNN(3), Rodrigo RIQUELME(1-7),  
Marcelo GARCIA(4-7), Darin CROFT(5) and André WYSS(6)

(1) Departamento de Geología, Universidad de Chile; rcharrie@cec.uchile.cl.

(2) IRD, 209-213, Rue La Fayette, 75010 Paris, gherail@paris.ird.fr.

(3) The Field Museum of Natural History, Chicago, USA. flynn@fmppr.fmnh.org.

(4) SERNAGEOMIN. Av. Sta. María 0104, Providencia, Santiago, Chile. mgarcia@sernageomin.cl.

(5) The Field Museum of Natural History, Chicago, and University of Chicago, USA.

(6) Department of Geological Sciences, University of California Santa Barbara, USA  
wyss@magic.geol.ucsb.edu.

(7) Convenio IRD-Departamento de Geología, Universidad de Chile.

**KEY WORDS:** Altiplano, Chile, syntectonic sedimentation, Cenozoic paleoenvironment.

### **INTRODUCTION**

Morphologic units in the northern Chilean Andes parallel one another and the offshore trench. These units are, from west to east: Coastal Cordillera, Central Depression, Precordillera, Western Cordillera and Altiplano (Fig. 1). The development of the west part of the Western Cordillera (Chapiquiña-Belén Ridge), in the Chilean Altiplano, was controlled by two diverging, trench-parallel systems of thrusts and folds, one located along the Precordillera and the other along the eastern side of the Western Cordillera. Although total shortening associated with these systems is only 12 to 14 km, their activity determined the development of fluvial and lacustrine basins which recorded the synorogenic paleoenvironmental evolution of this region. We describe here facies, geometry and chronology of the deposits associated with both systems and provide an explanation for the differing depositional environments developed on the Precordillera and the Western Cordillera.

## STRUCTURE OF THE PRECORDILLERA AND WESTERN CORDILLERA

A compressive episode began in this part of the Andes during the Early Miocene (approx. 18 Ma). Along the Precordillera a west-vergent thrust system developed in normal sequence: the Belén-Tignámar and Copaquilla-Tignámar thrust faults. Coarse sedimentary wedges were deposited proximal to these faults: the Joracane (18 to 16 Ma) and Huaylas (post-11 Ma to pre-5 Ma) Fms. Small displacements of parts of the Huaylas Ignimbrite (4.5 Ma), (which unconformably covers the Huaylas Fm.) suggest that compressive conditions lasted until the Pliocene. On the west side of the Chapiquiña-Belén Ridge an east-vergent thrust system caused the development of progressive unconformities and associated syntectonic sedimentary deposits, the Chucal Fm.

*Syntectonic fluvial deposits along the Precordillera: Joracane and Huaylas Fms.* Strong erosion of the relief generated by the activity of the Belén-Tignámar and the Copaquilla-Tignámar thrust-faults supplied abundant detrital fill for the depressed areas west of the faults. The Joracane Fm., close to the Belén-Tignámar fault, corresponds to coarse, proximal fluvial facies deposited by braided river systems with low sinuosity. No fossils (plants or animals) were reported from this unit.

The fluvial, upward fining and coarsening Huaylas Fm. is associated with the Copaquilla-Tignámar fault. The lower levels are composed of thin layers deposited in a flood plain environment. Grain size increases gradually upward; near the top it contains 50 cm in diameter blocks. Lower layers tilt slightly to the west, while the upper levels are horizontal. To the east they locally cover the Copaquilla-Tignámar fault as well as the deformed late Oligocene?-early Miocene Lupica Fm. To the west they cover the scarps at the foot of the fault and they overlap the Oxaya Fm. at the east-flank of the Oxaya Anticline. The thickness of the Huaylas Fm. varies considerably. It filled a network of west draining paleovalleys that reached the present Central Depression. The present topographic low in which most of the outcrops of the Huaylas Fm occur was never an endorheic basin. The lower levels contain fossil mammal remains (Notoungulates) (Bargo and Reguero, 1989; Salinas et al., 1991) of the Huayquerian (Salinas et al., 1991) South American Land Mammal Age (SALMA) of about 9 to 7 Ma (Flynn and Swisher, 1995).

*Syntectonic fluvial and lacustrine deposits on the east side of the Chapiquiña-Belén Ridge: Chucal Fm.* East of the Chapiquiña-Belén Ridge the Lupica Fm. is eastwardly deformed by a partially blind, east-vergent thrust system (Chucal thrust-system). This system, interpreted as a backthrusting of the west-vergent thrust system located along the Precordillera (Hérail et al., submitted), caused the development of progressive unconformities and associated syntectonic sedimentary deposits: lower member (post-21 Ma) and upper member (pre-11 Ma) of the Chucal Fm., that transitionally grades to the late to post-tectonic Lauca Fm. The uplift caused by the western fault (Chucal fault) of this system, exposed immediately west of the Chapiquiña-Belén Ridge, supplied the sediments that formed the Chucal Fm. and part of the Lauca Fm. The fluvial and lacustrine Chucal Fm. (Muñoz, 1991; Riquelme and Hérail, 1997) rest unconformably on the Lupica Fm. and is deformed by a growth fold. Although the Chucal and Lauca Fms. were deposited in fluvial and lacustrine environments, fluvial facies predominate in the Chucal Fm., while lacustrine environments

prevail in the Lauca Fm. The lower member of the Chucal Fm. begins with immature conglomerates composed of ignimbrite fragments, lavas of the Lupica Fm. and fluvial sandstones. Provenance is from the west and southwest. These are followed by siltstones, marls and limestones in decimeter thick layers deposited in flood plain and lacustrine environments. The lacustrine deposits contain well preserved plant stems and leaves. Vertebrate fossils have not been found in this sequence. The series continues with mudstones and siltstones deposited in a flood plain environment, and containing a rich mammal fauna. The upper member begins with a coarse fluvial sequence composed of green conglomerates, followed by sandstones and siltstones with limestone intercalations at the top, also deposited in flood plain and lacustrine (more restricted) environments ; it contains no mammal fossils. The sequence evolves upward toward higher energy deposits, which are also fossiliferous, and are covered to the west by the Chucal fault and to the east, in the basin, by the  $11.2 \pm 0.5$  Ma Chucal Ignimbrite. The Chucal Ignimbrite is covered by conglomerates composed of andesitic clasts, on top of which rests the Lauca Fm.

### FOSSIL MAMMAL CONTENT

An extensive mammalian fauna has been recently recovered from the Chucal Fm. (Charrier et al., 1994a; Flynn et al., 1999). Specimens were collected from different stratigraphic levels. The current faunal list includes: hegetotheriine hegetothere, at least 2 mesotheriine mesotheres, toxodontid, macraucheniid litoptern, chinchillid rodent, armadillo, turtle carapace pieces, and several bird bones (large and small size).

Taxa in the assemblages found in northernmost Chile range elsewhere from Santacrucian to Chasicuan or Huayquerian, with most overlapping in the "Friasian". As these are the northernmost Cenozoic mammalian fauna(s) known from Chile, these assemblages from the Huaylas and Chucal Formations permit comparisons of an extensive latitudinal series (more than 30 degrees) of middle Cenozoic faunas from west of the Andean crest (Casamayoran? and "Tinguirirican" [pre-Deseadan, post-Mustersan] through Santacrucian and the type "Friasian"). The occurrence of these faunas at a critically important modern biotic disjunction (Atacama Desert – Bolivian Orocline bending axis), directly west of well known Cenozoic faunas from a variety of paleoelevations in the Altiplano and other regions of Bolivia, may allow assessment of the biotic history along an east-west transect from Chile to eastern Bolivia.

### CONCLUSION

Tectonic evolution along the western Altiplano has occurred under prevailing compressive condition since 18 Ma creating a west-vergent thrust system and some east-vergent backthrusts associated with growth folds. These diverging systems generated an uplifted ridge located along the Precordillera and the present Western Cordillera. During development of these structures and uplift of the ridge, considerable amounts of syntectonic deposits containing mammalian remains accumulated at both sides of the ridge.

The west side of the ridge, was characterized geomorphologic conditions that favoring aridity and rapid drainage of rain and melt water toward the Central Depression, and explaining the relative scarcity of fossil faunas and the absence of fossil floras. On the east side, on the Chilean Altiplano, geomorphologic conditions favored the development of flood plains and shallow lakes with considerable accumulation of finer sediments containing abundant mammalian remains. This indicates that, as a consequence of the Middle Miocene tectonic activity, a partition of the environmental conditions occurred west and east of the present Western Cordillera. According to the paleofloras known from this Andean region, low altitudes in Chile (Charrier et al., 1994b and in Bolivia (Berry, 1919, 1922; Gregory-Wodzicki et al., 1998), this partition is better attributed to a topographic barrier, the Chapiquiña-Belén Ridge than to Andean uplift.

## REFERENCES

- Bargo, M.S. and Reguero, M.A., 1989. El primer registro de un mamífero fósil en el extremo septentrional de Chile. *Ameghiniana*, 26, 3-4, p.239.
- Berry, E.W.1919. Fossil plants from bolivia and their bearing upon the age of uplift of the eastern Andes. *proceeding U.S. National Museum*, 54, 103-164.
- Berry, E.W.1922. Late Tertiary plants from Jancocata, Bolivia. *The John Hopkins University studies in Geology*, 4, 145-202.
- Charrier, R., Muñoz, N., Wyss, A., Flynn, J.J. and Hérail, G. 1994a. Hallazgo de un húmero de *Toxodonte* (Mammalia) en la Formación Chucal (Oligoceno Tardío-Mioceno Inferior) en el Altiplano de Arica, Chile. *Proc. 7<sup>th</sup>. Congreso Geológico Chileno*, Concepción, 430-433.
- Charrier et al., 1994b. Edad y contenido paleoflorístico de la Formación Chucal y condiciones paleoclimáticas para el Oligoceno tardío-Mioceno Inferior en el Altiplano de Arica. *Proc. 7<sup>th</sup>. Congreso Geológico Chileno*, Concepción, 434-437.
- Flynn, J.J., Charrier, R., Hérail, G., Croft, D. and Wyss, A., 1999. The first Cenozoic mammal fauna from the Chilean Altiplano. (Conferencia simposio sobre paleontología de Vertebrados, La Paz, Museo...)
- Gregory-Wodzicki, K.M., McIntosh, W.C. and Velásquez, K., 1998. Climatic and tectonic implications of the late Miocene Jakokkota flora, Bolivian Altiplano. *Journal of South American Earth Sciences*, Vol. 11, N° 6, p. 533-560.