

Anti-inflammation assays

Wahrendorf M.-S., Sporer F., Wink M.

Institut für Pharmazeutische Biologie, Universität Heidelberg Im Neuenheimer Feld 364 69121 Heidelberg (Deutschland)
Email: Marit-Saskia.Wahrendorf@urz.uni-heidelberg.de

Key words: antiinflammation, *Harpagophytum procumbens*, HET-CAM-Assay, Lipoxygenase-Assay

The aim was to establish effective systems to measure the anti-inflammatory potential of ethnopharmacologically used substances. To avoid animal-experiments a modified HET-CAM-Assay (Hens-Egg-Test at the Chorion-Allantoin-Membrane), developed by Theisen and Luepke (0,1) is employed.

The second assay is targeted to a special enzyme system connected to inflammation, which measures lipoxygenase-activity via HPLC (2).

Anti-inflammatory activity of *Harpagophytum procumbens* has been demonstrated in different experiments in rats (3,4,5). In vitro experiments showed that the iridoid glycoside harpagoside inhibits the cyclooxygenase- as well as the lipoxygenasepathway of the arachidonic acid cascade (7). Therefore we have tested extracts of *Harpagophytum procumbens* in the modified HET and the LOX-enzyme-assay.

The plant was extracted three times to give a 60%-, 30%-, and 0%-ethanolic extract, dried and weighed. Its components are analysed and quantified via HPLC (RP-18 column, gradient elution, UV-detection at 280nm, external standart, after (6)).

Het-cam assay modified

Fresh fertile White Legorn eggs (Lohmann Selected Leghorn, LSL) were inoculated on day 9 with the substance supposed to have an antiinflammatory effect. A blank control and a positive (Aspisol) were injected as well. An average of 6 eggs per substance was taken. After further incubation for two hours the eggshell above the cell is removed and the CAM (Chorion-Allantois-Membrane) is exposed.

After dropping 0.3 ml SDS (sodiumdodecylsulfate) on the CAM, time is measured for certain reactions to appear. These were 1.) injection of blood into the vessels; 2.) haemorrhagia and 3.) lysis of the vessels.

A prolongation period of these events is indicative of anti-inflammatory effect.

All extracts showed antiinflammatory activity in low concentrations down to 0.001 mg/egg.

The 30%-ethanolic extract of *Harpagophytum procumbens* turned out to be slightly more potent than the 60%- followed by the 0%-ethanolic extract. (Figure 1)

Lipoxygenase assay

Leucocytes were isolated from human Buffy-Coat blood. After addition of the testsubstances the reaction is started by incubating them at 37°C with arachidonic acid and Calcium-Ionophore. Each sample contains around 106 Leucocytes.

The products, LTB₄, 15-, 12- and 5-HETEs, were measured in an isocratic HPLC-System using a mixture of water and methanol as solvent-system.

The hight of the peaks also gives an indication of quantitative effects. Peak areas of the products reflected the degree of enzyme inhibition.

In this Assay-system the 60%-ethanolic extract seems to be the most potent, followed by the 30 %- and then the 0%-extract. (Figure 2)

Conclusion

Aqueous and hydroalcoholic extracts of *Harpagophytum procumbens* exhibits a clear anti-inflammatory effect in our bioassa. A 30%-ethanolic extract is the most potent in the HET-CAM-assay, whereas lipogxygenase activity is inhibited maximally with the 60%-extract.

Considering costs and reproducibility in comparison to in vivo animal experiments HET-CAM-assay and lipoxygenase-assay are suitable systems for testing antiinflammatory activities of all sorts of subsances.

References

1. LUEPKE N.P. (1992) Toxikologisch-pharmakologische Prüfmöglichkeiten am bebrüteten Hühnerrei, In Schoeffl, Schulte-Herrmann, Tritthart (eds), *Möglichkeiten und Grenzen der Reduktion von Tierversuchen*, Springer-Verlag.
 2. THEISEN N. (1988) *Der modifizierte HET-CAM-Test als ein Modell zur Prüfung von Antiphlogistica*, Inaugural-Dissertation an der Universität Osnabrück.
 3. Nach Professor Dr.Dieter STEINHILBER, Institut für pharmazeutische Chemie, Universität Frankfurt, Marie-Curie-Str.9, 60439 Frankfurt
 4. LANHERS M.C., FLEURENTIN J., MORTIER F. et al (1992) Antiinflammatory and analgesic effects of an aqueous extract of *Harpagophytum procumbens*, *Planta Medica*, 58, 117-123.
 5. BAGHDIKIAN B., LANHERS M.C., FLEURENTIN J. et al (1997) An analytical study, anti-inflammatory and analgesic effects of *Harpagophytum procumbens* and *Harpagophytum zeyheri*, *Planta Medica*, 63, 171-176.
 6. ERDÖS A., FONTAINE R., FRIEHE H. et al (1978) Beitrag zur Pharmakologie und Toxikologie verschiedener Extrakte, sowie des Harpagosids aus *Harpagophytum procumbens* DC, *Planta Medica*, 34, 97-108.
 7. CHRUBASIK S., SPORER F., WINK M. (1996) Zum Harpagosidgehalt verschiedener Trockenextraktpulver aus *Harpagophytum procumbens*, *Forschende Komplementärmedizin*, 3, 6-11
- TIPPLER B., SYROVETS T., LOEW D. et al (1996) *Harpagophytum procumbens* : Wirkung von Extrakten auf die Eicosanoidbiosynthese in Ionophor A23187-stimuliertem menschlichen Vollblut, In Loew D.Rietbrock N. (eds): *Phytopharmaka II, Forschung und klinische Anwendung*, Darmstadt, Steinkopf-Verlag, 95-100.






Figure 1. Relative delay of injection/ haemorrhagia/ lysis by 60, 30, 0 % ethanolic extract of *Harpagophytum procumbens*




Figure 2. Inhibition of lipoxigenase activity by Harpagophytum procumbens in different extracts of human buffy coat blood