


# Excess $^{210}\text{Pb}$ and $^{210}\text{Po}$ in sediment from the Western North Pacific

Koh Harada

Yoko Shibamoto

## Introduction

Particles in most surficial marine sediments are mixed mostly by the biological activities on the sea floors and the mixing is called as



R/V Mirai in November 1998 using with a multiple corer (Figure 2). Immediately after recovery of the corer on the deck, the sediment core sample was sliced by 3 mm thickness for the top ten samples, by 6 mm thickness for the next 10 samples and by 12 mm thickness for the rest. The sliced samples were frozen and transferred to a land laboratory. In the laboratory, the samples were dried and powdered. Water content and porosity were calculated from weight loss after drying.




Figure 2  
Picture of the Multiple Corer in the R/V Mirai cruise.

### *Radiochemical analyses*

Radioactivities of  $^{226}\text{Ra}$  and  $^{210}\text{Pb}$  in the samples were determined by gamma spectrometry with a well-type intrinsic germanium detector. Efficiencies of the detector were calibrated by IAEA sediment standards.

Activity of  $^{210}\text{Po}$  in the samples was determined by alpha spectrometry using  $^{209}\text{Po}$  as a chemical yield monitor (Harada and Tsunogai, 1985). Aliquot amount of the sediment sample was leached in 6 M HCl solution and polonium was collected in hydroxide precipitate

from the leachate. Activities of  $^{209}\text{Po}$  and  $^{210}\text{Po}$  were determined by silicon surface barrier detectors after spontaneous deposition of polonium onto a silver disk from 0.1 M HCl that dissolved the precipitate.

## Results and discussion

### *$^{226}\text{Ra}$ and $^{210}\text{Pb}$ in the sediment at Stn. KNOT*

Vertical distributions of  $^{226}\text{Ra}$  and  $^{210}\text{Pb}$  in the two sediment cores from Stn. KNOT were shown in Figure 3. Concentration of  $^{210}\text{Pb}$  decreased rapidly from surface down to 1 cm depth, and then decreased gradually down to 6 cm depth. Below the depth, the concentration was almost constant, whereas there was a peak at 8 cm depth in the 1998 sample core. Contrastively, concentration of  $^{226}\text{Ra}$ , which is a precursor of  $^{210}\text{Pb}$ , was almost constant in the cores, ranging between 3.5 to 8.0 dpm.g<sup>-1</sup>.

The vertical profiles of  $^{226}\text{Ra}$  and  $^{210}\text{Pb}$  showed that there was excess  $^{210}\text{Pb}$  over its precursor  $^{226}\text{Ra}$  from surface down to 6 cm depth. This excess  $^{210}\text{Pb}$  must be supported by an input from outside of the sediment, which is an input of  $^{210}\text{Pb}$  from the water column by settling particles.

### *Fluxes of $^{210}\text{Pb}$ from the bottom water to the sediment*

Assuming a steady state condition, the  $^{210}\text{Pb}$  flux from the overlying water column to the sediment can be calculated from inventories of  $^{226}\text{Ra}$  and  $^{210}\text{Pb}$  in the sediment from following relation,

$$F_i = \lambda_i I_i \quad (1)$$

where  $\lambda_i$  is a decay constant of nuclide  $i$  (day<sup>-1</sup>) and  $I_i$  is an inventory in the sediment of the nuclide  $i$  (dpm.m<sup>-2</sup>).




Figure 3  
Vertical distributions of  $^{226}\text{Ra}$  and  $^{210}\text{Pb}$   
in the sediment core at Stn. KNOT.

The flux of  $^{210}\text{Pb}$  was calculated as 81.9 and 64.2  $\text{dpm}\cdot\text{m}^{-2}\cdot\text{day}^{-1}$  respectively from the October and November, 1998 profiles. This calculated flux can be compared with the flux observed by a sediment trap experiment. Unfortunately,  $^{210}\text{Pb}$  in the sediment trap samples have not been determined so far, whereas the experiment at Stn. KNOT was carried out by the other group. Judging from previous papers,  $^{210}\text{Pb}$  flux observed by the sediment trap in the North Pacific ranged from 20 to 100  $\text{dpm}\cdot\text{m}^{-2}\cdot\text{day}^{-1}$ . The calculated fluxes from the profiles in the sediment seem to be comparable with the fluxes observed by the sediment traps in the biologically productive area in the North Pacific.


### *$^{210}\text{Po}$ in the sediment*

Using the sediment samples collected in November 1998,  $^{210}\text{Po}$  was also determined, which is a daughter nuclide of  $^{210}\text{Pb}$ . The vertical

profile of  $^{210}\text{Po}$  showed in Figure 4 with  $^{210}\text{Pb}$ . Excess  $^{210}\text{Po}$  over  $^{210}\text{Pb}$  seems to exist in the top two sub-samples and between 7 and 10 cm depth. Generally speaking, the settling particles have high  $^{210}\text{Po}/^{210}\text{Pb}$  ratio from 3 to 10 and this was explained by preferential scavenging of  $^{210}\text{Po}$  from seawater to particulate matter. Although there still exists a possibility of calibration error of the counter because

the excess  $^{210}\text{Po}$  in the deeper samples is not explainable, the excess  $^{210}\text{Po}$  in the core top seems to be caused by higher particulate flux of  $^{210}\text{Po}$  from the water column than of  $^{210}\text{Pb}$ . The  $^{210}\text{Po}$  flux from the water column can be estimated from the inventory of the excess  $^{210}\text{Po}$  in the sediment in the same method as  $^{210}\text{Pb}$  flux mentioned above. The flux was calculated as  $590 \text{ dpm}\cdot\text{m}^{-2}\cdot\text{day}^{-1}$  showing the estimated  $^{210}\text{Po}/^{210}\text{Pb}$  ratio in the settling particle was 7, which is comparable or a little higher than the observed one in the North Pacific.

Figure 4  
Vertical  
distributions of  
 $^{210}\text{Pb}$  and  
 $^{210}\text{Po}$  in the  
sediment core  
collected in  
October.



### *Estimation of bioturbation rate from the vertical profiles of $^{210}\text{Pb}$ and $^{210}\text{Po}$*

As shown above, excess  $^{210}\text{Pb}$  and  $^{210}\text{Po}$  were detected in the sub-surface layers of the sediment core from the western North Pacific. Even though the excesses were supplied from the overlying water column by the settling particles, there must be some mechanisms in which surface sediment particles was mixed with sub-surface one since accumulation rate of pelagic sediment is very slow,  $<1 \text{ cm.kyr}^{-1}$ , compared to the time scale of mean life of the nuclides. The particle mixing by organisms on sea floor, bioturbation, seems to be predominant although the mixing by bottom water current could be occur. If the nuclides is carried to the interior of the sediment only by the bioturbation, the concentration of the nuclides in the sediment can be expressed by the following equation,

$$\frac{dA}{dt} = D_B \frac{d^2A}{dz^2} + S \frac{dA}{dz} - \lambda A \quad 2$$

where  $A$  is a concentration of nuclide  $i$  in the sediment,  $D_B$  is a bioturbation rate constant,  $S$  is a sediment accumulation rate and  $\lambda_i$  is a decay constant of nuclide  $i$ . Since the second term in the right side in the Eq.(2) is negligible, the steady state concentration of the nuclide  $i$  can be expressed as follows;

$$A = C \cdot e^{-\sqrt{\frac{\lambda_i}{D_B}} \cdot z}$$

The concentrations of  $^{210}\text{Pb}$  were plotted against depth semilogarithmically (Figure 5). Judging from the linearity, the sediments were separated into three layers, surface to 1 cm, 1 to 2.5 cm and 3 to 6 cm depth and the bioturbation coefficient in the layers was estimated to be 0.04, 0.4, 0.06  $\text{cm}^2.\text{yr}^{-1}$ , respectively. It is very curious that the second layer has the largest coefficient, however, the reason is still unknown. Using the excess  $^{210}\text{Po}$  profile, the coefficient was also calculated. The estimated value, 1.2  $\text{cm}^2.\text{yr}^{-1}$ , is significantly larger than one obtained from the excess  $^{210}\text{Pb}$ . To clarify the difference between the values from  $^{210}\text{Pb}$  and  $^{210}\text{Po}$ , further study is needed although the age dependent mixing of deep-sea sediments was reported (Smith *et al.*, 1993).




Figure 5  
Semi logarithmic plots of  $^{210}\text{Pb}$  and  $^{210}\text{Po}$   
in the sediment and estimation of  $D_B$ .

## Conclusion

In the sediment in the western North Pacific,

1. excess  $^{210}\text{Pb}$  and excess  $^{210}\text{Po}$  existed down to 6 and 1 cm depth, respectively,
2. particulate fluxes of  $^{210}\text{Pb}$  and  $^{210}\text{Po}$  from the overlying water were estimated as 70 and 590  $\text{dpm.g}^{-1}$ , respectively,
3. bioturbation mixing coefficient in the sediment was estimated to be from 0.004 to 0.4  $\text{cm}^2.\text{yr}^{-1}$  and the highest value was observed in 1 – 2.5 cm layer.



### Acknowledgements

The authors would like to thank scientists, technicians and crew members on R/V Mirai for their help of sample collection.

## Bibliography

- ALLER R. C., 1990 —  
"Bioturbation and manganese cycling in hemipelagic sediments".  
*In* H. Charnock *et al.* (eds) *The Deep Sea Bed: Its Physics, Chemistry and Biology*, Cambridge, University Press: 51-68.
- BERNER R. A., 1980 —  
*Early Diagenesis: A Theoretical Approach*. Princeton, University Press.
- DEMASTER D. J.,  
COCHRAN J. K., 1984 —  
Particle mixing rates in deep-sea sediments determined from excess  $^{210}\text{Pb}$  and  $^{32}\text{Si}$  profiles. *Earth Planet. Sci. Lett.*, 61: 257.
- EMERSON S. R., 1985 —  
"Organic carbon preservation in marine sediments". *In*: Sundquist E. T., Broecker W. S. (eds): *The Carbon Cycle and Atmospheric CO<sub>2</sub>: Natural Variations Archean to Present*, AGU Geophys: 78-86.
- HARADA K., TSUNOGAI S., 1985 —  
A practical method for the simultaneous determination of  $^{234}\text{Th}$ ,  $^{226}\text{Ra}$ ,  $^{210}\text{Pb}$  and  $^{210}\text{Po}$  in seawater. *J. Oceanogr. Soc. Japan*, 41: 98.
- HARADA K., TSUNOGAI S., 1986 —  
Fluxes of  $^{234}\text{Th}$ ,  $^{210}\text{Po}$  and  $^{210}\text{Pb}$  determined by sediment trap experiments in pelagic oceans. *J. Oceanogr. Soc. Japan*, 42: 192.
- NOZAKI Y., COCHRAN J. K.,  
TUREKIAN K. K., KELLER G., 1977 —  
Radiocarbon and  $^{210}\text{Pb}$  distribution in submersible-taken deep-sea cores from project FAMOUS. *Earth Planet. Sci. Lett.*, 34: 167.
- PENG T. H., BROECKER W. S.,  
BERGER W. H., 1979 —  
Rates of benthic mixing in deep-sea sediments as determined by radioactive tracers. *Quat. Res.*, 11: 141.
- SMITH C. R., POPE R. P.,  
DEMASTER D. J., MAGAARD L., 1993 —  
Age-dependent mixing of deep-sea sediments. *Geochim. Cosmochim. Acta*, 57: 1473.