

Temporal and spatial relationship between thick- and thin-skinned deformation in the thrust front of the Malargüe fold and thrust belt, southern Central Andes

Laura Giambiagi ¹, Florencia Bechis ¹, Víctor García ², & Alan Clark ³

¹ IANIGLA-CONICET, Parque San Martín s/n, Mendoza, 5500, Argentina. Email: lgiambia@lab.cricyt.edu.ar; fbechis@lab.cricyt.edu.ar

² Laboratorio de Modelado Geológico (LaMoGe) -Universidad de Buenos Aires, Pabellón II, Ciudad Universitaria, Capital Federal, 1428, Argentina. Email: victorg@gl.fcen.uba.ar

³ Department of Geological Sciences, Queen's University, Kingston, Ontario, Canada, K7L 3N6

KEYWORDS: Southern Central Andes, Malargüe fold and thrust belt, thick- and thin-skinned tectonics

INTRODUCTION

A new kinematic model, which integrates original detailed structural data and new Ar/Ar dating with previous surface and subsurface data and Ar/Ar dating, is proposed for the thrust front of the Malargüe fold and thrust belt (MFTB). The study area, located in the eastern slope of the Cordillera Principal, represents a key area in which to test thin- vs. thick-skinned models, because it affects the Late Triassic – Early Jurassic Atuel half-graben of the Neuquén rift basin (Fig. 1A). The goal of this study is to constrain the timing of uplift of the MFTB by defining of the deformation. We also address the question as to whether tectonic inversion of Mesozoic normal faults occurred first and the resulting basement thrust sheets propagated along a deep detachment into the sedimentary sequence in the foreland, or basement thrusting occurred out-of-sequence after the emplacement of shallow thrust sheets. The presence of pre-existing weaknesses related to rift structures and the occurrence of ductile layers in the cover would support both alternatives.

The MFTB extends from 34° to 36° S and forms part of the Cordillera Principal of the Southern-Central Andes. It comprises: Paleozoic metamorphic and plutonic rocks which constitute the basement of the belt; *Upper* Triassic to *Lower* Jurassic marine and continental rift sequences deposited in the Neuquén back-arc basin; *Upper* Jurassic to Cretaceous platform sequences; and Cenozoic sedimentary and volcanic rocks. This belt developed mainly during the Neogene in response to W-E compression and it has traditionally been interpreted as a thick-skinned orogenic wedge with inverted Jurassic normal faults (Maceda and Figueroa, 1995). Neogene synorogenic continental strata crop out in the Cuchilla de la Tristeza range and eastward, and are represented by three Miocene to Pliocene units separated by angular unconformities: the Agua de la Piedra Fm. (13-10.6 Ma: Baldauf, 1997), Loma Fiera Fm. (10-9.5 Ma: Baldauf, 1997) and Río Diamante Fm. (Pliocene: Combina and Nullo, 1997).




Figure 1. A) Shaded relief map of the Andes showing the study area. B) Simplified structural map of the thrust front of the Malargüe FTB, showing the main structures and the location of Ar-Ar dated pre-tectonic and post-tectonic volcanic and subvolcanic rocks.

SPATIAL RELATIONSHIP BETWEEN THICK- AND THIN-SKINNED STRUCTURES

The thrust front of the MFTB can be divided into two sectors (western and eastern) on the basis of palaeoenvironmental and tectonic relationships (Fig. 1B). Their mutual boundary is defined by the NNW-trending Borbollón-La Manga lineament, related to the Mesozoic rift system. In the *western sector*, outcropping rocks are predominantly Triassic-Jurassic rift sequences. Andean deformation is characterized by the presence of NNW-SSE and NNE-SSW, open and tight folds and low-angle and high-angle faults with both thick- and thin-skinned tectonic styles. Open folds developed in the hanging wall of moderate-high-angle reverse faults and are considered to be formed by inversion of older normal faults. Two structures, the La Manga and El Freno faults, are interpreted to be reactivated rift-related normal faults based on the basis of the highly variable thickness and facies of the rift sequences, the high cut-off angles along these reactivated faults, the presence of antithetic and synthetic faults reactivated in a reverse sense, and synextensional unconformities preserving the original extensional geometry. Seismic lines across the La Manga fault show that it has a convex-up geometry, cutting the basement-cover interface at a high angle and progressively decrease in dip upwards. This geometry strongly implies the inversion of a high-angle pre-existing normal fault by upward propagation of a steep basement fault

into the sedimentary cover. The El Freno fault has been interpreted as a NNE-trending high-angle blind fault, related by Lanés (2005) to the extensional period. The inversion of this fault is marked by a broad anticline in its hanging-wall. Associated with this thick-skinned structure, small-scale anticlines and synclines with angular hinges (kinks and box-folds) deform the Jurassic sequences, and low-angle thrusts formed above shallow detachments, in thin-skinned tectonic style.

The *eastern sector* corresponds to an emergent thrust front system, made up of several NNW-trending thrust sheets involving Cretaceous to Neogene strata in a thin-skinned tectonic style. The oldest sediments involved in the deformation are Cretaceous shales, evaporites and red beds. The Sosneado and Mesón faults uplift the Cuchilla de la Tristeza range and are low-angle thrust rooted into a shallow detachment. Toward the north, both faults split into several thrusts and produce stacking of the Cretaceous sequences (Kozłowski *et al.*, 1989).

We therefore infer that the tectonic evolution of the MFTB involved both thin-skinned tectonics along two shallow detachments within the Jurassic rift sequences (western sector) and Cretaceous strata (eastern sector) and basement involvement along a deeper detachment which accommodates stacking of basement thrust units.

TEMPORAL RELATIONSHIP BETWEEN THICK- AND THIN-SKINNED STRUCTURES

In order to constrain the age of the thick- and thin-skinned deformation, we construct a chronology of deformation based on structural relationships and Ar/Ar dating of tectonic and post-tectonic volcanic and subvolcanic rocks. Twelve samples of andesite and basalt were analysed by laser-induced $^{40}\text{Ar}/^{39}\text{Ar}$ step-heating procedure on hornblendes and whole-rocks.

Inversion of the Atuel half-graben (15 – 14 Ma): Contractional reactivation of the Atuel half-graben began with rigid displacement of the wedge of rift deposits and the underlying crystalline basement rocks along the La Manga fault. Fault displacement was dissipated in the cover units by folding. Porphyry dikes, assumed to be post-tectonically emplaced in the Cerro Chivato area by Baldauf (1997), indicate that the anticline associated with the first movement of the La Manga fault formed before 14 Ma. A maximum age for this deformation is given by the age of pre-tectonic subvolcanic rocks, cropping out in the Las Bardas creep, dated as 14.87 Ma (Fig. 1B).

Breakthrough of the La Manga fault into the sedimentary cover (13 – 11 Ma): After the partial inversion of the half-graben, faults emanating from the master fault broke through the entire sedimentary section and reached the surface. The time of breakthrough is well constrained by the age of the post-tectonic volcanics and by the angular unconformities between the synorogenic strata. In the thick-skinned area, deformation was accommodated by movement along the La Manga fault prior to 10.84 Ma, the age of the Cerro Tordilla post-tectonic volcanic rocks. In the thin-skinned area, cross-cutting relationships, together with emplacement ages, indicate that deformation and uplift of the Cuchilla de la Tristeza range along the Mesón and Sosneado thrusts must have occurred prior to 10.6 Ma, the age of the Laguna Amarga stock (10.56 ± 0.04 Ma - Baldauf, 1997) and that of the angular unconformity separating the Agua de la Piedra and Loma Fiera Formations (~10.6 Ma -

Internal deformation of the half-graben (11 - 9 Ma): The age of movement along the El Freno fault system, related at depth to the inversion of the pre-existing El Freno normal fault, is determined by the ages of pre-tectonic volcanic rocks (11.16 ± 0.28 Ma) and post-tectonic volcanics (9.11 ± 0.24 Ma). This indicates that the internal deformation of the half-graben occurred after the inversion of the La Manga normal fault.

CONCLUSIONS

The question whether shortening in the basement occurred first and was transmitted to the cover, or the cover detached first and basement thrusting occurred afterwards, has been addressed by detailed field structural data combined with Ar/Ar dating of pre- and post-tectonic volcanics and subvolcanic rocks in the MFTB. Comparison of the timing of deformation in the thick- and thin-skinned deformational areas strongly supports the hypothesis that the reactivation of normal faults was coeval with the activation of shallow detachments and low-angle thrusting in the thrust front of the MFTB. Low-angle thrusting interacted with high-angle faulting related to reversal of basement normal faults inherited from the extensional history of the foreland, indicating a deformation mode characterized by superimposed shallow and deep detachment tectonics. Within the thrust front of the MFTB, detachments occur at several stratigraphic horizons: a deep basement detachment related to the basement-involved thrusting, and shallow detachments located within the Jurassic and Cretaceous sequences. We propose that these detachments were active during the complex deformation of the thrust front, between 14 and 9 Ma.

REFERENCES

- Baldauf, P., 1997. Timing of the uplift of the Cordillera Principal, Mendoza Province, Argentina. M. S. Thesis, George Washington University, 356 p.
- Baldauf, P., Stephens, G. C., Nullo, F. E. and Combina, A. M., 1997. Tertiary uplift, magmatism and sedimentation of the Andes, Southern Mendoza province, Argentina. Geological Society of America, Annual Meeting with program 29: A48.
- Combina, A. M., and Nullo, F., 1997. Consideraciones tectosedimentarias sobre la Formación Río Diamante, Cordillera de los Andes, Argentina. Cuadernos de Geología Ibérica, 22: 305-320, Madrid.
- Kozlowski, E., Cruz, C., Condat, P., and Manceda, R., 1989. Interpretación del fallamiento de bajo ángulo en los sedimentos Cretácicos del río Diamante, provincia de Mendoza. I° Congreso Nacional de Exploración de Hidrocarburos, Actas I: 675-687.
- Lanés, S., 2005. Late Triassic to Early Jurassic sedimentation in northern Neuquén Basin, Argentina: Tectonosedimentary evolution of the first transgression. *Geologica Acta* (in press).
- Manceda, R. and Figueroa, D., 1995. Inversion of the Mesozoic Neuquén rift in the Malargüe fold-thrust belt, Mendoza, Argentina. In: A. J. Tankard, R. Suárez y H.J. Welsink (Eds): *Petroleum Basins of South America*. American Association of Petroleum Geologists, Memoir 62: 369-382.