

The magmatism in the easternmost sector of the Calama-Olocapato-El Toro transversal fault system in the Central Andes at 24° S: Geotectonic significance

M. Matteini⁵, A. Gioncada³, R. Mazzuoli³, V. Acocella¹, A. Dini², H. Guillou⁴, R. Omarini⁵, A. Uttini⁶, L. Vezzoli⁶, & N. Hauser⁵

1 Dip. Scienze Geologiche, Università Roma Tre, Italy

2 Istituto di Geoscienze e Georisorse CNR, Pisa, Italy

3 Dip. Scienze della Terra, Univ. Degli Studi di Pisa, Italy

4 Lab. des Sciences du Climat et de l'Environnement, CNRS, Gif-sur-Yvette, France

5 Facultad de Ciencias naturales, Universidad Nacional de Salta, Argentina

6 Dip. Scienze Chimiche e Ambientali, Univ. degli Studi dell'Insubria, Como, Italy

The easternmost sector of Calama-Olocapato-El Toro (COT) transversal fault system in the Central Andes at 24° S is characterised by the presence of a Miocene magmatic complex (Fig. 1) constituted by intrusive (Hongn et al. 2002) and volcanic rocks (Krallmann 1994, Hauser et al., 2005). Intrusive rocks consist of a laccolith-like intrusion having an U/Pb age on zircon of 14.4 Ma (Hongn et al. 2002) whereas new K/Ar age (this work) determinations indicate that the volcanic products emplaced in a span of time from 12 to 6 Ma. The lavas are located along fractures with ~E-W and ~NW-SE strike (Hauser et al. 2005) and the eruptive centres are distributed in an area of about 40 km². These Miocene igneous rocks are of particular scientific interest for their position at about 600 km from the Pacific trench and for the complex tectonic evolution of the area in which they emplaced. In fact in the last 10 Ma this sector of Central Andes was characterised by crustal thickening associated to extensive shortening. In this note, we report new geological and petrological data on the Almagro volcanic rocks (AVR from here) and Las Burras intrusion (LBI from here) that can allow us to: a) characterize the magmatism in trans-arc position very far from the trench b) to explain the mechanisms of pluton emplacement in a general compressive tectonic environment and 3) to define the role of the Calama-Olocapato-El Toro lineament (COT) in ascent of magmas.






Figure 1: Geological sketch map of Central Andes at about 24° S of latitude.

The volcanic rocks have a composition ranging from basaltic andesites to dacites at the boundary with rhyolites. The basaltic andesites contain as dominant mineral phases olivine, zoned plagioclase, clino- and orthopyroxenes whereas the andesites and dacites contain plagioclase, abundant amphibole, clino- and orthopyroxene with scarce olivine. The presence of stable amphibole in these intrusive



N-S faults. The N-S dextral faults seem to have controlled the rise of the monzogranite, forming a NE-SW trending releasing bend. The successive extension found along the NW-SE and the N-S systems can be interpreted as induced by the rise of magma along the COT.

The whole of the geochemical data suggest that LBI magmas could be generated by melting of a lithospheric mantle with relatively low Sr and high Nd isotopic ratios. The evolution processes which affected the LBI magmas (SiO₂ ranging between 54 and 68%) possibly occurred in a low radiogenic continental crust. The volcanics could be the result of a mixing process between a mantle-derived magma and crustal melts deriving by partial melting of Brazilian crust. The magma generating processes could be linked to an increase of isotherms due to the upflow of asthenosphere after the Quechua compressive phase and/or to a depressuring process due to the transtensional movements of the COT. In this case the COT played an important role either for the ascent of magmas.

References

- Kay, S.M., Mpodozis, C. and Coira, B., 2000. Neogene magmatism, tectonism, and mineral deposits of the Central Andes (22° to 33°S latitude). In: Skinner, B.J., Editor, , 1999. *Geology and Ore Deposits of Central Andes*. Soc. Econ. Geol., Spec. Publ. vol. 7, pp. 27–59.
- Krallman, A., 1994. Petrographische und geochemische Untersuchungen an jungen basischen Vulkaniten im Bereich des Calama-Olacapato-El Toro lineamentes östlich der vulkanischen Kette, NW Argentinien. *Clausthaler Geowissenschaftliche Dissertationen*, Heft 45; 1-150.
- Hauser, N., Matteini, M., Omarini, R., Mazzuoli, R., Vezzoli, A., Acocella, V., Uttini, A., Dini, A., Gioncada, A., 2005. Aligned extrusive andesitic domes in the southern sector of the Miocene Tastil-Las Burras volcanic complex, Eastern Cordillera, Salta, Argentina: new evidences for NNW-SSE extensional tectonics. XVI Congreso Geológico Argentino, La Plata 2005.
- Hongn, F.D., Tubía, J. M., Aranguren y Mon, R., 2002. La Monzodiorita Las Burras: un plutón mioceno en el batolito de Tastil, Cordillera Oriental, Argentina. XV Congreso Geológico Argentino, Acta 1, 128-133.
- Matteini, M., Mazzuoli, R., Omarini, R., Cas, R. and Maas, R., 2002. The geochemical variations of the upper Cenozoic volcanism along the Calama–Olacapato–El Toro transversal fault system in central Andes (~24°S): petrogenetic and geodynamic implications. *Tectonophysics*, 345: 211-227.
- Ort, M., Coira, B. and Mazzoni, M., 1996. Generation of a crust–mantle magma mixture: magma sources and contamination at Cerro Panizos, central Andes. *Contrib. Mineral. Petrol.* **123**, pp. 308–322