

Effects on invertebrates of traditional fishing by river poisoning in French Guiana

Impact sur les invertébrés de pêches traditionnelles par empoisonnement des rivières en Guyane française

O. Fossati⁽¹⁾, S. Mérigoux⁽²⁾, V. Horeau⁽³⁾ & M. Jégu⁽⁴⁾

(1) Correspondence: Odile Fossati, IRD-Université Lyon 1, Ecologie des hydrosystèmes fluviaux, 43 bd du 11 novembre 1918, 69622 Villeurbanne-Cedex – France <j.o.fossati@wanadoo.fr>

(2) Université Lyon 1 – France <sylvie.merigoux@univ-lyon1.fr>

(3) Hydreco, Kourou - Guyane française <hydreco-labops@wanadoo.fr>

(4) IRD - MNHN Paris - France <jegu@mnhn.fr>

Abstract

In the upper Maroni River (French Guiana), Wayana Amerindians fish by poisoning the rivers with toxins, mainly rotenone, extracted from lianas. We studied the impacts of this fishing practice on benthic invertebrate communities associated with Podostemaceae meadows in October 2000. Ephemeroptera or Diptera Simuliidae were dominant. The other Diptera, mainly Chironomidae, represented the third group of invertebrates, followed by Lepidoptera and Trichoptera. Annelida, mainly Oligochaeta, and Acarina were of little abundance, and Odonata and Coleoptera scarce. Densities were generally lower after fishing but no statistically significant effect was found. The toxic used by the Amerindians had no significant effect on invertebrate communities and the indirect impact for fishes is probably negligible.

Key-words

neotropical, rotenone, stream, toxicity

Résumé

Les indiens Wayanas du Haut Maroni (Guyane française) pêchent en empoisonnant les rivières avec des toxines (principalement de la roténone) extraites de lianes. Nous avons étudié l'effet de ces pratiques sur les peuplements invertébrés des herbiers à Podostemaceae en octobre 2000. Les Ephéméroptères ou les Diptères Simuliidae dominaient les peuplements. Les autres Diptères, principalement les Chironomidae, représentaient le troisième groupe d'invertébrés, suivis par les Lépidoptères et les Trichoptères. Les Annélides, principalement des Oligochètes, et les Acariens étaient généralement peu nombreux. Les Odonates et les Coléoptères étaient rares. Les densités étaient généralement plus faibles après les pêches, sans que les différences soient statistiquement significatives. Les toxiques utilisés par les Amérindiens ne semblent donc pas avoir d'effet significatif sur les peuplements invertébrés. L'impact indirect sur les poissons reste aussi probablement négligeable.

Key-words

cours d'eau, néotropical, roténone, toxicité

Introduction

In the Upper Maroni River (French Guiana), Wayana Amerindians fish by poisoning the rivers with toxins extracted from lianas such as *Derris elliptica* (Grenand & Moretti, 1982, Pagezy & Jégu, 2002). The active component of the poison, rotenone, is a well known fish toxicant (Schnick, 1974).

Growing on rocky substrates in fast flowing waters, Podostemaceae meadows form a complex

biotope in clear or black water streams draining the Precambrien shields in the Guianas, Venezuela and Brazil. A rich invertebrate fauna is associated with these meadows, composed mainly of insects larvae (Tavares *et al.*, 1998, Odinetz-collart *et al.*, 2001). Patrimonial fishes like the large phytophagous Serrasalminae (Teleostei : Characidae) feed on these invertebrate communities when juveniles (Py-Daniel & Jégu, 1996, Pagezy & Jégu, 2002).

The impact of this traditional fishing practice on benthic invertebrate communities was studied in October 2000, as part of an IRD (Institut de Recherche pour le Développement) and MNHN (National Museum of Natural History of Paris)'s project. The aim of this project was to describe the invertebrate communities in the Upper Maroni River and evaluate the impact of fish poisoning with rotenone on benthic invertebrate communities (Jégu *et al.*, 2003).

Study area and methods

Our study area was the Upper Maroni River Basin (Figure 1). Invertebrate communities were described before fishing in five sites near Antecume Pata (one on the Marouini River - M - and four on the Litany River - L1-L4) and in two areas in Saut Pierkourou, on the Upper Tampok (T1, T2). One large commercial fishing was observed on the Upper Tampok, where no fishing had been practiced for five years. Four familial fishing operations were studied near Antecume Pata, where fishing occurs each year, as soon as the water level is low.

Figure 1.

Mourera fluviatilis is the dominant Podostemaceae in the Upper Maroni River (Pagezy & Jégu, 2002). *Mourera fluviatilis* fragments (3 to 20 g each, 3 fragments in the Marouini and Litany Rivers sites, 5 fragments in the Tampok River sites, due to different operators) were taken in plants in a homogeneous meadow, in a rapid shallow area (10 - 30 cm depth), just before and 24 h after the fishing operations. The organic matter and invertebrates associated with the plant fragments were filtered (250 μ m mesh-size) and preserved with formaldehyde in the field. Organisms were determined and counted under a binocular microscope in the laboratory. Densities were reported in relation to fresh weight of plants (individuals.g⁻¹) and compared with t-tests ($p < 0.05$).

Results

Ephemeroptera (Leptohyphidae, Baetidae and Leptophlebiidae) were dominant in the Litani River (up to 70 % of the total invertebrates number in L4) and in one of the Tampok rapids (T2, 69 % - Table 1). However, Diptera Simuliidae were dominant in the Marouini River and in Tampok-1 site (90 and 62 % respectively), but were less abundant in the other rapids (0 - 17 %). The other Diptera, mainly Chironomidae, represented the third group of invertebrates (1 - 34 %). Lepidoptera and Trichoptera larvae were regular in the samples, although less numerous (roughly 1 - 11 %). Annelida, mainly Oligochaeta, and Acarina were of little abundance (less than 5 %). Odonata and Coleoptera were always scarce (less than 0.5 %).

Table 1.

Except for Litany-4, densities of invertebrates were lower after the fishing campaigns with rotenone (densities reduced by half in the Tampok 1) but not statistically (Figure 2). The community analysis showed a weak effect of the rotenone in the Tampok River. Some taxa, such as Philopotamidae, Trichoptera A, Hydropsychidae were not found after fishing.

Discussion - Conclusion

Ephemeroptera were abundant in our samples, although Diptera are generally dominant in Amazonian river invertebrate communities (Tavares *et al.* 1996, Odinetz-Collart *et al.* 2001). The large larvae of Lepidoptera and Trichoptera, although of little abundance, represented a major part of the biomass. In rapids of the Sinnamary River (French Guiana), aquatic invertebrates, mainly Diptera and Ephemeroptera, represented 71 % of the diet of the fishes (Horeau *et al.* 1998).

Effects of rotenone on fish communities have been well studied (Schnick, 1974), but references for invertebrates are scarce, and usually focus on standing waters. Invertebrates are generally less sensitive than fishes, except planctonic crustacea (*in* Schnick, 1974). In the laboratory, rotenone is more toxic to Daphnia (Cladocera) than to fishes (Rach *et al.*, 1988). In temperate areas, rotenone is known to have important effects on stream invertebrate communities: e.g., total loss of invertebrates along several kilometers (Dexter, 1965), loss of Ephemeroptera and Trichoptera (Binns, 1967), reductions of 34 to 100 % of individuals, depending on the invertebrate Order (Little, 1966, Helfrich, 1978). However, This effect was often assessed as weak (Morrison, 1977). It is generally accepted that the observed reductions are not critical to population survival because of possible re-colonization (*in* Schnick, 1974). In tropical areas (Papua-New-Guinea), rotenone was used to kill fishes in two streams and significantly reduced three invertebrates populations (a Baetidae, a Hydropsychidae, a Dixidae) but did not affect abundance and richness of the communities (Dudgeon 1990).

In our study, the toxics used by the Amerindians had no significant effect on invertebrate communities. The difference between the streams may be related to different fishing practices by the Indians □ they use probably more toxic in a remote river than near their village. A rapid recolonization through drift may compensate the effect of the toxic when only one reach is treated, like during traditional fishing operations. A higher level of identification could reveal sensitive species whereas a whole group is not sensitive. However, the right taxonomic level for this seems to be the species level (see Dudgeon 1990) and this level of identification is not yet possible for the Maroni fauna. In fact, this publication is the first one on invertebrates of the Upper Maroni River.

Although rotenone is toxic to invertebrates and fishes do feed on aquatic invertebrates, the indirect impact on fish communities, through effects on invertebrate communities, is probably negligible compared to direct mortality. However, our limited sampling does not allow strong inference. A more intense sampling effort, on both spatial and temporal scales, would lead to better statistical power to assess rotenone effects on invertebrates. .

Acknowledgements

This project was supported by the French Ministry of Environment and Sustainable Development (convention IRD / ECOFOR 2498). We thank the Wayana Amerindians for their welcome and the participants to the field assignment "Nivrée 2000", especially Bénéad Faujour (ENSAT Toulouse), Yves Fermon (MNHN Paris), and Hélène Pagezy (CNRS Paris). We also thank two anonymous reviewers for

their helpful comments.

Bibliography

- Binns, N.A., 1967. Effects of rotenone treatments on the fauna of the Green River, Wyoming. - *Wyoming Game Fish Com., Fish. Res. Bull.*, **1** : 1-114.
- Dexter, W.D., 1965. Some effects of rotenone treatment on the fauna of the Green River. - *Wyoming. West. Assoc. State Game Fish. Com.*, **45** : 193-197.
- Dudgeon, D., 1990. Benthic community structure and the effects of rotenone piscicide on invertebrate drift and standing stocks in two Papua New Guinea streams. - *Arch. Hydrobiol.*, **119** : 35-53.
- Grenand, P. & Moretti, C., 1982. Les nivrées ou plantes ichtyotoxiques de la Guyane française. - *J. Ethnopharmacology*, **6** : 139-160.
- Helfrich, L.A., 1978. Effects of rotenone on macrobenthic invertebrates of a Pennsylvania stream. - *Ann. Conf. Southeast Assoc. Fish Wildl. Agencies*, **32** : 401 - 408
- Horeau, V., Cerdan, P., Champeau, A. & Richard, S., 1998. Importance of aquatic invertebrates in the diet of rapids-dwelling fish in the Sinnamary River, French Guiana. - *J. tropic. Ecol.*, **14** : 851-864.
- Jégu, M., Carrière, S., Faujour, B., Fermon, Y., Fish-Müller, S., Fossati, O., Keith, P., Le Bail, P.Y., Meunier, F., Pagezy, H., Salaün, P. & Weber, C., 2003. *Typologie des pêches à la nivrée dans le Haut maroni et évaluation de leurs effets sur la faune aquatique en vue de la gestion de la ressource*. Report, Programme Ecosystème Tropicaux – French Ministry MEDD, 164 pp.
- Little, J.D., 1966. Reclamation of Pine Creek, Tennessee. - *Proc. Southeast. Assoc. Game Fish Com.* **19** (1965) : 302-315.
- Morrison, B.R.S. 1977. The effects of rotenone on the invertebrate fauna of three hill streams in Scotland. - *Fish. Manage.*, **8** : 128-139.
- Odinetz-Collart, O., Tavares, A.S. & Enriconi, A., 2001. Response of Podostomaceae aquatic biocenosis to environmental perturbations in central Amazonian waterfalls. - *Verh. internat. Verein. Limnol.*, **27** : 4063-4068.
- Pagezy, H. & Jégu, M., 2002. Valeur patrimoniale de Serrasalminae herbivores du Haut Maroni (Guyane française) : approche biologique et socioculturelle en pays wayana. - *Bull. fr. Pêche Piscic.*, **364** : 49-69.
- Py-Daniel, V. & Jégu, M., 1996. Peixes (Serrasalminidae) predadores dos imaturos de *Thrysopelema guianense* (Wise, 1911) (Diptera, Culicomorpha, Simuliidae), vetor da filária *Onchocerca volvulus* (Leuckart, 1893). - *Entomol. Vector.*, **3** : 27-29.
- Rach, J.J., Bills, T.D. & Marking, L.L., 1988. Acute and chronic toxicity of rotenone to *Daphnia magna*. - *U.S. Fish. Wildl. Serv. Invest. Fish Control*, **92** : 1-5.
- Schnick, R.A., 1974. A review of the literature on the use of rotenone in fisheries. - *U.S. Fish. Wildl. Serv. Lit. Rev.*, **74-15** : 130 p.
- Tavares, A.S., Enriconi, A. & Odinetz-Collart, O., 1998. Insect communities associated with aquatic macrophytes Podostemaceae in amazonian waterfalls. - *Verh. internat. Verein. Limnol.*, **26** : 2158-2161.

Table and figures

Table 1. Relative abundances (%) of invertebrates collected at the seven study sites (M = Marouini, L = Litany, T = Tampok).

Figure 1. Study area, with river indicated.

Figure 2. Overall densities of invertebrates (individuals.g of leaves⁻¹) on *Mourera fluviatilis* leaves before and after the fishing campaigns (mean ± standard deviation - L = Litany, 3 fragments of leaves ; T = Tampok, 5 fragments of leaves).