

The effect of salinity on sandy soils physical characteristics and rice root system development: a case study from Northeast Thailand

Hartmann, C.¹; H. Hao²; J.L. Maeght³; A.D. Noble⁴; A. Yuvaniyama³ and A. Polthanee⁵

Keywords: Sandy soils, saline patches, root development, soil porosity

Abstract

During the last several decades, saline patches have appeared in paddy fields located on sandy soils of Northeast Thailand this being due to capillary rise of underground saline water associated with elevated watertables. This soil salinisation decreases rice production and is a major threat for future agricultural production as these saline patches can spread making paddy fields highly saline. Under extreme saline conditions these saline patches are colonized by halophytes. Since the physical characteristics of sandy soils do not seem to be affected by salinity, research has focused predominantly on changes in chemical characteristics associated with soil degradation. The objective of this study is to determine whether i) salt concentration of the saline patches induces physical degradation in sandy soils and 2) if farmer strategies associated with organic matter spreading improves soil characteristics and root development of rice. In a severely affected area, we selected two neighbouring farm holds that had contrasting organic matter (OM) management strategies implemented during the last decade: OM was never used (OM⁰), differed from the farmer who routinely applied to saline patches (OM⁺). In each of the farmers holding, 4 fields were selected, each field containing one saline patch. In each plot, 3 areas were identified and were the focus of the study: P the middle of the saline patch where rice is unable to grow (bare soil), S at the edge of the saline patch where rice development is restricted and C, the surrounding area not affected by salinity where rice development and yield are considered as adequate. Saline patches have significantly reduced porosity (0-20cm). Reduced porosity is correlated with a drastic reduction in root development both frequency and depth of proliferation. Farmers' strategies associated with OM spreading increased soil porosity and improved root development regardless of any chemical improvement in the soil. Consequently, physical degradation cannot be neglected to characterize saline patches and rehabilitation techniques need to address both chemical and physical issues. The reason for poor plant development within saline patches is still unclear and requires further research. The study demonstrated that despite their low amount of clay, sandy soils are far from 'inert' under saline conditions.

¹ IRD – UR 176 'SOLUTIONS' – UMR 7618 'BIOEMCO' 46 rue d'Ulm – 75230 Paris Cedex 05 – France.

² Institute of Soil, Chinese Academy of Science, Nanjing, PR China.

³ Land Department Development, Office of Science for Land Development/IRD, Phaholyothin Road, Chatuchak, Bangkok 10900, Thailand.

⁴ International Water Management Institute (IWMI), c/o WorldFish Center Jalan Batu Maung, Batu Maung 11960 Bayan Lepas, Penang, Malaysia.

⁶ Associated Professor, Department of Agronomy, Faculty of Agriculture, Khon Kaen University, Thailand.

Management of Tropical Sandy Soils for Sustainable Agriculture



A holistic approach for sustainable development of problem soils in the tropics

27th November - 2nd December 2005
Khon Kaen, Thailand



Proceedings

Organizing Committee:

Christian Hartmann (IRD/LDD)	Chairman
Narong Chinabut (LDD)	co-Chairman
Andrew Noble (IWMI)	Secretary
Yuji Niino (FAO)	Treasurer
Taweesak Vearasilp (LDD)	co-Secretary
Anan Polthanee (KKU)	co-Secretary
Roland Poss (IRD)	co-Secretary

Scientific committee:

Andriantsona	(CIAT, Kenya)
Richard Bell	(Murdoch University, Australia)
Sue Berthelsen	(CSIRO, Australia)
Eric Blanchart	(IRD, France)
Ary Bruand	(ISTRO, France)
John Caldwell	(JIRCAS, Thailand)
Suraphol Chandrapatya	(IWMI, Thailand)
Hari Eswaran	(USDA, USA)
Martin Fey	(Stellenbosh University, South Africa)
Alfred Hartemink	(ISRIC, The Netherlands)
Christian Hartmann	(IRD, Thailand)
Irb Kheoruenromne	(Kasetsart University, Thailand)
Phil Moody	(Dep. Natural Resources and Mines, Australia)
Paul Nelson	(James Cook University, Australia)
Andrew Noble	(IWMI, Thailand)
William Payne	(Texas A&M University, USA)
Roland Poss	(IRD, France)
Robert Simmons	(IWMI, India)
Christian Valentin	(IRD, Laos)
Bernard Vanlauwe	(CIAT-TSBF, Kenya)
Hidenori Wada	(Japan)
Toshiyuki Wakatuki	(Kinki University, Japan)
Wanpen Wiriyakitnateekul	(LDD, Thailand)

NOTICE OF COPYRIGHT

All rights reserved. Reproduction and dissemination of material in this information product for educational or other non-commercial purposes are authorized without any prior written permission from the copyright holders provided the source is fully acknowledged. Reproduction of material in this information product for sale or other commercial purposes is prohibited without written permission of the copyright holders. Applications for such permission should be addressed to the Organizing Committee of the first Symposium on the Management of Tropical Sandy Soils through the Land Management Officer, FAO Regional Office for Asia and the Pacific, Maliwan Mansion, 39 Phra Atit Road, Bangkok 10200, Thailand.

ISBN 978-974-7946-96-3

For copies write to: Yuji Niino
Land Management Officer
FAO Regional Office for Asia and the Pacific
Maliwan Mansion, 39 Phra Atit Road
Bangkok 10200
THAILAND
Tel: (+66) 2 697 4000
Fax: (+66) 2 697 4445
E-mail: Yuji.Niino@fao.org



Under the auspices of:
International Union of Soil Science (IUSS)

organized by:
L'Institut de Recherche pour le Développement (IRD, France)
Land Development Department (LDD, Thailand)

co-organized by:
International Water Management Institute (IWMI)
Food and Agriculture Organization of the United Nations (FAO)
University of Khon Kaen, Faculty of Agriculture (KKU)

supported by:
CSIRO Land and Water, Australia
UNCCD
ISRIC

