

# INTRACONTINENTAL SEISMICITY AND NEOGENE DEFORMATION OF THE ANDEAN FOREARC IN THE REGION OF ARICA (18.5°S-19.5°S)

*Claire DAVID (1), Joseph MARTINOD (1, 2), Diana COMTE (1), Gérard HÉRAIL (2), Henri HAESSLER (3),*

(1) Universidad de Chile, Casilla 2777, Santiago, Chile (dcomte@dgf.uchile.cl, d.claire@dgf.uchile.cl)

(2) IRD, LMTG, 38 rue des 36 Ponts, 31400 Toulouse, France (jmartino@cec.uchile.cl, gherail@paris.ird.fr).

(3) IPGS, 5 rue Rene Descartes, 67084 Strasbourg Cedex France, (henri@sismo.u-strasbg.fr )

**KEY WORDS** : Forearc, Shortening, Tilting, Intraplate Seismicity, Altiplano.

## INTRODUCTION

The superficial crustal seismic activity in Chile is remarkably moderate. Almost all the Chilean earthquakes are located on the subduction plane, and most of the intracontinental seismicity that results from the growth of the Cordillera occurs on the Eastern side of the Andes, in Bolivia and Argentina. Geological data, indeed, in agreement with the seismological measurements, show that the Chilean forearc suffered very moderate deformations in the Neogene [e.g. Lamb et al., 1997; García et al., 1997], in spite most of the surrection of the Altiplano probably occurred at that time [e.g. Gregory-Wodzicki, 2000]. The only region in Chile where significant crustal seismicity has been reported in the forearc is the northernmost part of the country, near Arica, where a permanent seismic network evidences an intracontinental forearc seismicity that vanishes south of 19.7°S. In this work, we describe the main characteristics of this seismicity, the associated tectonics of the area, and its possible relations with the growth of the Altiplano.


## GEOLOGICAL DATA

The forearc and volcanic arc of the Andes near Arica can be divided in four main longitudinal morphological units, which are from west to east: (1) The Coastal Cordillera composed by Jurassic and Cretaceous volcanic rocks, (2) the Central Depression filled by Cenozoic sediments, (3) the Precordillera mainly composed by Miocene ignimbrites and detritic series overlying Mesozoic rocks, and (4) the Western Andean Cordillera behind which the present-day volcanic arc is developed. The Precordillera corresponds to a regular slope that joins the Central Depression whose mean elevation is 1000 m, with the 4000 m high Western Cordillera.


Active folds, thrusts and faults are scarce in the forearc of the Arica zone [e.g. Muñoz and Charrier, 1996; Lamb et al., 1997]. The Quaternary activity of the East-vergent thrust-faults and related growth anticlines that separate the Western Cordillera from the Altiplano (Lauca Basin) [Riquelme, 1998] is not well documented, although near Guallatire, the Quaternary sediment of the Lauca river is deformed. The so called “west vergent Thrust System” [Muñoz and Charrier, 1996] that emerges in the Putre-Belen area West of the Western



first for the profile P2 forearc events, then for those of profile P3, and finally for all the forearc seismicity. The three calculated stress tensors show the same compressive regime, with a maximum stress axis  $\sigma_1$  oriented N75E° and dipping 30° to the West. This result suggests that the intracontinental west-dipping seismic corresponds to an east-vergent reverse fault (Fig. 3).



**Figure 2** : Cross-sections showing the seismic events localized on profiles P2 and P3 (see Fig. 1 for the location of the cross sections). Focal mechanism solutions are projected on the northern hemisphere.




**Figure3** : Stress Tensor obtained from the intraplate forearc seismic events.

## DISCUSSION AND CONCLUSIONS

Forearc earthquakes are located between 2 km and 59 km depth with a large concentration of earthquakes at the largest depths, between 50 km and 59 km. The intracontinental seismicity belongs therefore both to the upper lithospheric mantle and to the crust of the South America plate, the depth of the Moho varying from 40 km under the coastal Cordillera to 60 km under the volcanic arc. The contrast between the seismic behavior of the forearc on the one hand, and that of the Western Cordillera and Altiplano on the other hand, is remarkable. The western Altiplano does not show any seismicity at depths larger than 10 km, while in the forearc earthquakes are observed from the surface up to 60 km depths. This different behavior probably results from the difference between the rheological profile of the forearc, that remains cold, rigid and brittle at large depths, and that of the Altiplano and Western Cordillera, that is characterized by a ductile low viscosity middle and lower crust. The west-dipping forearc seismic plan may correspond to the thermal and rheological boundary between the rigid forearc block and the soft crust of the magmatic arc (David, 2002).

This seismic plan emerges in the Western Cordillera (Fig. 4). Its position suggests that the superficial thrust-faults that have been reported on both sides of the Western Cordillera may be associated with this deep east-vergent reverse fault. The geometry of the superficial Western Cordillera tectonic structures does not constrain the vergence of the deep related thrust-fault. The Neogene thrust-faults of the Western Cordillera are minor accidents that do not accommodate much shortening, and they cannot have accommodated much of the surrection of the Altiplano. This surrection has essentially been accommodated west of the volcanic arc, by the weak tilting of the entire chilean forearc, and in the Altiplano by the flow of weak ductile crust coming from the

East and resulting from the underthrusting of the Brazilian Craton [Lamb et al., 1997]. Then, we propose that the deep thrust-fault that results in the shortening of the Western Cordillera in the Arica region dips to the West below the forearc. This fault may be expected to result from the westward flow of lower crust below the Altiplano. The surrection of the Altiplano resulting from this deep flow, in turn, may have been accommodated by the tilting of the forearc above this deep east-vergent thrust fault (Fig. 4).



**Figure 4** : Diagram illustrating the Neogene to recent kinematics of the western Margin of the Central Andes.

**ACKNOWLEDGEMENTS** This work was partially founded by FONDECYT 1020104.

## REFERENCES

- Comte, D., Dorbath L., Pardo M., Monfret T., Haessler H., L. Rivera, Michel Frogneux, B. Glass and C. Meneses, *Geophys. Res. Lett.*, 26, 1965-1968, 1999.
- Comte, D., H. Haessler, L. Dorbath, C. David, B. Glass, E. Correa, C. Meneses, J. Vergara, J. Tapia, I. Balmaceda and G. Hérail., *IX Congr. Geol. Chileno, Jornadas de Geofísica*, Puerto Varas, pp 409-412, 2000.
- David, C.; Comte, D.; Dorbath, L.; Haessler, H.; Frogneux, M.; Glass, B.; Balmaceda, I.; Correa, E.; Meneses, C.; Cruz, A.; Ruz, L., *European Geophysical Society, XXVI General Assembly*, Nice, France, 2001.
- David, C., Tesis de Magíster en Ciencias, Mención Geofísica, Universidad de Chile, 46 p., 2002.
- García, M., Hérail G., and Charrier R., *Actas VIII congreso geológico chileno*, 60-64, 1997.
- García, M., PhD thesis, Université de Grenoble, 2001.
- Gregory-Wodzicki, K., *Geol. Soc. Am. Bull.*, 112, 1091-1105, 2000.
- Haessler, H., Comte, D., L. Dorbath, C. David, B. Glass, E. Correa, C. Meneses, I. Balmaceda, A. Tapia, J. Vergara and G. Herail. *American Geophysical Union Fall Meeting*, San Francisco, Ca.-USA., EOS, 2000.;
- Klein F., Hypocentral location program HYPOINVERSE, *U. S. Geol. Survey Open File Rep.*, 78-694, 1978.
- Lamb, S., Hoke L., Kennan L. and Dewey J., *Geol. Soc. Spec. Pub.* , 121, 237-264, 1997.
- Muñoz, N., and Charrier R., *J. South Am. Earth Sci.*, 9, 171-181, 1996.
- Riquelme, R., Tesis de Magíster en Ciencias, Mención Geología, Universidad de Chile, 124 p., 1998.
- Rivera, L., and Cisternas A., *Bull. Seism. Soc. Am.*, 80, 600-614, 1990.
- Rochat, P., Hérail G., Baby P., and Mascle G., *C. R. Acad. Sci.*, 328, 189-195, 1999
- Rochat, P., PhD thesis, Université de Grenoble, 2000.

# Géodynamique andine Andean Geodynamics Geodinámica Andina

Résumés étendus  
Extended abstracts  
Resúmenes expandidos



**5th International Symposium**  
**Toulouse, France**  
**16-18 Sept. 2002**


**Organisateurs**

**Organizers**

**Organizadores**

**Institut de recherche  
pour le développement  
Paris**

**Université Paul Sabatier  
Toulouse  
France**



**GÉODYNAMIQUE ANDINE  
ANDEAN GEODYNAMICS  
GEODINAMICA ANDINA**

*5th International Symposium on Andean Geodynamics*

Université Paul Sabatier, Toulouse, France,  
16-18 Septembre 2002

**Résumés étendus  
Extended abstracts  
Resúmenes ampliados**

Organisateurs / Organizers / Organizadores  
Institut de recherche pour le développement  
Université Paul Sabatier

---

IRD  
INSTITUT DE RECHERCHE POUR LE DÉVELOPPEMENT  
Paris, 2002



**COMITÉ D'ORGANISATION**  
**COMITE ORGANIZADOR**  
**ORGANIZING COMMITTEE**

P. Baby (IRD-Toulouse), J. Darrozes (Univ. Paul Sabatier-Toulouse), J. Deramond (Univ. Paul Sabatier-Toulouse),  
B. Dupré (CNRS-Toulouse), J.-L. Guyot (IRD-Toulouse), G. Hérail (IRD-Toulouse),  
E. Jaillard (IRD-Quito), A. Lavenu (IRD-Toulouse), H. Miller (Univ. München),  
T. Monfret (IRD-Géoscience Azur), G. Wörner (Univ. Göttingen)

**Comité scientifique et représentants nationaux**  
**Comité Científico y Representantes Nacionales**  
**Scientific Advisory Board and National Representatives**

R. Armijo (IPG, Paris), J.-P. Avouac (CEA, Paris), R. Charrier (Univ. Chile, Santiago),  
J.-Y. Collot (IRD, Géoscience Azur), L. Dorbath (IRD, Strasbourg), S. Flint (Univ. Liverpool), B. France-Lanord  
(CNRS, Nancy), L. Fontboté (Univ. Genève), Y. Gaudemer (Univ. Paris VII), R. Gaupp (Univ. Jena),  
F. Hervé (Univ. Chile, Santiago), T.E. Jordan (INSTOC, Cornell), J. Mojica (Univ. Bogotá),  
O. Oncken (Univ. Potsdam), L. Ortlieb (IRD, Bondy), R.J. Pankhurst (Brit. Antarctic Surv.),  
V. Ramos (Univ. Buenos Aires), P. Ribstein (IRD, Paris),  
C. Robin (Univ. Clermont-Ferrand), S. Rosas (Univ. Lima), F. Sàbat (Univ. Barcelona),  
M. Schmitz (FUNVISIS, Caracas), R. Suárez Soruco (YPBF, La Paz),  
M. Rivadeneira (Petroproducción), W. Winkler (ETH, Zürich).

**APPUIS FINANCIERS**  
**FUNDINGS**  
**APPOYO FINANCIERO**

L'organisation de l'ISAG 2002 et les bourses accordées à un certain nombre de collègues  
latino-américains ont été possibles grâce au soutien financier de l'IRD  
(notamment de la Délégation à l'Information et à la Communication), de la région Midi-Pyrénées,  
de l'Université Paul Sabatier et de l'Andean Committee de l'ILP.