

LATE PLEISTOCENE AND HOLOCENE TEPHROSTRATIGRAPHY AND CHRONOLOGY IN SOUTHERN PERU

Etienne JUVIGNE(1), Jean-Claude THOURET(2), , Jersy MARIÑO(3), Marcela MOSCOL(4), A. LEGELEY-PADOVANI(5), Isabelle LOUTSCH(2)

(1)Laboratoire de Géomorphologie, Université de Liège, 4000 Liège, Belgique (ejuvigne@ulg.ac.be)

(2)Laboratoire Magmas et Volcans (UMR 6524 CNRS), Université Blaise Pascal, 63038 Clermont-Ferrand, France (thouret@opgc.univ-bpclermont.fr)

(3)Instituto Geofísico del Perú, Calle Clatrava 216, Urb. Camino Real, La Molina, Lima 12, Peru (irma@axil.igp.gob.pe)

(4)Université Montpellier II, Institut des Sciences de l'évolution, UMR CNRS 5554, 34095 Montpellier cedex 05 (mcmakosso@care2.com)

(5)Institut de Recherche pour le Développement, Centre Ile de France, 34 rue Henri-Varagnat, 93170 Bondy cedex, France (Annick.legeley-Padovani@bondy.ird.fr)

KEY WORDS: tephra, stratigraphy, chronology, Late Pleistocene, Holocene, Peru.

INTRODUCTION

Multidisciplinary investigations of stratigraphic sections and cores extracted from peat-bogs and lakes in the Western Cordillera of the southern Peruvian Andes, have enabled us to trace more than 40 tephras over the past 50,000 ¹⁴C years. At least eight pumice-fall deposits are widespread (>1,000 km²) and voluminous (≥1 km³), as shown in bold squares in Table 1.

LATE-GLACIAL AND HOLOCENE TEPHRAS

Two cores extracted from the *salar* of Laguna Salinas, 35 km east of Arequipa, include 7 tephra-fall deposits from Huaynaputina, Misti, and Ubinas over the past 15,000 years (Juvigné et al., 1997). Two recent tephra falls are found around the Ticsani domes: a dacitic pumice-fall deposit of 0.4 km³ has been dated ca. 11,600 yr BP; bombs and pumice preceding the youngest dome overlie the AD 1600 Huaynaputina ash. The recent explosive behaviour of Ubinas produced several tephra falls, including two widespread plinian pumice falls. The pumice-fall deposit >7840 yr BP and the ca. 980 yr BP-old pumice-fall deposit have a volume >1 km³ and may be linked to the formation of the summit caldera (Rivera et al., 1998). The peat sequence in the Sallalli peat-bog 10 km SE from the Nevado Sabancaya includes 4 tephra-fall layers: the AD 1600 Huaynaputina ash, an AD 1200-1400 Sabancaya/Ampato ash, a thin black ash probably from Misti ca. 2,370 yr BP, and a Sabancaya/Ampato ash of ca. 8,550 yr BP (Juvigné et al., 1998).

HISTORICAL TEPHRAS

The last subplinian explosive episode ca. 2,300-2,050 yr BP at Misti produced pumice-fall and flows ca. 1 km³ in volume. Spanish chronicles refer to an explosive eruption ca. AD 1440 and 1470: the small volume 'Pachacútec ash' affected the Incaic towns of Cayma and Chiguata (Thouret et al., 1999a, 2001a). The AD 1600 Huaynaputina plinian eruption produced the most widespread and voluminous (12 km³) pumice-fall deposit in the Andes in historical times (Thouret et al., 1999b). Tephra falls, pyroclastic flows, and surges disrupted life ~60 km around the volcano and ashfall was reported 250-500 km away in south Peru, west Bolivia and north Chile. Persistent eruptions of Nevado Sabancaya have dispersed a small volume of ash from May 1990 until 1998 as far as 20 km towards the east (Thouret et al., 1995, 2001b).

CONCLUSIONS

In sum, 1) heavy ashfall can recur every 500 to 1500 years on average but small ashfall can occur on a 100-years basis; 2) voluminous pumice-fall deposit can occur every 2000 to 4000 years on average; 3) three voluminous (≥ 1 km³) plinian eruptions occurred over the past 2,000 years; (4) the large-scale plinian and ignimbritic eruption of Huaynaputina produced 12 km³ of tephra and caused havoc and famine in southern Peru 400 years ago.

REFERENCES

- Juvigné E., Thouret J.-C., Gilot E., Gourgaud A., Legros F., Uribe M., Graf K., 1997. Etude téphrostratigraphique et bioclimatique du Tardiglaciaire et de l'Holocène de la Laguna Salinas, Pérou méridional. *Géographie physique et Quaternaire* (Canada), 51, 2: 219-231.
- Juvigné E., Thouret J.-C., Gilot E., Leclercq L., Gourgaud A., 1998. L'activité du volcan Nevado Sabancaya (Pérou) au cours de l'Holocène. *Quaternaire, Hommes et Volcans*, 9, 1, 45-51.

Géodynamique andine Andean Geodynamics Geodinámica Andina

Résumés étendus
Extended abstracts
Resúmenes expandidos



5th International Symposium
Toulouse, France
16-18 Sept. 2002


Organisateurs

Organizers

Organizadores

Institut de recherche
pour le développement
Paris

Université Paul Sabatier
Toulouse
France



**GÉODYNAMIQUE ANDINE
ANDEAN GEODYNAMICS
GEODINAMICA ANDINA**

5th International Symposium on Andean Geodynamics

Université Paul Sabatier, Toulouse, France,
16-18 Septembre 2002

**Résumés étendus
Extended abstracts
Resúmenes ampliados**

Organisateurs / Organizers / Organizadores
Institut de recherche pour le développement
Université Paul Sabatier

IRD
INSTITUT DE RECHERCHE POUR LE DÉVELOPPEMENT
Paris, 2002



COMITÉ D'ORGANISATION
COMITE ORGANIZADOR
ORGANIZING COMMITTEE

P. Baby (IRD-Toulouse), J. Darrozes (Univ. Paul Sabatier-Toulouse), J. Deramond (Univ. Paul Sabatier-Toulouse),
B. Dupré (CNRS-Toulouse), J.-L. Guyot (IRD-Toulouse), G. Hérail (IRD-Toulouse),
E. Jaillard (IRD-Quito), A. Lavenu (IRD-Toulouse), H. Miller (Univ. München),
T. Monfret (IRD-Géoscience Azur), G. Wörner (Univ. Göttingen)

Comité scientifique et représentants nationaux
Comité Científico y Representantes Nacionales
Scientific Advisory Board and National Representatives

R. Armijo (IPG, Paris), J.-P. Avouac (CEA, Paris), R. Charrier (Univ. Chile, Santiago),
J.-Y. Collot (IRD, Géoscience Azur), L. Dorbath (IRD, Strasbourg), S. Flint (Univ. Liverpool), B. France-Lanord
(CNRS, Nancy), L. Fontboté (Univ. Genève), Y. Gaudemer (Univ. Paris VII), R. Gaupp (Univ. Jena),
F. Hervé (Univ. Chile, Santiago), T.E. Jordan (INSTOC, Cornell), J. Mojica (Univ. Bogotá),
O. Oncken (Univ. Potsdam), L. Ortlieb (IRD, Bondy), R.J. Pankhurst (Brit. Antarctic Surv.),
V. Ramos (Univ. Buenos Aires), P. Ribstein (IRD, Paris),
C. Robin (Univ. Clermont-Ferrand), S. Rosas (Univ. Lima), F. Sàbat (Univ. Barcelona),
M. Schmitz (FUNVISIS, Caracas), R. Suárez Soruco (YPBF, La Paz),
M. Rivadeneira (Petroproducción), W. Winkler (ETH, Zürich).

APPUIS FINANCIERS
FUNDINGS
APPOYO FINANCIERO

L'organisation de l'ISAG 2002 et les bourses accordées à un certain nombre de collègues
latino-américains ont été possibles grâce au soutien financier de l'IRD
(notamment de la Délégation à l'Information et à la Communication), de la région Midi-Pyrénées,
de l'Université Paul Sabatier et de l'Andean Committee de l'ILP.