

# LATE PLEISTOCENE AND HOLOCENE TEPHROSTRATIGRAPHY AND CHRONOLOGY IN SOUTHERN PERU

*Etienne JUVIGNE(1), Jean-Claude THOURET(2), , Jersy MARIÑO(3), Marcela MOSCOL(4), A. LEGELEY-PADOVANI(5), Isabelle LOUTSCH(2)*

(1)Laboratoire de Géomorphologie, Université de Liège, 4000 Liège, Belgique (ejuvigne@ulg.ac.be)

(2)Laboratoire Magmas et Volcans (UMR 6524 CNRS), Université Blaise Pascal, 63038 Clermont-Ferrand, France (thouret@opgc.univ-bpclermont.fr)

(3)Instituto Geofísico del Perú, Calle Clatrava 216, Urb. Camino Real, La Molina, Lima 12, Peru (irma@axil.igp.gob.pe)

(4)Université Montpellier II, Institut des Sciences de l'évolution, UMR CNRS 5554, 34095 Montpellier cedex 05 (mcmakosso@care2.com)

(5)Institut de Recherche pour le Développement, Centre Ile de France, 34 rue Henri-Varagnat, 93170 Bondy cedex, France (Annick.legeley-Padovani@bondy.ird.fr)

**KEY WORDS:** tephra, stratigraphy, chronology, Late Pleistocene, Holocene, Peru.

## INTRODUCTION

Multidisciplinary investigations of stratigraphic sections and cores extracted from peat-bogs and lakes in the Western Cordillera of the southern Peruvian Andes, have enabled us to trace more than 40 tephras over the past 50,000 <sup>14</sup>C years. At least eight pumice-fall deposits are widespread (>1,000 km<sup>2</sup>) and voluminous (≥1 km<sup>3</sup>), as shown in bold squares in Table 1.

TEPHROSTRATIGRAPHY AND CHRONOLOGY IN SOUTHERN PERU, NORTH-CENTRAL ANDEAN VOLCANIC ZONE											
TABLE 1		SARA-SARA	COROPUNA	ANDALUJA-HUAMBO	AMPAKO MASSIF and NEVADO SARANEVA	EL MISTI	UBINAS	HUAYNAPUTINA	TICSANI	TUTUPACA	YUCAMANE
VOLCANOE TYPE PERIOD			STRATOVOLCANO and others	OROCOPAMA volcanic 1987	NEVADO SARANEVA lava domes	STRATOVOLCANO	STRATOVOLCANO	VELOCIDAD-CHITRA VELOCIDAD-CHITRA	VELOCIDAD-CHITRA	VELOCIDAD-CHITRA	VELOCIDAD-CHITRA
HISTORICAL	AD 1500			1990-98 ph / phreag / euc 1750 - 1784	1990-98 ph / phreag / euc 1750 - 1784	1784-87 1677 ph	active since 1852	Feb-March 1852		1900, 1852 1802 1780	1787
	Incaic			AD 1400-1600 at Chacayse cone	AD 1480 ± 20 ? AD 1680 ± 30 (PF) peñ-bog	AD 1440-20 at	at ph, primag 1677 at	pl, PF, ps (docel)		Sp, pt, at	pl, at
HOLOCENE	2000 yr BP		Andahu group			AD 1304-1398 at AD 553-865 at		AD 800 ± 60 PF			
	5000		2850 ± 50 at Kayoc cone 2675-2870 ± 50 at 4860 ± 50 at Ticsho and Paca Maxari cones	voluminous block-lava flows 5540 ± 40 at southwest flank		AD 1400 ± 160 PF PF 2300 ± 60 PF 2370 ± 90 at 2770 ± 190 at, PF 3890 ± 50 at 4750 ± 40 at 5200 ± 80 ps		Debris- avalanche deposit > 3670 ± 60			Tutupaca dome collapse pt
	10,000		Andahu group and Huambo			6320 ± 50 PF 8520 ± 80 at		> 7480 ± 40 R, pl	< 7480 ± 40 R, pl 8700 ± 150 P, pl		
LATE GLACIAL	15,000	P, pl	Oropampa group of subdued cones and lava flows	voluminous block-lava flows		~ 13,600 ± 240 TBI, pl, TBI, PL TBI, at, at	11,200 ± 70 11,480 ± 220 pl, PF, ps			11,600 ± 80 R, pl	pl
	20,000										
LATE PLEISTOCENE	25,000						20,860 ± 380-760 pl				
	30,000						24,840 ± 480 PF				
	35,000										
	40,000										
	45,000										
	50,000 yr BP										

LATE PLEISTOCENE TEPHRA

Late Pleistocene tephra have been dated on the west flank of Nevado Sara Sara between ca. 49,200 and 44,500 yr BP and ca. 44,000 yr BP on the south flank of Yucamane. Nevado Coropuna has probably been active before the Late Glacial; ashfalls in a soil section outside of the Last Glacial Maximum moraines yielded <sup>14</sup>C ages of ca. 27,200-37,370 yr BP. The composite Misti edifice comprises a stratovolcano termed Misti 1, partially overlapped by two stratocones termed Misti 2 and 3 (≤112 ka), and a summit cone Misti 4 ≤11 ka. Sustained explosive eruptions have delivered at least 12 pumice falls during the past ca. 50,000 years (Thouret et al., 2001b).

## LATE-GLACIAL AND HOLOCENE TEPHRAS

Two cores extracted from the *salar* of Laguna Salinas, 35 km east of Arequipa, include 7 tephra-fall deposits from Huaynaputina, Misti, and Ubinas over the past 15,000 years (Juvigné et al., 1997). Two recent tephra falls are found around the Ticsani domes: a dacitic pumice-fall deposit of 0.4 km<sup>3</sup> has been dated ca. 11,600 yr BP; bombs and pumice preceding the youngest dome overlie the AD 1600 Huaynaputina ash. The recent explosive behaviour of Ubinas produced several tephra falls, including two widespread plinian pumice falls. The pumice-fall deposit >7840 yr BP and the ca. 980 yr BP-old pumice-fall deposit have a volume >1 km<sup>3</sup> and may be linked to the formation of the summit caldera (Rivera et al., 1998). The peat sequence in the Sallalli peat-bog 10 km SE from the Nevado Sabancaya includes 4 tephra-fall layers: the AD 1600 Huaynaputina ash, an AD 1200-1400 Sabancaya/Ampato ash, a thin black ash probably from Misti ca. 2,370 yr BP, and a Sabancaya/Ampato ash of ca. 8,550 yr BP (Juvigné et al., 1998).

## HISTORICAL TEPHRAS

The last subplinian explosive episode ca. 2,300-2,050 yr BP at Misti produced pumice-fall and flows ca. 1 km<sup>3</sup> in volume. Spanish chronicles refer to an explosive eruption ca. AD 1440 and 1470: the small volume 'Pachacútec ash' affected the Incaic towns of Cayma and Chiguata (Thouret et al., 1999a, 2001a). The AD 1600 Huaynaputina plinian eruption produced the most widespread and voluminous (12 km<sup>3</sup>) pumice-fall deposit in the Andes in historical times (Thouret et al., 1999b). Tephra falls, pyroclastic flows, and surges disrupted life ~60 km around the volcano and ashfall was reported 250-500 km away in south Peru, west Bolivia and north Chile. Persistent eruptions of Nevado Sabancaya have dispersed a small volume of ash from May 1990 until 1998 as far as 20 km towards the east (Thouret et al., 1995, 2001b).

## CONCLUSIONS

In sum, 1) heavy ashfall can recur every 500 to 1500 years on average but small ashfall can occur on a 100-years basis; 2) voluminous pumice-fall deposit can occur every 2000 to 4000 years on average; 3) three voluminous ( $\geq 1$  km<sup>3</sup>) plinian eruptions occurred over the past 2,000 years; (4) the large-scale plinian and ignimbritic eruption of Huaynaputina produced 12 km<sup>3</sup> of tephra and caused havoc and famine in southern Peru 400 years ago.

## REFERENCES

- Juvigné E., Thouret J.-C., Gilot E., Gourgaud A., Legros F., Uribe M., Graf K., 1997. Etude téphrostratigraphique et bioclimatique du Tardiglaciaire et de l'Holocène de la Laguna Salinas, Pérou méridional. *Géographie physique et Quaternaire* (Canada), 51, 2: 219-231.
- Juvigné E., Thouret J.-C., Gilot E., Leclercq L., Gourgaud A., 1998. L'activité du volcan Nevado Sabancaya (Pérou) au cours de l'Holocène. *Quaternaire, Hommes et Volcans*, 9, 1, 45-51.

- Rivera M., Thouret J.-C. y Gourgaud A., 1998. Ubinas, el volcan más activo del Sur del Perú desde 1550: geología y evaluación de las amenazas volcánicas. *Boletín Sociedad Geológica del Perú*, 88, 53-71.
- Thouret J.-C., Gourgaud A., Uribe M., Rodríguez A., Guillaude R., Salas G., 1995. Geomorphological and geological survey, and spot remote sensing of the current activity of Nevado Sabancaya stratovolcano (south Peru): assessment for hazard-zone mapping. *Zeit. für Geomorphology*, 39, 4, 515-535.
- Thouret J.-C., Suni J., Eissen J.-Ph., Navarro P., 1999a. Assessment of volcanic hazards in the Arequipa area based on the eruptive history of Misti volcano, Southern Peru. *Zeitschrift für Geomorphology, Suppl.-Bd.114*, 89-112.
- Thouret J.-C., Davila J., and Eissen J.-Ph., 1999b. Largest historic explosive eruption in the Andes at Huaynaputina volcano, south Peru. *Geology*, 27, 5, 435-438.
- Thouret J.-C., Suni J., Finizola A., Fornari M., Legeley-Padovani, and Frechen M., 2001a. Geology of El Misti volcano near the city of Arequipa, Peru. Accepted to *Geological Society of America Bulletin*, June 2001.
- Thouret J.-C., Juvigné E., Loutsch I. et Chávez Chávez J.A., 2001b. Activité volcanique historique et sacrifices humains chez les Incas au Pérou méridional. In: E. Juvigné & JP Raynal, eds., *Tephres : chronologie/chronology, Archeologie/arquaeology*, p. 219-226, Les dossiers de l'Arqueo-Logis n°1, CDCRAD, Goudet.

# Géodynamique andine Andean Geodynamics Geodinámica Andina

Résumés étendus  
Extended abstracts  
Resúmenes expandidos



5th International Symposium  
Toulouse, France  
16-18 Sept. 2002

**Organisateurs**

**Organizers**

**Organizadores**

Institut de recherche  
pour le développement  
Paris

Université Paul Sabatier  
Toulouse  
France



**GÉODYNAMIQUE ANDINE  
ANDEAN GEODYNAMICS  
GEODINAMICA ANDINA**

*5th International Symposium on Andean Geodynamics*

Université Paul Sabatier, Toulouse, France,  
16-18 Septembre 2002

**Résumés étendus  
Extended abstracts  
Resúmenes ampliados**

Organisateurs / Organizers / Organizadores  
Institut de recherche pour le développement  
Université Paul Sabatier

---

IRD  
INSTITUT DE RECHERCHE POUR LE DÉVELOPPEMENT  
Paris, 2002



**COMITÉ D'ORGANISATION**  
**COMITE ORGANIZADOR**  
**ORGANIZING COMMITTEE**

P. Baby (IRD-Toulouse), J. Darrozes (Univ. Paul Sabatier-Toulouse), J. Deramond (Univ. Paul Sabatier-Toulouse),  
B. Dupré (CNRS-Toulouse), J.-L. Guyot (IRD-Toulouse), G. Hérail (IRD-Toulouse),  
E. Jaillard (IRD-Quito), A. Lavenu (IRD-Toulouse), H. Miller (Univ. München),  
T. Monfret (IRD-Géoscience Azur), G. Wörner (Univ. Göttingen)

**Comité scientifique et représentants nationaux**  
**Comité Científico y Representantes Nacionales**  
**Scientific Advisory Board and National Representatives**

R. Armijo (IPG, Paris), J.-P. Avouac (CEA, Paris), R. Charrier (Univ. Chile, Santiago),  
J.-Y. Collot (IRD, Géoscience Azur), L. Dorbath (IRD, Strasbourg), S. Flint (Univ. Liverpool), B. France-Lanord  
(CNRS, Nancy), L. Fontboté (Univ. Genève), Y. Gaudemer (Univ. Paris VII), R. Gaupp (Univ. Jena),  
F. Hervé (Univ. Chile, Santiago), T.E. Jordan (INSTOC, Cornell), J. Mojica (Univ. Bogotá),  
O. Oncken (Univ. Potsdam), L. Ortlieb (IRD, Bondy), R.J. Pankhurst (Brit. Antarctic Surv.),  
V. Ramos (Univ. Buenos Aires), P. Ribstein (IRD, Paris),  
C. Robin (Univ. Clermont-Ferrand), S. Rosas (Univ. Lima), F. Sàbat (Univ. Barcelona),  
M. Schmitz (FUNVISIS, Caracas), R. Suárez Soruco (YPBF, La Paz),  
M. Rivadeneira (Petroproducción), W. Winkler (ETH, Zürich).

**APPUIS FINANCIERS**  
**FUNDINGS**  
**APPOYO FINANCIERO**

L'organisation de l'ISAG 2002 et les bourses accordées à un certain nombre de collègues  
latino-américains ont été possibles grâce au soutien financier de l'IRD  
(notamment de la Délégation à l'Information et à la Communication), de la région Midi-Pyrénées,  
de l'Université Paul Sabatier et de l'Andean Committee de l'ILP.