

# VALLEY EVOLUTION, UPLIFT, VOLCANISM, AND RELATED HAZARDS IN THE CENTRAL ANDES OF SOUTHERN PERU

*Jean-Claude THOURET(1), Gerhard WÖRNER(2), A. FINIZOLA(1), and A. LEGELEY-PADOVANI(3)*

- (1) Laboratoire Magmas et Volcans, Université Blaise Pascal et CNRS, 63038 Clermont-Ferrand cedex, France (thouret@opgc.univ-bpclermont.fr)
- (2) Geoschemisches Institut, Goldschmidtstrasse 1, Universität Göttingen, 37077, Göttingen, Germany (gwoerne@dwdg.de)
- (3) IRD, Centre Ile de France, 34 rue Henri-Varagnat, 93193 Bondy cedex, France (A.Legeley-Padovani@bondy.ird.fr)

**KEY WORDS:** Andes, Peru, valley history, volcanoes, ignimbrites, uplift, canyons, downcutting, hazards.

## INTRODUCTION

Interpretation of satellite images, field work, and geochronology identify three principal types of volcanoes in the areas of Rio Cotahuasi-Rio Ocoña, and Arequipa in southern Peru (Figs. 1-3): (1) relatively youthful stratovolcanoes of Pleistocene to Recent age e.g. El Misti ( $\leq 0.8$  Ma) and older, larger complexes such as Nevado Coropuna, (2) deeply eroded stratovolcanoes of Plio-Quaternary age (4-1.5 Ma) such as Nevado Solimana and Chachani, and; (3) subdued 'shield' volcanoes of Late Miocene age. While the former are rather varied in petrography and hence chemical composition of erupted magmas, the latter tend to be comprised more of monotonous mafic andesites. Ignimbrites form of several hundreds of km<sup>2</sup> in area and several tens of km<sup>3</sup> in volume are also observed and have been dated at middle Miocene (13-14 Ma), Pliocene (5-2 Ma), and Pleistocene age (~1 Ma) (Thouret et al., 2001).

In the area of Arequipa, 13-14 M ignimbrites outcrop above the Jurassic basement in the Rio Chili valley that cuts the flank of the Western Cordillera (Fig. 2). The upper flanks of Rio Chili valley are built in part by lava flows and volcaniclastic sediments of Plio-Quaternary age, whose sources are the Chachani massif and El Misti. Downcutting by 250 m has been achieved within the past 3 My (Fig. 2).

In contrast, the headvalleys of the deepest canyons on Earth of Rios Ocoña, Cotahuasi and Colca NW of Arequipa, have been cut 2 km down in Miocene volcanic rocks and 1 km further down in Cretaceous intrusive and Jurassic sedimentary rocks. The headvalleys were repeatedly filled by pyroclastics and lava flows of Neogene age (< 1 Ma, Fig. 3) and have subsequently been recut below their original thalweg. These canyons are thus much deeper and older (middle Miocene) than Rio Chili.





Huaylillas plateau near Chuquibamba (Fig. 3). Downcutting also triggered flank failures on the Plio-Quaternary volcanoes, such as the southwest flank of Nevado Solimana. Subsequent debris avalanches have dammed the upper course of Rios Cotahuasi and Rio Huarcaya in Pleistocene time. Downcutting continues and valley flank and volcano instabilities pose a major hazard in these ultr-deep canyons. Further hazards are also related to potential dam breakouts that may trigger devastating debris flows toward the populated lower valleys.

## REFERENCES

Kennan L., 1999. Large-scale geomorphology in the Andes: interrelationships of tectonics, magmatism, and climate. *In*: M.A. Summerfield, ed., *Geomorphology and Global Tectonics*, p. 167-199, J. Wiley, New York.

Thouret J.-C., Finizola A., Fornari M., Suni J., Legeley-Padovani A., and Frechen M., 2001. Geology of El Misti volcano nearby the city of Arequipa, Peru. *Geological Society of America Bulletin*, vol. 113, n°12, 1593-1610.

# Géodynamique andine Andean Geodynamics Geodinámica Andina

Résumés étendus  
Extended abstracts  
Resúmenes expandidos



**5th International Symposium**  
**Toulouse, France**  
**16-18 Sept. 2002**


**Organisateurs**

**Organizers**

**Organizadores**

**Institut de recherche  
pour le développement  
Paris**

**Université Paul Sabatier  
Toulouse  
France**



**GÉODYNAMIQUE ANDINE  
ANDEAN GEODYNAMICS  
GEODINAMICA ANDINA**

*5th International Symposium on Andean Geodynamics*

Université Paul Sabatier, Toulouse, France,  
16-18 Septembre 2002

**Résumés étendus  
Extended abstracts  
Resúmenes ampliados**

Organisateurs / Organizers / Organizadores  
Institut de recherche pour le développement  
Université Paul Sabatier

---

IRD  
INSTITUT DE RECHERCHE POUR LE DÉVELOPPEMENT  
Paris, 2002



**COMITÉ D'ORGANISATION**  
**COMITE ORGANIZADOR**  
**ORGANIZING COMMITTEE**

P. Baby (IRD-Toulouse), J. Darrozes (Univ. Paul Sabatier-Toulouse), J. Deramond (Univ. Paul Sabatier-Toulouse),  
B. Dupré (CNRS-Toulouse), J.-L. Guyot (IRD-Toulouse), G. Hérail (IRD-Toulouse),  
E. Jaillard (IRD-Quito), A. Lavenu (IRD-Toulouse), H. Miller (Univ. München),  
T. Monfret (IRD-Géoscience Azur), G. Wörner (Univ. Göttingen)

**Comité scientifique et représentants nationaux**  
**Comité Científico y Representantes Nacionales**  
**Scientific Advisory Board and National Representatives**

R. Armijo (IPG, Paris), J.-P. Avouac (CEA, Paris), R. Charrier (Univ. Chile, Santiago),  
J.-Y. Collot (IRD, Géoscience Azur), L. Dorbath (IRD, Strasbourg), S. Flint (Univ. Liverpool), B. France-Lanord  
(CNRS, Nancy), L. Fontboté (Univ. Genève), Y. Gaudemer (Univ. Paris VII), R. Gaupp (Univ. Jena),  
F. Hervé (Univ. Chile, Santiago), T.E. Jordan (INSTOC, Cornell), J. Mojica (Univ. Bogotá),  
O. Oncken (Univ. Potsdam), L. Ortlieb (IRD, Bondy), R.J. Pankhurst (Brit. Antarctic Surv.),  
V. Ramos (Univ. Buenos Aires), P. Ribstein (IRD, Paris),  
C. Robin (Univ. Clermont-Ferrand), S. Rosas (Univ. Lima), F. Sàbat (Univ. Barcelona),  
M. Schmitz (FUNVISIS, Caracas), R. Suárez Soruco (YPBF, La Paz),  
M. Rivadeneira (Petroproducción), W. Winkler (ETH, Zürich).

**APPUIS FINANCIERS**  
**FUNDINGS**  
**APPOYO FINANCIERO**

L'organisation de l'ISAG 2002 et les bourses accordées à un certain nombre de collègues  
latino-américains ont été possibles grâce au soutien financier de l'IRD  
(notamment de la Délégation à l'Information et à la Communication), de la région Midi-Pyrénées,  
de l'Université Paul Sabatier et de l'Andean Committee de l'ILP.