

# Methylmercury production and exchanges in sediments of Lake Titicaca

Stéphane Guédron<sup>1,2</sup>, Stéphane Audry<sup>3</sup>, Dario Acha<sup>4</sup> and David Amouroux<sup>4,5</sup>.

<sup>1</sup>Institut de Recherche pour le Développement (IRD, ISTerre, UGA – Grenoble, France).

<sup>2</sup> Instituto de Investigaciones Químicas - Universidad Mayor de San Andres, La Paz, Bolivia.

<sup>3</sup> Géosciences Environnement Toulouse, Université Paul Sabatier, Toulouse, France.

<sup>4</sup> Laboratorio de Calidad Ambiental (LCA) – Instituto de Ecología - La Paz, Bolivia.

<sup>5</sup> LCABIE/IPREM, Université de Pau et des Pays de l'Adour, 64053 Pau cedex 9, France.

## Abstract :

Lake sediment is known to be an important source of the neurotoxic monomethylmercury (MMHg) for surface water (Fitzgerald et al., 2014), in which trophic transfer occur in both benthic and pelagic organisms (Watras et al., 1998). Geochemical processes involved in the methylation of mercury (Hg) generally occurs in surface sediment or biofilms developed at the sediment/water interface (SWI) and imply various micro-organism amongst which the most documented are sulfate reducing (SRB) and iron reducing (IRB) bacteria (Benoit et al., 1999). Such processes occur specifically during early diagenesis, when a part of deposited fresh organic matter (OM) is mineralized close to the SWI, with O<sub>2</sub>, NO<sub>3</sub><sup>-</sup>, Fe- and Mn-oxyhydroxides or SO<sub>4</sub><sup>2-</sup> acting as electron acceptors (De Lange, 1986; Froelich et al., 1979).


In this study, we provide a high resolution (mm scale) complete Hg speciation, major and trace elements in porewater and solid sediment from seven short core collected in Lake Titicaca.

Total Hg concentrations in sediment of Lake Titicaca are low in both shallow (depth < 10 m, THg = 23 ± 14 ng g<sup>-1</sup>, N= 110) and deep areas (depth > 10 m, THg = 52 ± 52 ng g<sup>-1</sup>, N= 101) with MMHg concentrations representing 2.9 ± 6.7 and 0.3 ± 0.1 %, in shallow and deep areas respectively. In opposition to sediment, MMHg concentrations are elevated in porewater, especially in shallow sulfate and carbonate rich sediments (2.2 ± 1.8 ng L<sup>-1</sup>), where SRB are the likely major methylating organisms. Calculation of diffusive fluxes, highlighted very high MMHg diffusive fluxes being 91.8 ± 13.8 and 13.2 ± 2.0 ng m<sup>-2</sup> d<sup>-1</sup> in shallow and deep areas, respectively. Such fluxes would represent around 30 % of the MMHg contribution to the water column of Lake Huiñaimarca. Therefore, the major outcome of this study is that although THg concentration in sediment are low, the shallow carbonate facies and sulfate rich sediment are a major source of MMHg to the water column of the small Lake Titicaca (Huiñaimarca).

## References:

- Benoit, J.M., Gilmour, C.C., Mason, R.P. and Heyes, A., 1999. Sulfide controls on mercury speciation and bioavailability to methylating bacteria in sediment pore waters. *Environ. Sci. Technol.*, 33(6): 951-957.
- De Lange, G.J., 1986. Early diagenetic reactions in interbedded pelagic and turbiditic sediments in the Nares Abyssal Plain (western North Atlantic): Consequences for the composition of sediment and interstitial water. *Geochimica et Cosmochimica Acta*, 50(12): 2543-2561.
- Fitzgerald, W.F., Lamborg, C.H. and Turekian, H.D.H.K., 2014. 11.4 - Geochemistry of Mercury in the Environment, Treatise on Geochemistry (Second Edition). Elsevier, Oxford, pp. 91-129.
- Froelich, P.N. et al., 1979. Early oxidation of organic matter in pelagic sediments of the eastern equatorial atlantic: suboxic diagenesis. *Geochimica et cosmochemica Acta*, 43: 1075-1090.
- Watras, C.J. et al., 1998. Bioaccumulation of mercury in pelagic freshwater food webs. *Sci. Tot. Environ.*, 219(2-3): 183-208.

# COLOQUIO INTERNACIONAL SOBRE LA CONTAMINACIÓN ACTUAL E HISTÓRICA EN LOS ECOSISTEMAS ACUÁTICOS ANDINOS



**La Paz, 3 al 5 de mayo de 2016**  
**Universidad Mayor de San Andrés, Cota Cota, La Paz**





## Proceedings

**International colloquium on current and ancient contamination in  
Andes aquatic ecosystems**

**Coloquio internacional sobre la contaminación actual y histórica  
en los ecosistemas acuáticos Andinos**

**Colloque international sur la contamination actuelle et historique  
des écosystèmes aquatiques andins**

**La Paz – May 3<sup>rd</sup> – 5<sup>th</sup> 2016**

**Universidad Mayor de San Andrés – Campus de Cota-cota, La Paz**

**Organization direction:**

Stéphane Guédron (ISTerre-IRD/UMSA): stephane.guedron@ird.fr

Dario Acha Cordero (LCA/UMSA): darioacha@yahoo.ca

Marc-Antoine Vella (IFEA): mav.vella@gmail.com

Oswaldo Eduardo Ramos Ramos (IIQ/UMSA) : rroe@kth.se

**Organization committee:**

Stéphane Guédron (ISTerre-IRD/UMSA): stephane.guedron@ird.fr

Dario Acha Cordero (LCA/UMSA): darioacha@yahoo.ca

David Amouroux (LCABIE-IPREM/CNRS): david.amouroux@univ-pau.fr

Marc-Antoine Vella (IFEA): mav.vella@gmail.com

Christophe Delaere (ULB): Christophe.Delaere@ulb.ac.be

Oswaldo Eduardo Ramos Ramos (IIQ/UMSA) : rroe@kth.se

Mauricio. R. Ormachea Muñoz (IIQ/UMSA) : mauormache@gmail.com

Jorge Quintanilla (IIQ/UMSA): ceedi77@gmail.com

David Point (GET-IRD/UMSA): david.point@ird.fr

Céline Duwig (LTHE-IRD/UMSA): celine.duwig@ird.fr

## General Planning

- May 3<sup>rd</sup> 2016: Contamination and eutrophication of Lake Titicaca

*AM session: Mercury biogeochemistry and contamination of aquatic ecosystems of the Andes region*

Keynote Lecture: Hg contamination in Latin America: the past is not what we think, nor the future (J.-R. Davee Guimarães).

*PM session: Chemical contamination, eutrophication and monitoring of Lake Titicaca and its watershed*

Keynote Lecture: Eutrophication of the Cohana Bay (D. Acha).

- May 4<sup>th</sup> 2016: Arsenic issues in the Andes

*AM session: Arsenic biogeochemistry and contamination of aquatic ecosystems of the Andes region*

Keynote Lecture: Arsenic contamination of groundwater (Chile) (G. Lobos).

*PM session:*

*Workshop 1: Arsenic and mercury speciation.*

*Workshop 2: Paleoenvironmental studies in the Andean altiplano.*

- May 5<sup>th</sup> 2016: Historical reconstructions of the human-climate interactions in the altiplano: implication of archeological purposes

*AM session: Paleo-*

*archives*

Keynote Lecture: Holocene Paleoclimatic and Paleoenvironmental History of the Lake Titicaca Basin (S. Fritz & P. Baker).

*PM session: Archeology: historical human environment interactions*

Keynote Lecture: Recent contribution of terrestrial and subaquatic archeological investigation in Lake Titicaca (C. Delaere & M-A. Vella).