



Distribution of pelagic fish eggs off the Senegalese Coast during an intense upwelling event in March 2014

Julian DÖRING^{1, *}, Timothée BROCHIER^{2,3}, Eric MACHU⁴, Xavier CAPET⁵, Oumar MANNÉ^{2,3}, Werner EKAU¹ and Patrice BREHMER^{2,3}

¹ Leibniz Centre for Tropical Marine Ecology, Fahrenheitstraße 6, 28359 Bremen, Germany

² Institut de Recherche pour le Développement (IRD), UMR 195 LEMAR, BP 1386 Dakar, Senegal

³ ISRA/CRODT, Centre PRH, BP 2221, Dakar, Senegal

⁴ IRD, UMR LPO - based at LPAO-SF, ESP, Université Cheikh Anta Diop, BP 5085, Dakar Fann, Sénégal

⁵ CNRS-IRD-Sorbonne Universités (UPMC Univ. Paris 6)-MNHN, LOCEAN/IPSL Laboratory, 4 Place Jussieu, Paris 75252, France

*Correspondance: Tél: (+221) 77 87 04 100; Courriel: julian.doering@zmt-bremen.de (J. DÖRING)

Reçu le 09/12/2014; publié le 15/05/2015

AWA © MS WP2_S2_4_42

Abstract

The upwelling ecosystem off the coast of Senegal is highly productive and usually dominated in biomass by *Sardinella aurita*. During the AWA cruise in March of 2014 (F/RV Thalassa off Mauritania, Senegal, and the Gambia) continuous egg samples were collected onboard with a CUFES (Continuous Underwater Fish Eggs Sampler). Laboratory analysis of these samples show that very few eggs collected had a size corresponding to *S. aurita* eggs. Instead, a large part of the eggs collected could be identified as *Engraulis encrasicolus*, *Sardina pilchardus*, or *Diplodus bellottii*. As of now, no data is available on spawning behavior and locations of these species in Senegalese waters. Additionally, the coastline (20 m isobath) South-East of Dakar was repeatedly sampled. The obtained results not only indicate a southward migration of *S. pilchardus* down to at least 14°30' N, when compared to data from the early 1980s, but also pose as a proof for continuous spawning activity of *S. pilchardus* and *E. encrasicolus* in the area South-East of Dakar. We suggest that the very intense upwelling event that occurred during the cruise was responsible for the spawning event of *S. pilchardus*, and *E. encrasicolus* and also for the absence of *S. aurita* eggs. Finally, it is highlighted that, especially for *D. Bellotii*, the highest egg densities were found very near the planed Dakar city wastewater emissary off Mbaou (South coast of Senegal).

Keywords: Continuous Underwater Fish Eggs Sampler, CUFES, clupeidae, Senegal.



Commission Sous-Régionale des Pêches
Sub-Regional Fisheries Commission



Book of Abstract International Conference ICAWA 2014

THE AWA PROJECT
Ecosystem Approach
to the management
of fisheries and the
marine environment
in West African waters

Cap-Vert

Mauritanie

Sénégal

Gambie

Guinée Bissau

Guinée

Sierra Leone



Bundesministerium
für Bildung
und Forschung



Institut de recherche
pour le développement



Liberté - Égalité - Fraternité
RÉPUBLIQUE FRANÇAISE
Ministère des Affaires étrangères
et du Développement international
Ministère de l'Enseignement
Supérieur et de la Recherche

Trilateral German-French-African research initiative

ISBN :978-2-9553602-0-1

Edited by:

Patrice Brehmer (IRD) & Hamady Diop (SRFC/CSRP)

With the collaboration of:

Marie Madeleine Gomez, Ndague Diogoul, Viviane Koutob, Peter Brandt, Bamol Ali Sow, Alban Lazar, Xavier Capet, Heino Fock, Carlos F. Santos, Eric Machu, Hamet Diadihou, Didier Jouffre, Ibrahima Diallo, Joern Schmidt, Amadou Gaye, Mahfoudhould Taleb Sidi, Yves Gouriou, Rafael Almar, Moussa Sall, Dominique Duval Diop, Modou Thiow, Ross Wanless, Jacob Gonzales-Solis Bou, Ibrahima Ly, Dienaba Beye Traoré, Marie Bonnin, Werner Ekau.

ISBN: 978-2-9553602-0-0

SRFC/CSRP – IRD ©2015

Cover design: AWA (BMBF – IRD) project

Logo and flyers: Laurent Corsini (IRD)

Sponsors ICAWA 2014

