

Rethinking the management of mycorrhizal soil infectivity to restore Mediterranean and tropical forest ecosystems

Robin DUPONNOIS
IRD, France

Lahcen OUAHMANE
Cadi Ayyad University of Marrakech, Morocco

Mohamed HAFIDI
Cadi Ayyad University of Marrakech, Morocco

Yves PRIN
CIRAD, France

Introduction

Desertification, one of the main causes of climate change, generally results from a variety of factors, including climatic variations and human activities. Among the man-mediated degradative activities, deforestation is considered to have a major impact by causing extinction, changes to climatic condition, desertification and the displacement of local populations (Defries et al. 2007).

Forest cover is an important source of protection from soil degradation and deforestation which impacts population structure, successional patterns and species diversity, generally inducing degradations in the physico-chemical and biological soil properties (Requena et al. 2001). These changes can be recorded in microbial functional capacity (microbial metabolism, biomass and composition, enzymatic activities and soil organic matter flux) which is mainly involved in soil quality and function (Chaer et al. 2009). Many studies have also shown that deforestation and soil cultivation alter soil microbial community structure (Bossio et al. 2005) and may lead to reduction in microbial biodiversity (Chaer et al. 2009).

Among components of soil microbiota, mycorrhizal fungi are known to be essential key components of sustainable soil-plant systems, especially in arid ecosystems (Duponnois et al. 2011). The mycorrhizal symbiosis mobilizes and transports nutrients to roots (Smith & Read, 2008), reduces water stress (Augé, 2001) and improves soil aggregation in eroded soils (Caravaca et al. 2002). It has also been reported that arbuscular mycorrhizal (AM) fungi affect the diversity of plant communities (van der Heijden et al. 1998) and influence relationships between plants (van der Heijden et al. 1998).

Since trees are a primary source of protection from soil degradation, many afforestation programs have been undertaken but with generally low performance in terms of productivity and seedling survival after outplanting (Duponnois et al. 2005a). These deficiencies have usually been recorded in Mediterranean semi-arid areas known to have low bioavailable phosphate content and high phosphate retention capacities (Duponnois et al. 2011). This environmental context represents the most favourable conditions for AMF's potential to increase plant growth without any mineral fertilizers (Rodriguez & Sanders, 2015). Unfortunately, this microbial resource has been largely neglected despite numerous studies focused on this symbiosis.

The native inoculum potential of AM fungi in arid and semi-arid Mediterranean ecosystems is generally limited which, in turn, prevents plant establishment and growth (Smith & Read 2008). It is necessary to apply mycorrhizal inoculation technologies or to manage native AM fungus communities to replace or reinforce the mycorrhizal potential in these degraded areas (Duponnois et al. 2011).

Approaches to AM application in forestry practices

This chapter aims to describe different practical approaches to integrate the AM symbiosis in forestry practices through a “reductionist” approach (also named Controlled mycorrhization) (Inoculation of optimized AM fungal strains to improve the plant growth in unfriendly conditions) or a “holistic” approach (Suitable management of AM fungal diversity for ensuring AM fungi – dependent ecosystem services) (Rodriguez & Sanders, 2015). Each of these cultural practices will be illustrated by results from field experiments performed in Mediterranean and tropical areas.

The “reductionist” approach (controlled mycorrhization)

In recent decades, considerable research has been made by using specific mycorrhizal fungal strains to enhance outplanting performances with forest tree species (Caravaca et al. 2002). Hence, numerous studies have reported the beneficial effects on plant growth resulting from AM fungal inoculation during the nursery plantation. Among all the AM fungal strains tested in these experiments, *Rhizophagus irregularis* has attracted great interest because of (1) its world-wide distribution, (2) its high genetic variability and variation in effects on plant growth and (3) its ability to be produced in an *in vitro* system.

Some of its impacts on the growth of different tree species in controlled conditions are reported in table 1. Most of these experiments have been performed in controlled conditions and few studies have clearly demonstrated the benefits

Table 1
Impact of *R. irregularis* on the growth of tree species
in controlled conditions after different times of cultivation

Tree species	Time (months)	Impact of <i>R. irregularis</i> on plant growth		References
		On shoot biomass	On root biomass	
<i>Acacia holosericea</i>	4	+ 77.5% ⁽¹⁾	+ 122.7% ⁽²⁾	Duponnois et al. (2005b)
<i>Fraxinus uhdei</i>	3	+ 250%	+ 180%	Ambriz et al. (2010)
<i>Cupressus atlantica</i>	3	+ 48.7%	+ 155.2%	Ouahmane et al. (2007)
<i>Acacia mangium</i>	3	+ 20.2%	+ 50%	Weber et al. (2005)
<i>Citrus aurantium</i>	5	+ 273.3%	+ 66.3%	Nemec & Vu (1990)
<i>Acacia tortilis</i>	4	+ 499.2%	+ 78.3%	André et al. (2003)
<i>Parkia biglobosa</i>	2	+7.1%	+ 25.5%	Guissou et al. (1998)
<i>Tamarindus indica</i>	2	+6.7%	+ 36%	Guissou et al. (1998)
<i>Zizyphus mauritiana</i>	2	+ 130.3%	+ 152%	Guissou et al. (1998)
<i>Alnus cordata</i>	2	+ 441%	+ 1407%	Monzon & Azcon (2001)
<i>Alnus incana</i>	2	+ 644%	+ 1041.2%	Monzon & Azcon (2001)
<i>Alnus glutinosa</i>	2	+ 996.7 %	+ 1194%	Monzon & Azcon (2001)
<i>Acacia senegal</i>	3	+ 171.4%	+ 127.3%	Ndoye et al. (2013)
<i>Phoenix dactylifera</i>	5	+ 140.4%	+ 165.4%	Baslam et al. (2014)

⁽¹⁾ (Shoot biomass of mycorrhizal plants / Shoot biomass of non mycorrhizal plants) x 100. ⁽²⁾ (Root biomass of mycorrhizal plants / Root biomass of non mycorrhizal plants) x 100.

of fungal inoculation in the field. The degree of mycorrhizal responses on a reforestation site depends on the status of fungal colonization at planting, and the persistence of introduced fungi and other biotic and abiotic factors at the planting site (Duponnois et al. 2011).

Hence, the use of AM fungi and plants adapted to the local environmental conditions may be a prerequisite for the success of reforestation programmes (Duponnois et al. 2005a). The potential effect of mycorrhizal inoculation with native AM fungi on the survival rates and early growth performance in the field of Mediterranean tree species (i.e. cypress, carob) has been assessed in a few studies (Manaut et al. 2015). The results showed the high potential of this approach by sustainably improving the growth and nutrient status of both tree species and also by inducing a positive soil microbial environment for nutrient cycling and environmental stress resistance (figs. 1 & 2).


The “holistic” approach

It has been reported that certain shrubs react positively to the survival and growth of other neighboring plant species by creating a better environmental habitat with low stresses from high radiation and temperature as well as from soil nutrient and moisture deficiencies (Callaway & Walker, 1997) named “the nurse-plant syndrome” (Niering et al. 1963). The ecological facilitation between plant species results in the patchy distribution of the vegetation commonly observed in Mediterranean areas, especially in degraded ecosystems (Callaway & Walker, 1997).

Hence it has been suggested that the use of nurse plants as planting microhabitats in Mediterranean degraded ecosystems could promote the survival and development of native tree species and constitute an alternative reforestation technique compared to the standard practices (Duponnois et al. 2011). The “fertility islands” or “resource islands” (Schlesinger et al. 1996) resulting from the establishment of these nurse plants show a higher arbuscular mycorrhizal (AM) soil infectivity compared to the adjacent soil away from plant influence (Duponnois et al. 2011), which can improve plant growth and survival in arid conditions, by increasing the supply of nutrients to the plants (especially for soil P uptake) (Smith & Read, 2008), enhancing soil aggregation in eroded soils (Caravaca et al. 2002) and reducing water stress (Augé, 2001).

After three years’ plantation, it was reported that the association between *C. atlantica* and a nurse plant, *L. stoechas*, enhanced the growth of *C. atlantica* and provided better soil microbial characteristics compared to the control treatment (fig. 3) (Duponnois et al. 2011). AM mycelium network, total microbial activity, dehydrogenase activity, phosphate-solubilizing fluorescent pseudomonads and N, P nutrient uptake by *C. atlantica*, were significantly higher in the presence of *L. stoechas*. This pioneer shrub facilitated the early establishment of Cypress seedlings by improving soil microbial characteristics and AM fungus community development. Since the facilitative effect of one

A. HEIGHT



B. COLLAR DIAMETER




Figure 1

Height and collar diameter of carob outplants in the field, either inoculated with AM fungi (■) or non-inoculated (control □). An asterisk indicates a significant ($P < 0.05$) difference between the two treatments for a given year (From Manaut et al. 2015).




Figure 2

Cumulative mortality of carob outplants in the field, either inoculated with AM fungi (■) or non-inoculated (control □) during the three years of plantation (From Manaut et al. 2015).




Figure 3

Time course changes in plant height (expressed in cm) of *C. atlantica* outplants growing under natural conditions in the High Atlas Mountains (Morocco), either non-inoculated (Control) (●) or associated with *L. stoechas* plants (■). Symbols represent means (\pm standard error of the mean). An asterisk indicates that the difference between the height of uninoculated *C. atlantica* and *C. atlantica* associated with *L. stoechas* is significant in the corresponding month according to the Newman Keul's test ($p < 0.05$).

plant species on another increases with abiotic stress (Callaway, 1995), the benefits of this technique would be useful in reforestation programs undertaken to rehabilitate degraded areas in the Mediterranean region (Duponnois et al. 2011). Other shrub species have been identified for their potential nursing effects on Mediterranean tree species (fig. 4).

Conclusion

These data show that the management of the mycorrhizal soil infectivity through different cultural approaches (reductionist or holistic approaches) has large potentialities to improve the performances of afforestation programmes, especially in Mediterranean and Tropical areas. This biological tool must be used according to the biological characteristics of the targeted areas (physico-chemical characteristics, biological characteristics) in order to reach sustainable objectives in forest ecosystem productivity and resistance. Hence ecological approaches at community and population scales must be encouraged with a view to better informed management of AM fungi in order to propose practical solutions to manage forest ecosystems in a sustainable manner.




Figure 4

Growth responses of *Acacia raddiana* seedlings to the soil origins collected under shrub species native from Moroccan arid areas after 4 months' culture in glasshouse conditions (Unpublished data). HL: *Helianthemum lupii*; ON: *Ononis natrix*; HS: *Haloxylon scoparium*; RR: *Retama retama*; WA: *Withania adpressa*; LS: *Lavandula sp.*; LAS: *Launea sp.*; CL: *Cleome sp.*; CT: *Convolvulus trautianus*; AA: *Artemisia herba alba*; AS: *Astericus sp.*; SHC: Soil non influenced by plants.

References

AMBRIZ, E., BAEZ-PEREZ, A., SANCHEZ-YANEZ, J.M., MOUTOGLIS, P. & VILLEGAS, J. (2010)

Fraxinus–Glomus–Pisolithus symbiosis: Plant growth and soil aggregation effects. *Pedobiologia*, 53: 369-373.

- ANDRÉ, S., NEYRA, M. & DUPONNOIS, R. (2003) Arbuscular Mycorrhizal Symbiosis Changes the Colonization Pattern of *Acacia tortilis* spp. *Raddiana* Rhizosphere by Two Strains of Rhizobia. *Molecular Ecology*, 45: 137-144.
- AUGÉ, R.M. (2001) Water relations, drought and vesiculararbuscular mycorrhizal symbiosis. *Mycorrhiza*, 11: 3-42.
- BASLAM, M., QADDOURY, A. & GOICOECHEA, N. (2014) Role of native and exotic mycorrhizal symbiosis to develop morphological, physiological and biochemical responses coping with water drought of date palm, *Phoenix dactylifera*. *Trees*, 28: 161-172.
- BOSSIO, D.A., GIRVAN, M.S., VERCHOT, L., BULLIMORE, J., BORELLI, T., ALBRECHT, A., SCOW, K.M., BALL, A.S., PRETTY, J.N. & OSBORN, A.M. (2005) Soil microbial community response to land use change in an agricultural landscape of Western Kenya. *Microbial Ecology*, 49: 50-62.
- CALLAWAY, R.M. (1995) Positive interactions among plants. *Botanical Review*, 61: 306-349.
- CALLAWAY, R.M. & WALKER, L.R. (1997) Competition and facilitation: a synthetic approach to interactions in plant communities. *Ecology*, 78: 1958-1965.
- CARAVACA, F., BAREA, J.M., FIGUEROA, D. & ROLDAN, A. (2002) Assessing the effectiveness of mycorrhizal inoculation and soil compost addition for reforestation with *Olea europaea* subsp. *sylvestris* through changes in soil biological and physical parameters. *Applied Soil Ecology*, 20: 107-118.
- CHAER, G., FERNANDES, M., MYROLD, D. & BOTTOMLEY, P. (2009) Comparative resistance and resilience of soil microbial communities and enzyme activities in adjacent native forest and agricultural soils. *Microbial Ecology*, 58: 414-424.
- DEFRIES, R., ACHARD, F., HEROLD, M., MURDIVARSO, D., SCHLAMADINGER, B. & DESOUZAIR, C. (2007) "Earth observations for estimating greenhouse gas emissions from deforestation in developing countries". *Environmental Science Policy*, 10 (4): 385-394.
- DUPONNOIS, R., FOUNOUNE, H., MASSE, D. & PONTANIER, R. (2005A) Inoculation of *Acacia holosericea* with ectomycorrhizal fungi in a semiarid site in Senegal: growth response and influences on the mycorrhizal soil infectivity after 2 years plantation. *Forest Ecology & Management*, 207: 351-362.
- DUPONNOIS, R., COLOMBET, A., HIEN, V. & THIOULOUSE, J. (2005B) The mycorrhizal fungus *Glomus intraradices* and rock phosphate amendment influence plant growth and microbial activity in the rhizosphere of *Acacia holosericea*. *Soil Biology & Biochemistry*, 37: 1460-1468.
- DUPONNOIS, R., OUAHMANE, L., KANE, A., THIOULOUSE, J., HAFIDI, M., BOUMMEZZOUGH, A., PRIN, Y., BAUDOIN, E., GALIANA, A. & DREYFUS, B. (2011) Nurse shrubs increase the early growth of *Cupressus* seedlings by enhancing belowground mutualism and soil microbial activity. *Soil Biology & Biochemistry*, 43: 2160-2168.
- GUISSOU, T., BÂ, A.M., OUADBA, J.M., GUINKO, S. & DUPONNOIS, R. (1998) Responses of *Parkia biglobosa* (Jacq.) Benth, *Tamarindus indica* L. and *Zizyphus mauritiana* Lam. to arbuscular mycorrhizal fungi in a phosphorus-deficient sandy soil. *Biology and Fertility of Soils*, 26: 194-198.
- MANAUT, N., SANGUIN, H., OUAHMANE, L., BRESSAN, M., THIOULOUSE, J., BAUDOIN, E., GALIANA, A., HAFIDI, M., PRIN

- NEMEC, S. & VU, J.C.V. (1995)**
Effects of soil phosphorus and *Glomus intraradices* on growth, non-structural carbohydrates, and photosynthetic activity of *Citrus aurantium*. *Plant & Soil*, 128: 257-263.
- NIERING, W.A., WHITTAKER, R.H. & LOWE, C.H. (1963)**
The saguaro: a population in relation to environment. *Science*, 142: 15-23.
- REQUENA, N., PEREZ-SOLIS, E., AZCON-AGUILAR, C., JEFFRIES, P. & BAREA J.M. (2001)**
Management of indigenous Plant-Microbe Symbioses aids restoration of desertified ecosystems. *Applied & Environmental Microbiology*, 67: 495-498.
- RODRIGUEZ, A. & SANDERS, I.R. (2015)**
The role of community and population ecology in applying mycorrhizal fungi for improved food security. *The ISME Journal*, 9: 1053-1061.
- SCHLESINGER, W.H., RAIKES, J.A., HARTLEY, A.E. & CROSS, A.F. (1996)**
On the spatial pattern of soil nutrients in desert ecosystems. *Ecology*, 7: 364-374.
- SMITH, S.E. & READ, D.J. (2008)**
Mycorrhizal Symbiosis, third ed. Academic Press, London, UK.
- VAN DER HEIJDEN, M.G.A., KLIRONOMOS, J.N., URSIC, M., MOUTOGLIS, P., STREITWOLF-ENGEL, R., BOLLER, T., WIEMKEN, A. & SANDERS, I.R. (1998)**
Mycorrhizal fungal diversity determines plant biodiversity, ecosystem variability and productivity. *Nature*, 396: 72-75.
- WEBER, J., DUCOSSO, M., YEE THAM, F., NOURISSIER-MOUNTOU, S., GALIANA, A., PRIN, Y. & LEE, S.K. (2005)**
Co-inoculation of *Acacia mangium* with *Glomus intraradices* and *Bradyrhizobium* sp. in aeroponic culture. *Biology & Fertility of Soils*, 41: 233-239.

The Mediterranean Region under Climate Change

A Scientific Update

AllEnvi

Alliance nationale de recherche
pour l'Environnement

The Mediterranean Region under Climate Change

A Scientific Update

Preface by

Hakima EL HAÏÉ

Postface by

Driss EL YAZAMI

Address by

HSH the Prince ALBERT II of Monaco

IRD ÉDITIONS
INSTITUT DE RECHERCHE POUR LE DÉVELOPPEMENT

Marseille, 2016

Revision and translation

Daphne Goodfellow

Andrew Morris

Graphics

Michelle Saint-Léger

With the collaboration of:

Desk

Gris Souris

Layout

Desk

Cover layout

Michelle Saint-Léger

Page layout

Pierre Lopez

Coordination production

Catherine Plasse

Cover illustrations

© Météo France – RGB composite imagery, *METEOSAT-10*, 07/04/2016 at 12 UTC.

© IRD/B. Moizo – The town of Chefchaouen, Morocco.

© Ifremer/D. Lacroix – The port of Bizerte, Tunisia.

© IRD/J.-P. Montoroi – Olive trees, Seblet Ben Ammar, Tunisia.

La loi du 1^{er} juillet 1992 (code de la propriété intellectuelle, première partie) n'autorisant, aux termes des alinéas 2 et 3 de l'article L. 122-5, d'une part, que les « copies ou reproductions strictement réservées à l'usage du copiste et non destinées à une utilisation collective » et, d'autre part, que les analyses et les courtes citations dans le but d'exemple ou d'illustration, « toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'auteur ou de ses ayants droit ou ayants cause, est illicite » (alinéa 1^{er} de l'article L. 122-4).

Cette représentation ou reproduction, par quelque procédé que ce soit, constituerait donc une contrefaçon passible des peines prévues au titre III de la loi précitée.

© **IRD, 2016**

ISBN : 978-2-7099-2219-7

This book, coordinated by AllEnvi, is published on the occasion of the 22nd Conference of the Parties to the United Nations Framework Convention on Climate Change (COP22, Marrakech, 2016)

Scientific Direction

Stéphanie Thiébault
Jean-Paul Moatti

Scientific Committee

Isabella Annesi-Maesano	Véronique Ducrocq	Pascal Marty
Yildiz Aumeeruddy-Thomas	François Dulac	Yunne-Jai Shinne
Robert Barouki	Benoît Fauconneau	Jean-François Soussana
Gilles Boulet	Eric Gaume	Emmanuel Torquebiau
Jean-Luc Chotte	Jean-François Guégan	Jean-Denis Vigne
François Clin	Joël Guiot	
Wolfgang Cramer	Eric Hamonou	
Michel Crépon	Denis Lacroix	

Editorial Committee

Marie-Lise Sabrié
Elisabeth Gibert-Brunet
Thomas Mourier

AllEnvi

AllEnvi, the French National Alliance for Environmental Research, is tasked with making the great environmental transitions work, coordinating French research into major societal issues such as food, water, climate and territories. AllEnvi i) sets policy guidelines and research priorities for advance planning before approaching funding agencies, ii) supports the emergence and structuring of research organizations, iii) coordinates innovation and technology transfer policies between public research operators, businesses and industries, and iv) contributes to the European research environment and international programme development.

Alliance nationale de recherche pour l'environnement, AllEnvi coordonne la recherche française sur les enjeux des grands défis sociétaux que sont l'alimentation, l'eau, le climat et les territoires pour réussir les grandes transitions environnementales. AllEnvi i) définit les orientations et priorités de recherche pour la programmation à l'amont des agences de financement, ii) soutient l'émergence et la structuration d'infrastructures de recherche, iii) coordonne les politiques d'innovation et valorisation entre opérateurs publics de la recherche, entreprises et industries, et iv) participe à l'Europe de la recherche et favorise l'émergence de programmes internationaux.

Executive Secretary/Sécrétariat exécutif :

Benoit Fauconneau
Christine Douchez
Elisabeth Gibert-Brunet