



GAMBIERDISCUS TOXICUS AND CIGUATERA

M. DURAND-CLÉMENT, P. BOURDEAU* et Danielle PESANDO

*Unité 303 INSERM, BP 3, 06230 Villefranche sur Mer, France; * Laboratoire de Parasitologie, Maladies parasitaires, Zoologie appliquée, Ecole Nationale Vétérinaire de Nantes, BP 3013, 44087 Nantes Cedex 03, France*

In 1977, Yasumoto et al. found ciguatoxin-like and maitotoxin-like toxins in samples of algae containing large amounts of *Gambierdiscus toxicus*. Up to this date this benthic dinoflagellate is considered as the main toxin producer yielding to ciguateric fishes via the marine food chain.

We have conducted a study of *G. toxicus* populations at Saint Barthélémy from 1985 to 1987. Because this Caribbean island is highly affected by ciguatera fish poisoning, it is a good model for studying the link between *G. toxicus* toxicity and poisoning incidence.

We have found that *G. toxicus* was located all around St Barthélémy at low cell densities from 5 to 500 cells per gram of wet macrophytes, with perhaps a maximum of abundance in Spring. Photonic and electronic microscopy showed no difference in regard to Pacific specimens.


Using the mouse bioassay (intraperitoneal injection), toxicities of crude methanolic extracts (mostly MTX) were found to be very close (1000 cells of *G. toxicus* are necessary to kill a 20g mouse within 24 h.), whereas after separation on silicagel column, the toxicity of the chloroform-methanol (9:1) fraction (CTX-like) was higher in the caribbean sample (around 1% instead of 0,1%). This difference may be due either to a lost of toxicity of the Gambier strain during the culture period or to the presence of different bacterial communities. Research is in progress to elucidate this point.

Our study could be summarized by : - high incidence of ciguatera in both Gambier islands and St Barthélémy but -large cell density, low CTX-like toxicity in the former, -low cell density, higher CTX-like toxicity in the latter. Therefore both toxicity and abundance of *G. toxicus* should be taken into account for evaluating the role of *G. toxicus* in ciguateric areas.

Troisième Symposium sur les substances naturelles d'intérêt biologique de la région Pacifique-Asie

Nouméa, Nouvelle-Calédonie, 26-30 Août 1991

ACTES



Editeurs : Cécile DEBITUS, Philippe AMADE,
Dominique LAURENT, Jean-Pierre COSSON