



A uniform, quality controlled Surface Ocean CO₂ Atlas (SOCAT)

B. Pfeil^{1,2,3}, A. Olsen^{1,2,4,5}, D. C. E. Bakker⁶, S. Hankin⁷, H. Koyuk⁸, A. Kozyr⁹, J. Malczyk¹⁰, A. Manke⁷, N. Metzler¹¹, C. L. Sabine⁷, J. Akl^{12,13}, S. R. Alin⁷, N. Bates¹⁴, R. G. J. Bellerby^{15,16,2}, A. Borges¹⁷, J. Boutin¹¹, P. J. Brown^{6,18}, W.-J. Cai¹⁹, F. P. Chavez²⁰, A. Chen²¹, C. Cosca⁷, A. J. Fassbender²², R. A. Feely⁷, M. González-Dávila²³, C. Goyet²⁴, B. Hales²⁵, N. Hardman-Mountford^{26,*}, C. Heinze^{1,2,5,16}, M. Hood²⁷, M. Hoppema²⁸, C. W. Hunt²⁹, D. Hydes³⁰, M. Ishii³¹, T. Johannessen^{1,2}, S. D. Jones³², R. M. Key³³, A. Körtzinger³⁴, P. Landschützer⁶, S. K. Lauvset^{1,2}, N. Lefèvre¹¹, A. Lenton¹², A. Lourantou¹¹, L. Merlivat¹¹, T. Midorikawa³⁵, L. Mintrop³⁶, C. Miyazaki³⁷, A. Murata³⁸, A. Nakadate³⁹, Y. Nakano³⁸, S. Nakaoka⁴⁰, Y. Nojiri⁴⁰, A. M. Omar^{5,16}, X. A. Padin⁴¹, G.-H. Park⁴², K. Paterson^{12,13}, F. F. Perez⁴¹, D. Pierrot⁴², A. Poisson²⁴, A. F. Ríos⁴¹, J. M. Santana-Casiano²³, J. Salisbury²⁹, V. V. S. S. Sarma⁴³, R. Schlitzer²⁸, B. Schneider⁴⁴, U. Schuster⁶, R. Sieger²⁸, I. Skjelvan^{1,2,16}, T. Steinhoff³⁴, T. Suzuki⁴⁵, T. Takahashi⁴⁶, K. Tedesco^{47,**}, M. Telszewski^{48,**}, H. Thomas⁴⁹, B. Tilbrook^{12,13,50}, J. Tjiputra^{1,2}, D. Vandemark²⁹, T. Veness^{12,13}, R. Wanninkhof⁵¹, A. J. Watson⁶, R. Weiss⁵², C. S. Wong⁵³, and H. Yoshikawa-Inoue³⁸

¹Geophysical Institute, University of Bergen, Bergen, Norway

²Bjerknes Centre for Climate Research, Bergen, Norway

³PANGAEA Data Publisher for Earth & Environmental Science, University of Bremen, Bremen, Germany

⁴Institute of Marine Research, Bergen, Norway

⁵Uni Bjerknes Centre, Bergen, Norway

⁶School of Environmental Sciences, University of East Anglia, Norwich, UK

⁷Pacific Marine Environmental Laboratory, National Oceanic and Atmospheric Administration, Seattle, Washington, USA

⁸Joint Inst. for the Study of the Atmosphere and Ocean, University of Washington, Seattle, Washington, USA

⁹Carbon Dioxide Information Analysis Center, Oak Ridge, Tennessee, USA

¹⁰Jetz Laboratory, Department of Ecology and Evolutionary Biology, Yale University, New Haven, Connecticut, USA

¹¹Université Pierre et Marie Curie, LOCEAN/IPSL, Paris, France

¹²CSIRO Wealth from Oceans Flagship, Hobart, Tasmania, Australia

¹³Centre for Australian Weather and Climate Research, Hobart, Tasmania, Australia

¹⁴Bermuda Institute of Ocean Sciences, Ferry Reach, Bermuda

¹⁵Norwegian Institute for Water Research, Bergen, Norway

¹⁶Uni Research AS, Bergen, Norway

¹⁷University of Liège, Chemical Oceanography Unit, Institut de Physique, Liège, Belgium

¹⁸British Antarctic Survey, Cambridge, UK

¹⁹Department of Marine Sciences, University of Georgia, Athens, Georgia, USA

²⁰Monterey Bay Aquarium Research Institute, Moss Landing, California, USA

²¹Institute of Marine Geology and Chemistry, National Sun Yat-sen University, Kaohsiung, Taiwan

²²School of Oceanography, University of Washington, Seattle, Washington, USA

²³Univ. de Las Palmas de Gran Canaria, Facultad de Ciencias del Mar, Las Palmas de Gran Canaria, Spain

²⁴Inst. de Modélisation et d'Analyse en Géo-Environnement et Santé, Univ. de Perpignan, Perpignan, France

²⁵Oregon State University College of Oceanic and Atmospheric Sciences, Corvallis, USA

²⁶CSIRO, Marine and Atmospheric Research, Wembley, Western Australia, Australia

²⁷Intergovernmental Oceanographic Commission, UNESCO, Paris, France

²⁸Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, Germany

- ²⁹Ocean Process Analysis Lab, University of New Hampshire, Durham, New Hampshire, USA
³⁰National Oceanography Centre, Southampton, UK
³¹Japan Meteorological Agency, Meteorological Research Institute, Tsukuba, Japan
³²Tyndall Centre for Climate Change Research, Norwich, UK
³³Atmospheric and Oceanic Sciences, Princeton University, Princeton, New Jersey, USA
³⁴GEOMAR, Helmholtz Centre for Ocean Research, Kiel, Germany
³⁵Nagasaki Marine Observatory, Nagasaki, Japan
³⁶MARIANDA, Kiel, Germany
³⁷Faculty of Environmental Earth Science, Hokkaido University, Hokkaido, Japan
³⁸Japan Agency for Marine–Earth Science and Technology, Yokosuka, Japan
³⁹Marine Division, Global Environment and Marine Department, Japan Meteorological Agency, Tokyo, Japan
⁴⁰National Institute for Environmental Studies, Tsukuba, Japan
⁴¹Instituto de Investigaciones Marinas de Vigo, Consejo Superior de Investigaciones Científicas, Vigo, Spain
⁴²Cooperative Inst. for Marine and Atmospheric Studies, Rosenstiel School for Marine and Atmospheric Science, Univ. of Miami, Miami, Florida, USA
⁴³National Institute of Oceanography, Regional Centre, Visakhapatnam, India
⁴⁴Leibnitz Research Institute for Baltic Sea Research, Warnemünde, Germany
⁴⁵Marine Information Research Center, Japan Hydrographic Association, Tokyo, Japan
⁴⁶Lamont Doherty Earth Observatory, Palisades, New York, USA
⁴⁷International Ocean Carbon Coordination Project (IOCCP), Silver Spring, Maryland, USA
⁴⁸International Ocean Carbon Coordination Project (IOCCP), Inst. of Oceanology of Polish Academy of Sciences, Sopot, Poland
⁴⁹Department of Oceanography, Dalhousie University, Halifax Nova Scotia, Canada
⁵⁰Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart, Tasmania, Australia
⁵¹Atlantic Oceanographic and Meteorological Laboratory, National Atmospheric and Oceanographic Administration, Miami, Florida, USA
⁵²Scripps Institution of Oceanography, La Jolla, California, USA
⁵³Institute of Ocean Sciences, Sidney, British Columbia, Canada
*formerly at: Plymouth Marine Laboratory, Plymouth, UK
**formerly at: International Ocean Carbon Coordination Project (IOCCP), Intergovernmental Oceanographic Commission of UNESCO, Paris, France

Correspondence to: B. Pfeil (benjamin.pfeil@gfi.uib.no)

Received: 24 June 2012 – Published in Earth Syst. Sci. Data Discuss.: 14 August 2012

Revised: 6 February 2013 – Accepted: 20 February 2013 – Published: 4 April 2013

Abstract. A well-documented, publicly available, global data set of surface ocean carbon dioxide (CO₂) parameters has been called for by international groups for nearly two decades. The Surface Ocean CO₂ Atlas (SOCAT) project was initiated by the international marine carbon science community in 2007 with the aim of providing a comprehensive, publicly available, regularly updated, global data set of marine surface CO₂, which had been subject to quality control (QC). Many additional CO₂ data, not yet made public via the Carbon Dioxide Information Analysis Center (CDIAC), were retrieved from data originators, public websites and other data centres. All data were put in a uniform format following a strict protocol. Quality control was carried out according to clearly defined criteria. Regional specialists performed the quality control, using state-of-the-art web-based tools, specially developed for accomplishing this global team effort. SOCAT version 1.5 was made public in September 2011

and holds 6.3 million quality controlled surface CO₂ data points from the global oceans and coastal seas, spanning four decades (1968–2007). Three types of data products are available: individual cruise files, a merged complete data set and gridded products. With the rapid expansion of marine CO₂ data collection and the importance of quantifying net global oceanic CO₂ uptake and its changes, sustained data synthesis and data access are priorities.

Data coverage

Repository-Reference: doi:10.1594/PANGAEA.767698
 Available at: www.socat.info
 Coverage: 80° N to 79° S and 0–360°
 Location Name: Global Ocean
 Date/Time Start: 16 November 1968
 Date/Time End: 31 December 2007

1 Motivation

The net absorption of CO₂ by the oceans, caused by rising atmospheric CO₂ concentrations since the industrial revolution, has been responsible for removing CO₂ equivalent to approximately 50% of the fossil fuel and cement manufacturing emissions or about 30% of the total anthropogenic emissions, including land use change (Sabine et al., 2004). Because of the availability of the carbonate ion, an important species of the dissolved inorganic carbon pool, and carbonate sediments, the oceans have a tremendous CO₂ uptake capacity and will, on timescales of ten to hundred thousand years, absorb all but a small fraction of the fossil CO₂ that has been and will be emitted (Archer et al., 1997). Meanwhile the changes in ocean CO₂ uptake, relying on factors such as ocean circulation and biology, will be among the decisive factors for the evolution of future atmospheric CO₂ concentrations and climate development (e.g., Friedlingstein et al., 2006; Riebesell et al., 2009).

Presently there are two types of globally coordinated efforts that seek to resolve the dynamics of ocean CO₂ uptake through observations: repeat hydrography and surface ocean CO₂ observations (Gruber et al., 2010; Sabine et al., 2010). While repeat hydrography aims to assess variations in the ocean inventory of CO₂ on decadal timescales, surface ocean observations may resolve variations on seasonal to interannual timescales due to the higher sampling frequency. This high sampling frequency has been made possible by the advent of autonomous instruments and sensors for the near-continuous determination of surface water CO₂, which may be installed on commercial seagoing vessels giving an observational repeat rate of a few weeks, depending on ship schedule (Cooper et al., 1998; Pierrot et al., 2009), or on moorings (Merlivat and Brault, 1995; DeGrandpré et al., 2000; Friederich et al., 2008; Wada et al., 2011). Moorings or drifting platforms provide observations on sub-diurnal timescales (e.g., Körtzinger et al., 2008; Leinweber et al., 2009; Merlivat et al., 2009; Parard et al., 2010), while underway observations increase spatial coverage.

These technological developments have led to a rapid increase in new surface ocean CO₂ data being collected each year. This is reflected in the number of data underlying the successive surface ocean *p*CO₂ (partial pressure of CO₂) climatologies of Takahashi et al. (1997, 2002, 2009a, b, 2011), increasing from 0.25 million for the 1997 edition to 5.2 million in 2011. Presently over a million observations are being made each year (Sabine et al., 2010). In order to deal with these data effectively and to maximise their scientific use, the international ocean carbon research community initiated the Surface Ocean CO₂ Atlas (SOCAT) project in 2007 (IOCCP, 2007). The aims of SOCAT were threefold. Firstly, SOCAT aimed to merge all available surface ocean CO₂ data into one uniformly formatted, quality controlled, publicly available database with regular updates. The second aim of SOCAT was to secure the long-term storage of each data set




Figure 1. Number of data points per year during the period 1968 to 2007 included in SOCAT version 1.5.

together with its required documentation (metadata). Finally, the community sought to realise a transparent and traceable approach for the handling, quality control and integration of surface ocean CO₂ data, which may be managed by the community on a routine basis in the future.

The first version of SOCAT (version 1.5) was made public on 14 September 2011 during “The ocean carbon cycle at a time of change: Synthesis and Vulnerabilities” meeting at the UNESCO (United Nations Educational, Scientific and Cultural Organization), Paris (Bakker et al., 2012). This SOCAT version comprises 6.3 million surface water CO₂ data from 1851 voyages from 1968 to 2007 covering the global oceans and coastal seas (Figs. 1 and 2). Three data products are available: (1) cruise data files of quality controlled surface water *f*CO₂ (fugacity of CO₂, similar to partial pressure) data and including the reported CO₂ values as reported by the investigator, (2) globally and regionally aggregated files of these *f*CO₂ data, and (3) a collection of gridded products providing averaged *f*CO₂ with minimal interpolation (Sabine et al., 2013). This article describes the history of SOCAT (Sect. 2), the procedures adopted in SOCAT for retrieving data (Sect. 3), for formatting (Sect. 3) and quality controlling these data (Sect. 4). The article introduces SOCAT data products and where they can be accessed (Sect. 5). An accompanying article (Sabine et al., 2013) describes the gridding procedures. The SOCAT website (www.socat.info) provides documentation on SOCAT, as well as links to sites with SOCAT data products. This article concludes with lessons learned from this first SOCAT version and recommendations for future SOCAT releases (Sect. 6).




Figure 2. Distribution of data in SOCAT version 1.5 per decade: (a) 1970s, (b) 1980s, (c) 1990s and (d) 2000s.

2 History and organisation of SOCAT

2.1 History of SOCAT

In the late 1990s attempts were made by the SCOR-IOC (Scientific Committee on Oceanic Research-Intergovernmental Oceanographic Commission) committee on ocean CO₂, the forerunner of the IOCCP (International Ocean Carbon Coordination Project), to assemble a comprehensive, well documented, publicly available data set of surface ocean *f*CO₂ for the global oceans and coastal seas. Efforts for encouraging data submission to a central location, the Carbon Dioxide Information Analysis Center, were partly successful. In 2004 the marine carbon community agreed on recommendations for the reporting of surface water CO₂ data and metadata (IOCCP, 2004). However, most data gatherers did not strictly follow these. Only a subset of all global surface water CO₂ data were made publicly available via CDIAC, with many data only available via the investigators, institute websites and national or world data centres.

Over the past decades several attempts have been made to establish a global surface ocean CO₂ database. In the late 1990s, Taro Takahashi from Lamont-Doherty Earth Observatory (LDEO) compiled an initial data set and updated this collection in 2002 and every year from 2007 onwards (Takahashi et al., 1997, 2002, 2009a, 2011). The primary reason

for this effort was the creation of global climatologies of air-sea CO₂ fluxes (Takahashi et al., 1997, 2002, 2009b). This LDEO database was made public in 2007 and is currently being updated on an annual basis. The data treatment is based upon Takahashi's long experience. The LDEO database includes *p*CO₂ from discrete and continuous measurements. The most recent version of the LDEO data set has 5.2 million *p*CO₂ data from the global oceans and coastal seas from 1957 to 2010 (Takahashi et al., 2011).

In 2001, Bakker began to assemble a surface ocean CO₂ data set by putting public data from CDIAC into a uniform format, as part of the European Union (EU) project ORFOIS (Origin and fate of biogenic particle Fluxes in the Ocean and their Interaction with the atmospheric CO₂ concentration as well as the marine Sediment). Pfeil and Olsen streamlined and expanded this effort within the EU project CarboOcean from 2005 onwards. They compiled public surface ocean CO₂ data held at CDIAC, PANGAEA – Data Publisher for Earth & Environmental Science (an International Council for Science (ICSU) World Data Center, formerly the World Data Center for Marine Environmental Sciences, WDC-MARE) and elsewhere into a common format *f*CO₂ database based on the recommended formats for data and metadata reporting (IOCCP, 2004).

Table 1. Meetings for SOCAT version 1.5. Abbreviations are explained in the text and acknowledgements.

Timing	Meeting description	Location	Reference
04/2007	Surface Ocean CO ₂ Variability and Vulnerability (SOCOVV) workshop. Initiation of SOCAT.	UNESCO, Paris, France	IOCCP (2007)
12/2007	First technical meeting. Discussion of procedures.	Delmenhorst and Bremen, Germany	NA
06/2008	Second technical meeting. Discussion of data inclusion, QC flags, LAS-based data QC.	UNESCO, Paris, France	IOCCP (2008)
01/2009	SOCAT Coastal regional workshop	GEOMAR, Kiel, Germany	IOCCP (2009a)
03/2009	SOCAT Pacific regional workshop	NIES, Tsukuba, Japan	IOCCP (2010a)
06/2009	SOCAT Atlantic and Southern Ocean regional workshop	UEA, Norwich, UK	IOCCP (2009b)
02/2010	SOCAT Equatorial Pacific, North Pacific and Indian Ocean regional workshop	Tokyo, Japan	NA
06/2010	SOCAT Southern Ocean and Indian Ocean regional workshop	CSIRO, Hobart, Australia	IOCCP (2010b)
09/2011	The Ocean Carbon Cycle at a Time of Change: Synthesis and Vulnerabilities. Public release of SOCAT version 1.5.	UNESCO, Paris, France	IOCCP (2013)

Table 2. Groups and key participants in SOCAT. Figure 3 shows the SOCAT regions.

Group	Area	Lead(s) and Key Participants
Global		Bakker (chair), Olsen, Pfeil, Hankin, Koyuk, Kozyr, Malzcyk, Metzl, Pierrot, Sabine, Telszewski
Coastal regions	< 400 km from land; north of 30° S	Borges, Chen
North Atlantic	north of 30° N, incl. Atlantic Arctic	Schuster
Tropical Atlantic	30° N to 30° S	Lefèvre
North Pacific	north of 30° N, incl. Pacific Arctic	Nojiri
Tropical Pacific	30° N to 30° S	Feely
Indian Ocean	north of 30° S	Sarma
Southern Ocean	south of 30° S, incl. coastal waters	Tilbrook, Metzl

The Surface Ocean CO₂ Atlas was initiated at the Surface Ocean CO₂ Variability and Vulnerability (SOCOVV) meeting by the international ocean carbon research community (Table 1) (IOCCP, 2007). The SOCAT project agrees well with the objectives of the joint Carbon Implementation Plan of the Surface Ocean Lower Atmosphere Study (SOLAS) and Integrated Marine Biogeochemistry and Ecosystem Research (IMBER) (IMBER, 2005). SOCAT was given the specific objectives of developing two data products (IOCCP, 2007):

- A quality controlled $f\text{CO}_2$ data set made publicly available on a regular basis following agreed procedures and regional review;
- A gridded product consisting of monthly surface $f\text{CO}_2$ means (including number of data points and standard deviation) on a 1° latitude by 1° longitude grid with no interpolation.

A gridded surface ocean $f\text{CO}_2$ product was deemed to be more useful than air–sea CO₂ flux estimates for modelling and other purposes (IOCCP, 2007). Regional groups and a global group for coordination were formed (Table 2).




Figure 3. Oceanic and coastal regions in SOCAT (Table 1). The white collar surrounding land masses indicates the coastal regions. Coastal regions north of 30° S were quality controlled by the coastal group, while coastal regions south of 30° S were quality controlled by the Southern Ocean group.

A series of meetings was held in which SOCAT gradually took shape and in which the regional groups coordinated their work (Table 1) (IOCCP, 2007, 2008, 2009a, b, 2010a, b).

The SOCAT community evaluated existing data compilations and selected the data collection by Pfeil and Olsen as the basis for SOCAT (IOCCP, 2008). The focus for SOCAT has been the assembly of publicly available data (including metadata), standardisation of the file formats, recalculation of consistent and uniform surface water $f\text{CO}_2$ data, and basic and secondary level quality control (Sects. 3, 4 and 5).

SOCAT is independent from the LDEO database (Takahashi et al., 2011), but has a large overlap in its original data. SOCAT only includes surface water CO₂ values, measured in near-continuous operation or in discrete samples with an equilibrator system or a spectrophotometer and reported as $x\text{CO}_2$ (CO₂ mixing ratio), $p\text{CO}_2$ or $f\text{CO}_2$ (Sect. 3). SOCAT does not include $f\text{CO}_2$ recalculated from dissolved inorganic carbon, alkalinity or pH.

2.2 SOCAT groups

Roughly 45 international, seagoing marine carbon scientists and data managers from 12 countries actively participated in the assembly and quality control of SOCAT version 1.5. These participants were organised into regional groups and a global group (Table 2). The regional groups were responsible for quality control of the data in their region. Regional groups were formed for the coastal seas (north of 30° S), the North Atlantic Ocean (north of 30° N, including the Arctic Ocean), the Tropical Atlantic Ocean (30° N to 30° S), the North Pacific Ocean (north of 30° N, including the Pacific

Arctic Ocean), the Tropical Pacific Ocean (30° N to 30° S), the Indian Ocean (north of 30° S) and the Southern Ocean (south of 30° S, including coastal waters). Coastal regions were initially defined by bathymetry (shallower than 200 m) for regions north of 30° S (IOCCP, 2008). This definition was later replaced by a criterion of distance from a major land mass (less than 400 km) in order to better reflect the environmental significance of these regions as continental margins. Figure 3 shows these oceanic and coastal regions in SOCAT.

SOCAT has been a large, complex undertaking and has involved activities focused on: data retrieval, assembling data in a uniform format, recalculating surface water $f\text{CO}_2$ using the same agreed-upon protocol, defining SOCAT QC criteria, developing the QC cookbook and Matlab QC code, making SOCAT available via the Live Access Server (LAS) for QC and public release, data QC, gridding SOCAT, making SOCAT documentation and products available via the web, designing the SOCAT logo, internal communication, organisation of SOCAT meetings, and liaising with the international marine carbon community. Numerous colleagues have played a role in these activities (Table 3). The SOCAT global group initially had five members and has gradually been expanded to reflect the increasing complexity of the tools and products in SOCAT (Table 2).

3 SOCAT data assembly

3.1 Data sources and instrumentation

SOCAT includes 6.3 million $f\text{CO}_2$ data points measured in all ocean areas from 1968 to 2007. Most of these data were gathered from the online sources at CDIAC (30 % of the cruises) and PANGAEA (10 %), as well as from institute and project websites (37 %). The remaining cruises (23 %) were obtained directly from the data originators. Almost half of the cruises (45.7 %) originated in the USA. Other significant contributors are based in Japan (20.1 %), Norway (9.6 %), the United Kingdom (7.4 %), Germany (5.8 %), France (4.5 %), Belgium (2.4 %), Canada (1.6 %), Spain (1.5 %), Australia (1.2 %) and the Netherlands (0.3 %).

The data in SOCAT are a synthesis of 4 decades of seagoing fieldwork by numerous scientists from 12 countries. Various instruments have been used to obtain these data and only the basic principles will be summarised here. Further information is available in the metadata, which accompany individual cruise files at PANGAEA (doi:10.1594/PANGAEA.769638) (Sect. 5.2).

The seawater $f\text{CO}_2$ values included in SOCAT have been measured according to one of the following two principles: (1) analysis of the CO₂ content in an air sample in equilibrium with a large volume of seawater or (2) calculation of the seawater $f\text{CO}_2$ from the colour response of an acid-base indicator dye (sulfonephthalein) in contact with seawater across a CO₂ permeable membrane. The analysis of the CO₂ content in an air sample in equilibrium with a large volume

of seawater is recommended in the standard work by Dickson et al. (2007). The CO₂ concentration in the air sample is determined through either gas chromatography (Weiss et al., 1981) or infrared analysis (Takahashi, 1961). The equilibration of air and water can be carried out in an equilibrator in a flow-through system (Takahashi, 1961; Wanninkhof and Thoning, 1993; Cooper et al., 1998; Pierrot et al., 2009). Only the latter (i.e., continuous data) are included in SOCAT. Flow-through systems combined with a non-dispersive infrared (IR) detector are by far the most common type in operation. Flow-through systems are routinely deployed on commercial vessels (e.g., Cooper et al., 1998; Olsen et al., 2008; Watson et al., 2009), research vessels (e.g., Lefèvre et al., 1994; Skjelvan et al., 1999; Bakker et al., 2008), and on moored platforms (e.g., Friederich et al., 2008; Wada et al., 2011). Intercomparison experiments have taken place on a number of occasions (e.g., Körtzinger et al., 1996, 2000).

The indicator-based, spectrophotometric determination of *f*CO₂ has been developed for moored and drifting platforms (Lefèvre et al., 1993). Prominent examples of these are the CARIOCA (Carbon Interface Ocean Atmosphere) buoy (Merlivat and Brault, 1995) and the SAMI (Submersible Autonomous Moored Instrument) *p*CO₂ instrument (DeGrandpré et al., 2000). These instruments have been deployed in many ocean regions (e.g., Hood et al., 1999; Bakker et al., 2001; Körtzinger et al., 2008; Lefèvre et al., 2008).

3.2 Data harmonisation and basic quality control

All data files available for SOCAT were first converted to a common file structure. This also included discarding data not directly relevant for surface ocean CO₂, e.g., meteorological parameters like wind speed and direction, whenever these were supplied in the file. Next, the unit of each parameter was checked and converted into the agreed standard unit, if required (e.g., conversion of atmospheric pressure from atmospheres to hPa, and of latitude and longitude to decimal degrees). For around 10 % of the cruises, different versions of the data had been obtained from various sources. In these cases only the most recent version was included in SOCAT in consultation with the data originator.

Basic, primary quality control was carried out at this stage. Outliers and unrealistic values in date, time, position, intake temperature, salinity, atmospheric pressure and surface water CO₂ were identified. The criteria were that ship speeds calculated from position should be realistic, that atmospheric pressures should be within 800 hPa and 1100 hPa and that the dates should exist. Rapid changes in intake or equilibrator temperature of several degrees, in salinity of several units or in surface *f*CO₂ of several hundreds of micro-atmospheres were also questioned, except for data in coastal or ice-covered regions. Whenever several such data points were encountered, the data originator was contacted and this often resulted in resubmission of an updated (corrected) ver-

sion. In some cases several iterations were required, making this a time-consuming task. In a few cases interaction with the data originator was not possible, and obviously bad data were removed from the data file.

In version 2 of SOCAT this class of quality control will be used to assign quality flags to individual data points, using the conventions of the World Ocean Circulation Experiment (WOCE): flag 2 (good), flag 3 (questionable) or 4 (bad). Only a very small number of WOCE flags 3 and 4 are found in the version 1.5 data collection.

3.3 *f*CO₂ (re-)calculations

The final stage of the SOCAT data assembly was the (re-)calculation of *f*CO₂ values at sea surface (or intake) temperature in order to ensure a uniform representation of CO₂ concentration. The conversions from *x*CO₂ and *p*CO₂ were carried out using a single set of equations with a clear hierarchy for the preferred CO₂ input parameter (Table 4) (Pfeil and Olsen, 2009). We used the equations recommended by Dickson et al. (2007),

$$p\text{CO}_2 = x\text{CO}_{2T_{\text{equ}}}^{\text{dry}} (P_{\text{equ}} - p\text{H}_2\text{O}), \quad (1)$$

for the conversion of dry CO₂ mole fraction to partial pressure at 100 % humidity, where P_{equ} is the pressure in the equilibrator. The water vapour pressure $p\text{H}_2\text{O}$ is calculated as

$$p\text{H}_2\text{O} = \exp\left(24.4543 - 67.4509\left(\frac{100}{TK_{\text{equ}}}\right) - 4.8489 \ln\left(\frac{TK_{\text{equ}}}{100}\right) - 0.000544S\right), \quad (2)$$

where TK_{equ} is the measurement (or equilibrator) temperature in Kelvin and S is sample salinity. For the conversion of *p*CO₂ values into *f*CO₂ the equation is

$$f\text{CO}_2 = p\text{CO}_2 \times \exp\left\{\frac{\left[B(\text{CO}_2, TK_{\text{equ}}) + 2(1 - x\text{CO}_{2T_{\text{equ}}}^{\text{wet}})^2 \delta(\text{CO}_2, TK_{\text{equ}})\right] \times P_{\text{equ}}}{R \times TK_{\text{equ}}}\right\}, \quad (3)$$

where $x\text{CO}_{2T_{\text{equ}}}^{\text{wet}}$ is the wet CO₂ mole fraction at the equilibrator temperature. The virial coefficients for CO₂, $B(\text{CO}_2, TK_{\text{equ}})$ and $\delta(\text{CO}_2, TK_{\text{equ}})$ (cm³ mol⁻¹), are given by

$$B(\text{CO}_2, TK_{\text{equ}}) = -1636.75 + 12.0408TK_{\text{equ}} - 3.2795710^{-2}TK_{\text{equ}}^2 + 3.1652810^{-5}TK_{\text{equ}}^3 \quad (4)$$

$$\delta(\text{CO}_2, TK_{\text{equ}}) = 57.7 - 0.118TK_{\text{equ}}. \quad (5)$$

Whenever conversion of the measurement (equilibrator) temperature (T_{equ}) to the sea surface temperature (SST) was required, we used the equation of Takahashi et al. (1993) with both temperatures in the same unit:

$$f\text{CO}_2^{\text{SST}} = f\text{CO}_2^{\text{equ}T} \exp(0.0423(\text{SST} - T_{\text{equ}})). \quad (6)$$

Table 3. Activities and key participants in SOCAT.

Activity	Key Participants
Data retrieval	Pfeil, Hood, Bakker
Data ingestion	Pfeil
Uniform format	Olsen, Pfeil
Defining QC (quality control) criteria	Wanninkhof, Olsen, Schuster, meeting participants (IOCCP, 2008, 2009b)
QC cookbook	Olsen, Metzl
Coastal mask	Hales, Olsen, Hankin
Matlab [®] QC code	Pierrot, Olsen
Live Access Server (LAS)	Hankin, Malczyk, Koyuk
Data QC	SOCAT regional groups
Conflicting cruise flags	Bakker
Gridding	Sabine, Fassbender, Manke
Logo	Brown
SOCAT webpage	Pfeil, Koyuk, Bakker
Online SOCAT products	Kozyr, Koyuk, Hankin, Pfeil
ODV (Ocean Data View) for SOCAT	Schlitzer
SOCAT meetings	Nojiri, Borges, Wallace, Schuster, Bakker, Tilbrook, Hood, Tedesco, Telszewski, Brévière, Maddison
Internal coordination	Hood, Bakker, Koyuk
Coordination with marine community	Hood, Tedesco, Telszewski

Table 4. Surface water CO₂ parameters reported in the original data files, which have been used for the calculation of recommended *f*CO₂ (*f*CO₂_rec) at sea surface (or intake) temperature (Pfeil and Olsen, 2009). The parameters are listed in order of preference (with index 1 as the favourite). The index has been reported in the SOCAT global and regional output files as “fCO₂_source” (Table 5). Ancillary parameters have been used for NCEP (National Centers for Environmental Prediction) atmospheric pressure (Kalnay et al., 1996) and WOA (World Ocean Atlas) salinity (Antonov et al., 2006) (Sect. 3.3) in cases of incomplete data reporting.

Index	Reported CO ₂ parameter	Unit	Data percentage (%)	Extra variable
1	xCO ₂ water_equi_dry	μmol mol ⁻¹	57.5	
2	xCO ₂ water_SST_dry	μmol mol ⁻¹	6.6	
3	pCO ₂ water_equi_wet	μatm	6.4	
4	pCO ₂ water_SST_wet	μatm	2.9	
5	fCO ₂ water_equi	μatm	0.3	
6	fCO ₂ water_SST_wet	μatm	8.4	
7	pCO ₂ water_equi_wet ¹	μatm	0.4	NCEP Pressure
8	pCO ₂ water_SST_wet ¹	μatm	13.8	NCEP Pressure
9	xCO ₂ water_equi_dry ²	μmol mol ⁻¹	0.2	WOA Salinity
10	xCO ₂ water_SST_dry ²	μmol mol ⁻¹	1.2	WOA Salinity
11	xCO ₂ water_equi_dry ¹	μmol mol ⁻¹	0.0*	NCEP Pressure
12	xCO ₂ water_SST_dry ¹	μmol mol ⁻¹	2.2	NCEP Pressure
13	xCO ₂ water_equi_dry ^{1,2}	μmol mol ⁻¹	0.0*	NCEP Pressure, WOA Salinity
14	xCO ₂ water_SST_dry ^{1,2}	μmol mol ⁻¹	0.2	NCEP Pressure, WOA Salinity

¹ Atmospheric pressure was not reported in the original data file.² Salinity was not reported in the original data file.

* Not used for data reporting as an approach with a lower index was available.

Table 5. Content of the individual data cruise files (version 1.4) and the global and regional concatenated files (version 1.5) in SOCAT¹ (Sects. 5.1, 5.2 and 5.3).

Content of the individual cruise data files, version 1.4 ¹	Content of the regional and global synthesis files, version 1.5 ¹	Unit	Description
Event	cruise_ID	–	Expocode
Campaign	cruise_name	–	Cruise name*
Date/Time	NA	–	yyyy-mm-dd:hh:mm (ISO 8601) ¹
NA	Yr	Year	Year
NA	Mon	Month	Month
NA	Day	Day	Day
NA	Hour	Hour	Hour (GMT/UTC)
NA	Min	Minute	Minute ¹
Longitude	Longitude	° E	Longitude (0 to 360)*
Latitude	Latitude	° N, ° S	Latitude (–90 to 90)*
Depth Water	DepthW	m	Intake depth* ²
Temp	Temp	°C	Sea surface temperature*
Sal	Salinity	–	Sea surface salinity*
T_{equ}	Temperature_equi	°C	Temperature at equilibrator*
PPPP	Pressure_atm	hPa	Atmospheric pressure*
P_{equ}	Pressure_equi	hPa	Pressure in the equilibrator*
Sal interp	woa_sss	–	Salinity from WOA 2005
Press atmos interp	ncep_slp	hPa	Atmospheric pressure from 6-hourly NCEP/NCAR data
Bathy depth interp	ETOPO2	m	Bottom depth from ETOPO2 (2006)
xCO ₂ water_equ_dry*	–	μmol mol ^{–1}	xCO ₂ (water) at equilibrator temperature (dry air)*
fCO ₂ water_SST_wet*	–	μatm	fCO ₂ (water) at sea surface temperature (wet air)*
pCO ₂ water_SST_wet*	–	μatm	pCO ₂ (water) at sea surface temperature (wet air)*
xCO ₂ water_SST_dry*	–	μmol mol ^{–1}	xCO ₂ (water) at sea surface temperature (dry air)*
fCO ₂ water_equ_wet*	–	μatm	fCO ₂ (water) at equilibrator temperature (wet air)*
pCO ₂ water_equ_wet*	–	μatm	pCO ₂ (water) at equilibrator temperature (wet air)*
fCO ₂ water_SST_wet	fCO ₂ _rec	μatm	Recommended fCO ₂ , calculated for the SOCAT protocol (Table 3)
–	fCO ₂ _source	–	The algorithm for calculating fCO ₂ _rec (Index in Table 3)
–	gvco2	μmol mol ^{–1}	Atmospheric xCO ₂ from GLOBALVIEW-CO2 (2008)
–	Julian_day_GMT	Day	Day of year (1 for 1 January)
–	WOCE_flag	–	WOCE flag for fO ₂ _rec
Origin of values	doi	–	Digital object identifier to the individual cruise file and metadata
–	Averaged	–	Indicator that data was averaged for version 1.5 ¹

* refers to data reported by the data originator.

¹ Individual cruise data files in version 1.4 may contain multiple entries for a given time stamp. Multiple entries for a given time stamp have been averaged in the global and regional concatenated files in version 1.5 (Sects. 5.1, 5.2 and 5.3).

² If the intake depth has not been reported by the data originator, we assume an intake depth of 5 m.

The Takahashi et al. (1993) temperature correction is preferred, as it does not require knowledge of the alkalinity and dissolved inorganic carbon content of the water and was determined for isochemical conditions, while other temperature corrections (Gordon and Jones, 1973; Weiss et al., 1982; Copin-Montégut, 1988, 1989; Goyet et al., 1993) were not.

Altogether 6 different surface ocean CO₂ parameters were reported by the data originators, notably $x\text{CO}_2$, $p\text{CO}_2$ and $f\text{CO}_2$, either at sea surface (or intake) temperature or at equilibrator (or measurement) temperature (Table 4). The (re-)calculations of $f\text{CO}_2$ at sea surface temperature were implemented following these strict guidelines:

1. Whenever possible, (re-)calculate $f\text{CO}_2$.
2. The preferred starting point for the calculations is $x\text{CO}_2$, next $p\text{CO}_2$, and finally $f\text{CO}_2$.
3. Minimize the use of external data required to complete the calculations.

Thus, $f\text{CO}_2$ was recalculated if $x\text{CO}_2$, $p\text{CO}_2$, and $f\text{CO}_2$, as well as all parameters required to calculate $f\text{CO}_2$ were available in the file. However, $f\text{CO}_2$ was not recalculated if $f\text{CO}_2$ was reported, but pressure or salinity were not, as Eqs. (1), (2) and (3) could not be applied without resorting to external data. If only surface water $f\text{CO}_2$ at sea surface temperature was provided (as is the case for CARIOCA data and other spectrophotometric measurements), no recalculation was carried out. If $f\text{CO}_2$ was not provided, $f\text{CO}_2$ was always calculated, even if use of external data was necessary. Table 4 lists the parameters that went into the $f\text{CO}_2$ calculations and the preference (or hierarchy) of the different calculation methods. The $f\text{CO}_2$ values, which have been (re-)calculated following the preferred method (lowest index number in Table 4), are reported as the recommended $f\text{CO}_2$ ($f\text{CO}_2\text{_rec}$) values in each SOCAT output file (Table 5). The calculation method is indicated (as $f\text{CO}_2\text{_source}$) in the regional and global synthesis files of SOCAT version 1.5 (Table 5).

Two external parameters were used for the recalculations of $f\text{CO}_2$, when necessary: climatological monthly mean salinity was obtained from the World Ocean Atlas (WOA) 2005 (Antonov et al., 2006). Sea level pressure (SLP) was acquired from the NCEP/NCAR (National Centers for Environmental Prediction/National Center for Atmospheric Research) project (Kalnay et al., 1996), provided on a 6 hourly, global, 2.5° latitude by 2.5° longitude grid. Whenever NCEP/NCAR SLP or reported atmospheric pressure was used in the calculations (as opposed to equilibrator pressure), 3 hPa were added to account for the slight overpressure normally maintained in ships (Takahashi et al., 2009b). Surface water CO₂ data without accompanying SST were suspended from SOCAT, as $f\text{CO}_2$ is highly sensitive to temperature fluctuations.

3.4 Naming convention

Each cruise was assigned a unique cruise identifier, an Expocode (Swift, 2008), to remove the ambiguities of the commonly used informal cruise names and to identify duplicate versions of data. The first two characters of a twelve-character Expocode identify the country code of the vessel and are followed by the two-character National Oceanographic Data Center (NODC) vessel code. The final eight characters denote the starting date of the measurements of the cruise (as YYYYMMDD). For instance, *06MT19920510* means that this cruise was conducted on the German (06) research vessel *Meteor* (MT) and that the first measurement was reported for 10 May 1992. Both the Expocode and the original cruise name are provided in all SOCAT output files (Sect. 5), such that cruises can be retrieved using the Expocode as well as the vessel specific or investigator specific naming convention (M21/3 for the above example). The Expocode has not been used for buoys, since no NODC vessel code is available for these.

4 SOCAT secondary quality control

An important aim of SOCAT was to establish and implement community agreed secondary quality control (QC) procedures for $f\text{CO}_2$ data. Procedures for secondary QC were established at several SOCAT workshops (IOCCP, 2008, 2009b, 2010b) and were summarized in the SOCAT QC Cookbook (Olsen and Metzl, 2009). Secondary quality control was carried out by the SOCAT regional groups. The following sections provide an overview on the secondary quality control procedures in SOCAT and their implementation.

4.1 SOCAT secondary quality control procedures

Secondary quality control was carried out in SOCAT by assigning a quality flag to each cruise. The cruise flags provide information on the expected quality of the $f\text{CO}_2$ data in the different cruises. These are based on (i) an evaluation of the procedures and instruments used to measure the data, (ii) the availability of documentation enabling this evaluation, i.e., metadata, (iii) (whenever possible) a comparison with other data collected in the same region in the same period, and (iv) an assessment of data quality. The cruise flags and the formal criteria used to assign them are provided in Table 6. Only cruises with cruise flags A, B, C or D are included in the SOCAT products (Sect. 5).

In order to achieve cruise flag A, the data had to be accompanied by “complete metadata documentation”, the measurement techniques had to follow “approved methods or SOP criteria” (Standard Operating Procedures), extended QC had to be carried through and deemed acceptable, and the data would have to reasonably compare to other data from the same region. Moving from the A through to the D flag implies that the data did not meet with one or several of these

Table 6. Criteria for assigning cruise flags, based on the expected quality of the recommended $f\text{CO}_2$ data (revised after Olsen and Metzl, 2009). All criteria need to be met for assigning a cruise flag. SOP is Standard Operating Procedures (Dickson et al., 2007). QC is quality control.

Cruise flag (ID)	Criteria
A (11)	1. Followed approved methods or SOP criteria and 2. Metadata documentation complete and 3. Extended QC was deemed acceptable and 4. A comparison with other data was deemed acceptable.
B (12)	1. Followed approved methods or SOP criteria and 2. Metadata documentation complete and 3. Extended QC was deemed acceptable.
C (13)	1. Did not follow approved methods or SOP criteria and 2. Metadata documentation complete and 3. Extended QC was deemed acceptable (including comparison with other data if possible).
D (14)	1. Did or did not follow approved methods or SOP criteria and 2. Metadata documentation incomplete and 3. Extended QC was deemed acceptable (including comparison with other data if possible).
S (Suspend)	1. Did or did not follow methods or SOP criteria and 2. Metadata documentation complete or incomplete and 3. Extended QC revealed non-acceptable data 4. Data are being updated.
X (15) (Exclude)	The cruise (data set) duplicates another cruise (data set) in SOCAT.
N (No flag)	No cruise flag has yet been given to this cruise.
U (Update)	The cruise data have been updated. No cruise flag has yet been given to the revised data.

criteria. Hence, if the data were found to be sufficiently documented, obtained according to approved methods, and the data quality was deemed acceptable, but the data could not be compared to other data from the same region (since no other data were available), they would be assigned cruise flag B. If the sampling techniques did not follow approved methods, a flag C was assigned, and if the metadata documentation was incomplete, the data were assigned a cruise flag D. In addition to the A to D flags, it was intended that a flag F should be assigned to cruises for which the extended QC revealed that the data were non-acceptable. In practice, however, such cruises were suspended (Flag S). The flag S was assigned to cruises which were suspended with the aim to update the cruise data in future (often by the individual PIs after SOCAT QC revealed issues that could be fixed) and the flag X was assigned to data that were identified as duplicates of other data in SOCAT. To streamline the workflow we also used flags N, for newly added cruises with no cruise flag yet, and U, for cruises that had been updated.

4.1.1 Approved methods or SOP criteria

By approved methods or SOP criteria, required for flags A and B, we mean the recommendations of a 2002 workshop on underway $f\text{CO}_2$ systems (Atlantic Oceanographic and Mete-

orological Laboratory, 2002), as well as those of Dickson et al. (2007). Adhering to these methods results in $f\text{CO}_2$ data with an accuracy of 2 μatm or better (Olsen and Metzl, 2009). Seven SOP criteria need to be fulfilled for a cruise flag A or B in SOCAT:

1. The data are based on $x\text{CO}_2$ analysis, not $f\text{CO}_2$ calculated from other carbon parameters, such as pH, alkalinity or dissolved inorganic carbon;
2. Continuous CO_2 measurements have been made, not discrete CO_2 measurements;
3. The detection is based on an equilibrator system and is measured by infrared analysis or gas chromatography;
4. The calibration has included at least 2 non-zero gas standards, traceable to World Meteorological Organisation (WMO) standards;
5. The equilibrator temperature has been measured to within 0.05 °C accuracy;
6. The intake seawater temperature has been measured to within 0.05 °C accuracy;
7. The equilibrator pressure has been measured to within 0.5 hPa accuracy.

Note that criterion 1 also needs to be fulfilled for flags C and D. A satisfactory comparison with other data was required for flag A. This was carried out using either comparison with all other data previously obtained in the region or, if available, a set of formally defined cross-overs. The formally defined cross-overs were identified using a criterion that combined separation in space and separation in time into a single value. The algorithm that was used treated 1 day of separation in time as equivalent (heuristically) to 30 km of separation in space, i.e., if dx is the distance between points from two cruises in km, and dt is the separation between the same two points in days, then the separation between these two points would be given as $\sqrt{[dx^2 + (dt/30)^2]}$. The cross-over distance separating two cruises, dc , is the smallest value found comparing all pairs of points between two cruises. If a cross-over distance between two cruises was zero, a cruise had in most cases been erroneously duplicated, and the oldest version of the cruise data was excluded (flag X) in consultation with the data originator. Where the cross-over distance was relatively small, meaningful QC insights were often found by comparing observations from the two cruises. The LAS (Sect. 4.2.2) offered QC operators a means to compare cruise pairs with a small cross-over distance between them. No strict criteria were defined for judging the quality and significance of cross-overs.

The comparison with all data collected in a region was implemented in the Matlab QC toolbox for SOCAT (Sect. 4.2.3). This toolbox prompts the user to define the region of interest on a map, hence allowing the QC operator to use his/her expert knowledge of the regional characteristics during this process. When the region has been defined, the toolbox produces figures that compare the data subject to QC with all other data in the region in the time- and space domains, as well as in SST-space.

4.1.2 Metadata

Complete metadata documentation was required for flags A, B and C. By complete we mean that all the following information must be supplied:

1. The investigator;
2. The vessel;
3. The temporal coverage;
4. The analytical method;
5. The type of reported CO₂ data ($x\text{CO}_2$, $p\text{CO}_2$, $f\text{CO}_2$);
6. The number of CO₂ standards used with their approximate CO₂ mixing ratio and traceability;

7. A list of sensors and their accuracy, notably for:

- a. The equilibrator and seawater intake temperature;
- b. The equilibrator pressure.

Salinity does not need to be highly accurate for meeting the 2 μatm criterion, as the sensitivity of the $x\text{CO}_2$ to $f\text{CO}_2$ calculation is small (for example $x\text{CO}_2$ of 360 ppm at 20 °C and 1 atm yields $f\text{CO}_2$ of 347.22 μatm and 347.24 μatm at salinity 30 and 35, respectively). The metadata information had to either appear in the metadata themselves or in a publication cited in the metadata.

4.1.3 Extended quality control deemed acceptable

Flags A, B, C and D all required an extended QC with acceptable results. This extended QC included checks of the sampling positions and time, atmospheric pressures, salinity, intake and equilibrator temperatures, as well as recommended $f\text{CO}_2$ data, and included also a comparison with other data from the same region, if possible. The parameters were checked for range and occurrence of sudden, unrealistic jumps, and data from multiple streams were compared whenever possible (equilibrator, atmospheric and NCEP pressure; measured and WOA salinity; intake and equilibrator temperatures). Criteria for comparison of the intake and equilibrator temperatures were defined by the Southern and Indian Ocean SOCAT groups at their joint workshop in 2010 (IOCCP, 2010b):

- Warming should be less than 3 °C;
- Warming rate should be less than 1 °C h⁻¹, unless a rapid temperature front is apparent;
- Warming outliers should be less than 0.3 °C, compared to background data.

Apart from these, no strict criteria for QC were defined for the extended QC across all SOCAT groups. This will be improved in future versions of SOCAT.

If the data from a campaign were by-and-large of unacceptable quality, a cruise flag S was assigned. Whenever a large number (> 50, as a guideline) of non-acceptable data were found, the data file was suspended (Flag S), while the data contributor was invited to submit a suitably revised version of the data. If revised data were made available before the SOCAT quality control had been completed and were deemed of good quality, the data were included in version 1.5. Other resubmitted data will be included in the quality control for future SOCAT versions.

If it was not possible to establish contact with the data originator, or if the number of unacceptable data was sufficiently small (typically less than 50), WOCE flags 3 (questionable) or 4 (bad) were assigned to each unacceptable $f\text{CO}_2$ recommended value (Table 5). While WOCE flags 3

and 4 were assessed during version 1 quality control, virtually all such flags were unintentionally reset to flag 2 (good) in the version 1 data products. The WOCE flags 3 and 4 assigned during version 1 quality control will be applied in the SOCAT version 2 products.

4.2 Secondary quality control in practice

4.2.1 Secondary quality control by the regional groups

The regional groups had the responsibility for secondary QC of all cruises crossing their region. Regional SOCAT QC operators carried out secondary quality control and assigned flags to each cruise during the QC process upon evaluation of the data and metadata. The recommended $f\text{CO}_2$ and supporting data were made available via the Live Access Server during quality control. Data were evaluated according to the procedures outlined above. The QC was carried out in a variety of ways, either online via the LAS (Sect. 4.2.2) or offline (Sect. 4.2.3).

4.2.2 Live Access Server for quality control

The Live Access Server is a web server designed by NOAA PMEL (National Oceanic and Atmospheric Administration, Pacific Marine Environmental Laboratory) to provide access to geo-referenced scientific data (<http://ferret.pmel.noaa.gov/LAS>). Cruise data and metadata were ingested into a relational database and made available to the regional teams for evaluation through a version of the LAS, which had been enhanced with SOCAT quality control tools. Contents of the database included recommended $f\text{CO}_2$ values, ancillary parameters, cruise metadata, and reference variables drawn from other sources (Sect. 5.4). The LAS enabled QC operators to query the data collection using criteria of region, time period, seasonality, cruise and ship identifiers, and ranges of data values. The scientists could select data from one or more cruises, evaluate the data within the LAS and/or download subsets as compressed files for offline QC. The LAS offered QC evaluation tools, such as interactive property-property plots and co-inspection of cruises identified by the cross-over analysis (Sect. 4.2.1). The LAS provided access to the cruise metadata, which was evaluated as part of the QC. It also allowed uploading of ancillary documentation about the cruises and QC findings. The QC operators entered cruise flags and WOCE flags with comments explaining the rationale for their evaluations on the LAS during quality control. These cruise flags and comments are available via the Cruise Data Viewer (Sect. 5.5). The system alerted QC operators, when conflicting QC evaluations had been entered, allowing SOCAT scientists to evaluate and resolve these conflicts.

4.2.3 Offline quality control

A set of Matlab routines for data evaluation was available for offline QC (Olsen and Pierrot, 2010). These routines create a

series of plots, enabling QC. This toolbox prompts the user to define the region of interest on a map, hence allowing the QC operator to use his/her expert knowledge of the regional characteristics during the QC process. When the region has been defined, the toolbox produces figures that compare the data subject to QC with all other data in the region in the time- and space domains as well as in sea surface temperature space. Examples include a series of property-property plots, a series of plots of property versus time, and a series of plots comparing the $f\text{CO}_2$ data for the cruise subject to QC with all other data obtained in the region (as defined by the QC operator).

4.2.4 Conflicting cruise flags

Most cruises cross multiple regions, e.g., the coastal region and the North Atlantic Ocean. In SOCAT QC, a cruise needed to receive a cruise flag for each region that it crosses. A final check in the quality control consisted of checking conflicting cruise flags (Bakker). Most “conflicting” cruise flags reflected the absence of quality control in one region. These conflicts were resolved by carrying out appropriate QC and entering the missing cruise flags. Few truly conflicting cruise flags were encountered and in all cases a satisfactory solution was found.

4.3 Secondary quality control in practice

4.3.1 Secondary quality control by the regional groups

The regional groups had the responsibility for secondary QC of all cruises crossing their region. Regional SOCAT QC operators carried out secondary quality control and assigned flags to each cruise during the QC process upon evaluation of the data and metadata. The recommended $f\text{CO}_2$ and supporting data were made available via the Live Access Server during quality control. Data were evaluated according to the procedures in the SOCAT cookbook (Olsen and Metz, 2009). The QC was carried out in a variety of ways, either online via the LAS (Sect. 4.3.2) or offline (Sect. 4.3.3).

4.3.2 Live Access Server for quality control

The Live Access Server is a web server designed by NOAA PMEL (National Oceanic and Atmospheric Administration, Pacific Marine Environmental Laboratory) to provide access to geo-referenced scientific data (<http://ferret.pmel.noaa.gov/LAS>). Cruise data and metadata were ingested into a relational database and made available to the regional teams for evaluation through a version of the LAS, which had been enhanced with SOCAT quality control tools. Contents of the database included recommended $f\text{CO}_2$ values, ancillary parameters, cruise metadata, and reference variables drawn from other sources (Sect. 5.4). The LAS enabled QC operators to query the data collection using criteria of region, time period, seasonality, cruise and ship identifiers, and ranges of

data values. The scientists could select data from one or more cruises, evaluate the data within the LAS and/or download subsets as compressed files for offline QC. The LAS offered QC evaluation tools, such as interactive property-property plots and co-inspection of cruises identified by the cross-over analysis (Sect. 4.2.1). The LAS provided access to the cruise metadata, which had to be evaluated as part of the QC. It also allowed uploading of ancillary documentation about the cruises and QC findings. The QC operators entered cruise flags and WOCE flags with comments explaining the rationale for their evaluations on the LAS during quality control. The flags and comments are available via the Cruise Data Viewer (Sect. 5.5). The system alerted QC operators, when conflicting QC evaluations had been entered, allowing SOCAT scientists to evaluate and resolve these conflicts.

4.3.3 Offline quality control

A set of Matlab routines for data evaluation was available for offline QC (Olsen and Pierrot, 2010). These routines create a series of property-property plots, enabling QC operators to compare data from cruises in the same region. The $f\text{CO}_2$ is plotted and colour coded according to the input parameter used ($x\text{CO}_2$, $p\text{CO}_2$, $f\text{CO}_2$) in the (re-)calculation of recommended $f\text{CO}_2$ (Sect. 3.3). Examples include a figure comparing the $f\text{CO}_2$ versus sea surface temperature of a particular cruise to that for other cruises in the region. A second plot compares the monthly average and spread of the data in a box plot.

4.3.4 Suspended cruises and conflicting cruise flags

During the primary and secondary quality control, cruises were suspended from SOCAT (cruise flag “S” in Table 6), as minor and major flaws in the CO₂ data or in the data necessary for the (re-)calculation of $f\text{CO}_2$ became apparent. Data contributors were informed of these suspensions and were invited to resubmit their data upon making relevant corrections to the original data. In many cases data were resubmitted to SOCAT.

Most cruises cross multiple regions, e.g., the coastal region and the North Atlantic Ocean. In SOCAT QC, a cruise needed to receive a cruise flag for each region that it crosses. A final check in the quality control consisted of checking conflicting cruise flags (Bakker). Most “conflicting” cruise flags reflected the absence of quality control in one region. These conflicts were resolved by carrying out appropriate QC and entering the missing cruise flags. Few truly conflicting cruise flags were encountered and in all cases a satisfactory solution was found.

5 SOCAT products and tools

5.1 SOCAT cruises, versions, and time stamps

SOCAT data are publicly available via the SOCAT website (www.socat.info) as individual cruise data files (SOCAT version 1.4) (Sect. 5.2) and as regional and global, concatenated files (SOCAT version 1.5) (Sect. 5.3). SOCAT versions 1.4 and 1.5 include all cruises with a cruise flag A, B, C or D. A table of these cruises is available at [doi:10.1594/PANGAEA.769638](https://doi.org/10.1594/PANGAEA.769638) and provides information about the investigator, research vessel, Expocode, original cruise naming, metadata (as reported by the investigator), and temporal and geographical coverage. Through PANGAEA SOCAT is fed into the ICSU World Data System (WDS). The Global Earth Observation System of Systems (GEOSS), which is being built by the Group on Earth Observations (GEO), makes SOCAT available to other research communities.

The individual cruise data files (version 1.4) record observation time stamps at a resolution of integer minutes, rounding off the seconds, when they were available. Some cruises have multiple recommended $f\text{CO}_2$ values for a given time stamp (around 5 % of the observations). Individual cruise data files (Sect. 5.2) contain all recommended $f\text{CO}_2$ data, including multiple values per minute. However, handling multiple entries for the same time stamp can be problematic for some software programs. The SOCAT global group decided to average multiple entries within a given minute for the regional and global synthesis files (Sect. 5.3) as a pragmatic solution to this issue.

Table 5 lists the contents of the SOCAT files in versions 1.4 and 1.5. Matlab code by Pierrot and Landschützer for reading these files is available via the SOCAT website or directly at CDIAC (<http://cdiac.ornl.gov/ftp/oceans/SOCATv1.5/>).

5.2 Individual cruise data files (version 1.4)

Individual cruise data files (version 1.4) with cruise flags A, B, C and D are available via PANGAEA ([doi:10.1594/PANGAEA.767698](https://doi.org/10.1594/PANGAEA.767698)). These cruise data files include all recommended $f\text{CO}_2$ data with WOCE flags 2 (good), 3 (questionable) and 4 (bad), without listing these WOCE flags. Cruise data files archived at PANGAEA have not been averaged to remove multiple entries per minute (Sect. 5.1).

The individual cruise data files provide access to the metadata, the original CO₂ parameter(s) (as reported by the investigator), which were used to (re-)calculate $f\text{CO}_2$ (Sect. 3.3), and the (re-)calculated and quality controlled $f\text{CO}_2$ data. The files contain these additional parameters: WOA salinity (Antonov et al., 2006), NCEP/NCAR sea level pressure (Kalnay et al., 1996) and ETOPO2 (2006) bathymetry. Each individual cruise data file has been assigned a digital object

identifier (doi) for citation and transparency. Table 5 lists the parameters in the cruise data files.

5.3 SOCAT global and regional files (version 1.5)

Regional and global concatenated files (version 1.5) have been merged from the individual cruise data files for a subset of SOCAT parameters (Table 5). These concatenated files only contain recommended $f\text{CO}_2$ data with a WOCE flag 2 from cruises with a flag A, B, C or D. Table 5 lists the parameters in these regional and global synthesis files. Some changes have been applied relative to SOCAT version 1.4 (Sects. 5.1 and 5.2). Notably, multiple entries with the same time stamp were averaged for the global and regional synthesis files (Sect. 5.1).

Additional parameters have been added to the regional and global, concatenated files. These include Julian day (day of year), interpolated atmospheric $x\text{CO}_2$ extracted from GLOBALVIEW-CO₂ (2008), WOA salinity, NCEP/NCAR sea level pressure and ETOPO2 bathymetry. The global and regional files specify which reported CO₂ variable was used for (re-)calculation of recommended $f\text{CO}_2$ (Sect. 3.3; Table 5). Every line of the concatenated files contains a doi-string, which provides a link to the individual cruise data file with the original CO₂ parameter(s) and metadata at PANGAEA (Sect. 5.2).

The regional and global concatenated files (version 1.5) are publicly available as “compressed zip” text files via CDIAC (<http://cdiac.ornl.gov/ftp/oceans/SOCATv1.5/>), in Ocean Data View (ODV) format (<http://odv.awi.de/en/data/ocean/socat.v15.fco2.data/>) and via the interactive Cruise Data Viewer (Sect. 5.4). NetCDF files (Eaton et al., 2011) will be made available in the future. The text files exist as one very large global file, and as subset files per region, with no overlap between the regions. The latter means that data of a given cruise may have been divided into several regional files (for example North Atlantic, Tropical Atlantic and Coastal region).

5.4 Cruise Data Viewer (version 1.5)

The LAS Cruise Data Viewer provides interactive access to SOCAT version 1.5 on a Live Access Server. It provides all of the output capabilities described in Sect. 4 as tools for the SOCAT QC-ers, except for the ability to enter QC flags and comments. The Cruise Data Viewer also supplies variables from other sources that provide scientific context useful to users of the $f\text{CO}_2$ data: atmospheric $x\text{CO}_2$ values interpolated from GLOBALVIEW-CO₂ (2008), WOA 2005 salinity, NCEP/NCAR sea level pressure values (Kalnay et al., 1996), and bathymetry from ETOPO2 (2006).

The Cruise Data Viewer allows the inclusion of WOCE flag 3 (questionable) or 4 (bad) data when viewing or downloading data. When subsets are downloaded from the Cruise Data Viewer, each data line contains a doi-string that links di-

rectly to the relevant cruise data file with its original reported CO₂ parameters at PANGAEA. A “Table of Cruises” is available from the Cruise Data Viewer and lists the cruise flags, QC comments and SOCAT QC-er for each cruise. The Cruise Data Viewer can be accessed via the SOCAT website or directly at http://ferret.pmel.noaa.gov/SOCAT_cruise_viewer/.

5.5 Gridded products (version 1.5)

The gridded products provide values at a 1° latitude by 1° longitude resolution using monthly, annual, decadal and monthly climatological timescales, and at a 0.25° latitude by 0.25° longitude with monthly time resolution for coastal analysis (Sabine et al., 2013). The recommended $f\text{CO}_2$ with a WOCE flag 2 were gridded by two algorithms: (1) averages giving equal weight to each observation in a cell, and (2) averages giving equal weight to each cruise that passed through a cell. Mean, extremes and standard deviations of $f\text{CO}_2$ are provided. Other statistical measures include the number of cruises per cell, the number of observations per cell and measures of the degree to which the $f\text{CO}_2$ averaged values may be biased from the cell centre. The SOCAT version 1.5 gridded products have not been corrected for any temporal increase in surface water $f\text{CO}_2$. Gridded fields are available as NetCDF files from CDIAC (http://cdiac.ornl.gov/ftp/oceans/SOCATv1.5/SOCATv1.5_Gridded_Dat/) and via the interactive Gridded Data Viewer. For more details, refer to the accompanying paper by Sabine et al. (2013).

5.6 Gridded Data Viewer (version 1.5)

The interactive LAS Gridded Data Viewer enables users to explore the gridded SOCAT fields. The viewer displays maps and time series for the specific region or period selected. Sequences of fields can be viewed as animations. Simple statistics such as means, extremes, variance and counts, may be requested of the data. By requesting counts of the number of observations and cruises, a user is able to explore the global coverage of the SOCAT collection. Figure 4 obtained by this means, illustrates the north-south distribution of cruises in the years 2000 through 2007. The gridded viewer also supplies 1° latitude by 1° longitude marine surface variables from ICOADS (2008) that provide useful scientific context when exploring $f\text{CO}_2$: surface air temperature, sea level pressure, sea surface temperature, and surface wind speed. The Gridded Data Viewer can be accessed at (http://ferret.pmel.noaa.gov/SOCAT_gridded_viewer/) or via the link on the SOCAT website.

6 Lessons learned and outlook

6.1 Lessons learned

SOCAT has taken four years to be put together and has been a large, international, collaborative effort of the marine carbon




Figure 4. The number of SOCAT data points per 1° latitude by 1° longitude grid box for the years 2000 to 2007 shown as a function of latitude. This figure highlights the paucity of data in the Southern Hemisphere.

research community. SOCAT version 1.5 is the culmination of much hard work in data collection, data assembly and quality control by many seagoing marine carbon scientists around the world.

Lessons learned and improvements for future SOCAT releases have been discussed at the Surface Ocean CO₂ Data-to-Flux workshop (IOCCP, 2013). The lessons include a strong need for automating SOCAT with respect to data submission, metadata submission and quality control. The automation and other improvements will reduce the amount of work required for creating SOCAT data products and SOCAT quality control, while at the same time speeding up the whole process with the aim to provide regular updates.

The SOCAT global group, upon consultation with regional group leaders, has decided to start work on SOCAT version 2, while in parallel automating SOCAT for version 3. Data submission to SOCAT version 2 was closed on 31 December 2011. SOCAT version 2 products will report time in seconds as reported in the original data files to remove the need to calculate averaged data. Regular SOCAT releases are envisaged, e.g., every two years from SOCAT version 3 onwards. Such regular future SOCAT releases will require sustained funding for key players.

Colleagues are strongly encouraged to make public their surface water $f\text{CO}_2$ data and accompanying documentation from the global oceans and coastal seas, preferably via CDIAC (<http://cdiac.ornl.gov/oceans/submit.html>) for inclusion in future SOCAT releases. Data and metadata should be reported in the IOCCP (2004) recommended formats, which are also listed on the CDIAC website.

6.2 Automation of SOCAT procedures

Automation of the submission of data and metadata will include prompt feedback to the data originator on unrealistic data and property-property plots of the data, such that the data originator can carry out primary and initial secondary quality control. Such automation will facilitate harmonisation of the data for SOCAT and will strongly reduce the number of cruises suspended from SOCAT during secondary quality control.

In the future, new cruises will be added to the LAS at regular (e.g., two monthly) intervals, enabling QC operators to carry out regular SOCAT QC. The Live Access Server will be modified to automatically generate typical property-property plots for secondary QC. The LAS will be enhanced with features to enter cruise flags and QC comments for multiple cruises (e.g., on the same vessel).

6.3 SOCAT products for assessing the ocean carbon sink

The release of SOCAT version 1.5 represents a milestone in ocean carbon research. Research using SOCAT will highlight the response of surface water $f\text{CO}_2$ and the oceanic CO₂ sink to increasing levels of atmospheric CO₂ in a changing climate. The SOCAT products can be used in studies of spatial and temporal (seasonal, interannual and decadal) variability and trends in surface water $f\text{CO}_2$. The SOCAT products will enable validation of model distributions of surface water $f\text{CO}_2$ and air–sea CO₂ fluxes. SOCAT will aid process studies of oceanic $f\text{CO}_2$ variability, e.g., in the North Atlantic, in the Pacific Ocean, in coastal seas, in the Arctic Ocean, in seasonally ice-covered Southern Ocean regions, near remote islands and oceanographic fronts. The SOCAT products may be used to create monthly basin-wide $f\text{CO}_2$ maps for the most data-rich basins by a range of techniques such as neural networks, statistical techniques and algorithms (e.g., Lefèvre et al., 2005; Telszewski et al., 2009). These $f\text{CO}_2$ maps can be used for calculating basin-wide monthly CO₂ air–sea fluxes, which may constrain atmospheric inversions for global atmospheric carbon budgets. Study of length scales of $f\text{CO}_2$ variability will provide information on the minimum sampling coverage required for quantifying the oceanic CO₂ sink with sufficient accuracy (e.g., Lenton et al., 2009). It is expected that the regular SOCAT releases will become a crucial tool in quantification of changes in oceanic CO₂ uptake and in global climate research. Increasing the number of surface ocean CO₂ data has in the past significantly modified the estimate of the oceanic CO₂ sink (e.g., Takahashi et al., 2009b). SOCAT and its future development will contribute to further enhance the reliability of such assessments.

Acknowledgements. SOCAT is promoted by IOCCP, the Surface Ocean Lower Atmosphere Study, and the Integrated Marine Biogeochemistry and Ecosystem Research program. Douglas Wallace (Dalhousie University, Canada, and former SOLAS chair), Emilie Brévière (SOLAS executive officer) and Lisa Maddison (IMBER deputy executive officer) have strongly encouraged SOCAT. Support for SOCAT has been received from the Bjerknes Centre for Climate Research, the University of Bergen, Uni Research (Norway), the US National Oceanic and Atmospheric Administration, the University of Washington, Oak Ridge National Laboratory (US), the University of East Anglia (UEA, UK), PANGAEA – Data Publisher for Earth & Environmental data (Germany), the Alfred Wegener Institute for Polar and Marine Research (Germany), the Centre National de la Recherche Scientifique (France), the Research Council of Norway (CARBON-HEAT), the US National Science Foundation (OCE-1068958), the international Scientific Committee on Oceanic Research (SCOR, US, OCE-0938349), the European Union integrated projects CarboOcean (GOCE 511176-1) and CarboChange (FP7 264879), the UK Ocean Acidification Research Programme (NE/H017046/1; funded by the Natural Environment Research Council, the Department for Energy and Climate Change and the Department for Environment, Food and Rural Affairs) and the UK National Centre for Earth Observation. Support for SOCAT meetings has been received from IOCCP, IMBER, the National Institute for Environmental Studies (NIES, Japan), the Commonwealth Scientific and Industrial Research Organisation (CSIRO, Australia), GEOMAR (Germany) and the European Cooperation in Science and Technology (COST) Action 735 (UK). This is publication no. A417 for the Bjerknes Centre for Climate Research. This study is a contribution to the Center for Climate Dynamics (SKD) within the Bjerknes Centre for Climate Research.

Edited by: F. Schmitt

References

- Antonov, J. I., Locarnini, R. A., Boyer, T. P., Mishonov, A. V., and Garcia, H. E.: World Ocean Atlas 2005, in: Volume 2: Salinity, edited by: Levitus, S., NOAA Atlas NESDIS 62, US Government Printing Office, Washington, DC, 182 pp., 2006.
- Archer, D., Kheshgi, H., and Maier-Reimer, E.: Multiple timescales for neutralization of fossil fuel CO₂, *Geophys. Res. Lett.*, 24, 405–408, 1997.
- Atlantic Oceanographic and Meteorological Laboratory (AOML): Underway pCO₂ Systems workshop, AOML National Oceanographic and Atmospheric Administration, Miami, 2–3 October 2002, <http://www.aoml.noaa.gov/ocd/gcc/uwpc2/workshops/> (last access: 13 March 2012), 2002.
- Bakker, D. C. E., Etcheto, J., Boutin, J., Dandonneau, Y., and Merlivat, L.: Variability of surface water fCO₂ during seasonal upwelling in the equatorial Atlantic Ocean as observed by a drifting buoy. *J. Geophys. Res.*, 106, 9241–9253, doi:10.1029/1999JC000275, 2001.
- Bakker, D. C. E., Hoppema, M., Schröder, M., Geibert, W., and de Baar, H. J. W.: A rapid transition from ice covered CO₂-rich waters to a biologically mediated CO₂ sink in the eastern Weddell Gyre, *Biogeosciences*, 5, 1373–1386, doi:10.5194/bg-5-1373-2008, 2008.
- Bakker, D. C. E., Pfeil, B., Olsen, A., Metzl, N., Sabine, C. L., Hankin, S., Koyuk, H., Kozyr, A., Malczyk, J., Manke, A., and Telszewski, M.: Global data products help assess changes to the ocean carbon sink, *Eos*, 93, 125–126, doi:10.1029/2012EO120001, 2012.
- Cooper, D. J., Watson, A. J., and Ling, R. D.: Variations of pCO₂ along a North Atlantic shipping route (UK to the Caribbean): A year of automated observations, *Mar. Chem.*, 60, 147–164, 1998.
- Copin-Montégut, C.: A new formula for the effect of temperature on the partial pressure of CO₂ in seawater, *Mar. Chem.*, 25, 29–37, 1988.
- Copin-Montégut, C.: Corrigendum. A new formula for the effect of temperature on the partial pressure of CO₂ in seawater, *Mar. Chem.*, 27, 143–144, 1989.
- DeGrandpré, M. D., Baehr, M. W., and Hammar, T. R.: Development of an optical chemical sensor for oceanographic applications: the submersible autonomous moored instrument for seawater CO₂, in: *Chemical sensors in oceanography*, edited by: Varney, M. S., Gordon and Breach Science Publishers, Amsterdam, the Netherlands, 123–142, 2000.
- Dickson, A. G., Sabine, C. L., and Christian, J. R. (Eds.): Guide to best practices for ocean CO₂ measurements, PICES Special Publication 3, IOCCP Report 8, 191 pp., 2007.
- Eaton, B., Gregory, J., Drach, B., Taylor, K., Hankin S., Caron, J., Signell, R., Bentley, P., Rappa, G., Höck, H., Pamment, A., and Juckes, M.: NetCDF Climate and forecast (CF) metadata conventions version 1.6, <http://cf-pcmdi.llnl.gov/documents/cf-conventions/1.6/cf-conventions.html#discrete-sampling-geometries>, 2011.
- ETOPO2: 2-minute Gridded Global Relief Data (ETOPO2v2), US Dept. of Commerce, National Oceanic and Atmospheric Administration, National Geophysical Data Center, <http://www.ngdc.noaa.gov/mgg/fliers/06mgg01.html>, 2006.
- Friederich, G. E., Ledesma, J., Ulloa, O., and Chavez, F. P.: Air-sea carbon dioxide fluxes in the coastal southeastern tropical Pacific, *Prog. Oceanogr.*, 79, 156–166, 2008.
- Friedlingstein, P., Cox, P., Betts, R., Bopp, L., Von Bloh, W., Brovkin, V., Cadule, P., Doney, S., Eby, M., Fung, I., Govindasamy, B., John, J., Jones, C., Joos, F., Kato, T., Kawamiya, M., Knorr, W., Lindsay, K., Matthews, H. D., Raddatz, T., Rayner, P., Reick, C., Roeckner, E., Schnitzler, K.-G., Schnur, R., Strassmann, K., Weaver, A. J., Yoshikawa, C., and Zeng, N.: Climate-carbon cycle feedback analysis, results from the C4MIP model intercomparison, *J. Climate*, 19, 3337–3353, 2006.
- GLOBALVIEW-CO2: Cooperative Atmospheric Data Integration Project – Carbon Dioxide, 2008 version, NOAA ESRL, Boulder, Colorado, <http://www.esrl.noaa.gov/gmd/ccgg/globalview/co2/>, 2008.
- Gordon, L. L. and Jones, L. B.: The effect of temperature on carbon dioxide partial pressure in seawater, *Mar. Chem.*, 1, 317–322, 1973.
- Goyet, C., Millero, F. J., Poisson, A., and Shafer, D. K.: Temperature dependence of CO₂ fugacity in seawater, *Mar. Chem.*, 44, 205–219, 1993.
- Gruber, N., Körtzinger, A., Borges, A., Claustre, H., Doney, S. C., Feely, R. A., Hood, M., Ishii, M., Kozyr, A., Monteiro, P., Nojiri, Y., Sabine, C. L., Schuster, U., Wallace, D. W. R., and Wanninkhof, R.: Plenary paper: Towards an integrated observing system for ocean carbon and biogeochemistry at a time of change,

- in: Proceedings of OceanObs'09: Sustained ocean observations and information for society, Vol. 1, Venice, Italy, 21–25 September 2009, edited by: Hall, J., Harrison, D. E., and Stammer, D., ESA Publication WPP-306, 18 pp., doi:10.5270/OceanObs09, 2010.
- Hood, E. M., Merlivat, L., and Johannessen, T.: *f*CO₂ variations and air-sea flux of CO₂ in the Greenland Sea gyre using high frequency time series from CARIOCA drift-buoys, *J. Geophys. Res.*, 104, 20571–20583, doi:10.1029/1999JC900130, 1999.
- ICoads: International Comprehensive Ocean-Atmosphere Data Set, 1 Degree, Release 2.5. NOAA, ESRL PSD, Boulder, Colorado, <http://www.esrl.noaa.gov/psd/data/gridded/data.coads.1deg.html>, 2008.
- IMBER: Joint SOLAS-IMBER Ocean Carbon Research Implementation plan, IMBER Report 1, SOLAS Report, 41 pp., 2005.
- IOCCP: Ocean surface *p*CO₂, data integration and database development workshop, National Institute for Environmental Studies, Tsukuba, Japan, 14–17 January 2004, IOCCP Report 2, www.ioccp.org, 2004.
- IOCCP: Surface Ocean CO₂ Variability and Vulnerabilities Workshop. UNESCO, Paris, France, 11–14 April 2007. IOCCP Report 7, www.ioccp.org, 2007.
- IOCCP: Surface Ocean CO₂ Atlas Project, UNESCO, Paris, France, IOCCP Report 9, www.ioccp.org, 2008.
- IOCCP: Joint workshop SOCAT Coastal Regional Group & COST Action 735 Working Group 3. IFM-GEOMAR, Kiel, Germany, 22–23 January 2009, www.ioccp.org, 2009a.
- IOCCP: SOCAT Atlantic and Southern Oceans Regional Meeting, University of East Anglia, Norwich, UK, 25–26 June 2009, IOCCP Report 13, www.ioccp.org, 2009b.
- IOCCP: SOCAT Pacific Regional Meeting, National Institute for Environmental Studies, Tsukuba, Japan, 18–20 March 2009, IOC Workshop Report 221, IOCCP Report 12, www.ioccp.org, 2010a.
- IOCCP: Southern and Indian Ocean SOCAT Workshop, CSIRO Marine Laboratories, Hobart, Tasmania, Australia, 16–18 June 2010, IOC Workshop Report 234, IOCCP Report 21, www.ioccp.org, 2010b.
- IOCCP: Surface Ocean CO₂ Data-to-Flux workshop. UNESCO, Paris, France, 12–13 September 2011, www.ioccp.org, IOCCP Report, in preparation, 2013.
- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds, R., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *B. Am. Meteorol. Soc.*, 77, 437–470, 1996.
- Körtzinger, A., Thomas, H., Schneider, B., Gronau, N., Mintrop, L., and Duinker, J. C.: At-sea intercomparison of two newly designed underway *p*CO₂ systems – Encouraging results, *Mar. Chem.*, 52, 133–145, 1996.
- Körtzinger, A., Mintrop, L., Wallace, D. W. R., Johnson, K. M., Neill, C., Tilbrook, B., Towler, P., Inoue, H., Ishii, M., Shaffer, G., Torres, R. F., Ohtaki, E., Yamashita, E., Poisson, A., Brunet, C., Schauer, B., Goyet, C., and Eiseheid, G.: The international at-sea intercomparison of *f*CO₂ systems during the R/V Meteor cruise 36/1 in the North Atlantic Ocean, *Mar. Chem.*, 72, 171–192, 2000.
- Körtzinger, A., Send, U., Lampitt, R. S., Hartman, S., Wallace, D. W. R., Karstensen, J., Villagarica, M. G., Linás, O., and DeGrandpré, M. D.: The seasonal *p*CO₂ cycle at 49° W 16.5° N in the northeastern Atlantic Ocean and what it tells us about biological productivity, *J. Geophys. Res.*, 113, C04020, doi:10.1029/2007JC004347, 2008.
- Lefèvre, N., Ciabrin, J. P., Michard, G., Briant, B., Duchaffaut, M., and Merlivat, L.: A new optical sensor for *p*CO₂ measurements in seawater, *Mar. Chem.*, 42, 189–198, 1993.
- Lefèvre, N., Andrié, C., Dandonneau, Y., Reverdin, G., and Rodier, M.: *p*CO₂, chemical properties, and estimated new production in the equatorial Pacific in January–March 1991, *J. Geophys. Res.*, 99, 12639–12654, 1994.
- Lefèvre, N., Watson, A. J., and Watson, A. R.: A comparison of multiple regression and neural network techniques for mapping in situ *p*CO₂ data, *Tellus B*, 57, 375–384, 2005.
- Lefèvre, N., Guillot, A., Beaumont, L., and Danguy, T.: Variability of *f*CO₂ in the Eastern Tropical Atlantic from a moored buoy, *J. Geophys. Res.*, 113, C01015, doi:10.1029/2007JC004146, 2008.
- Leinweber, A., Gruber, N., Frenzel, H., Friederich, G. E., and Chavez, F. P.: Diurnal carbon cycling in the surface ocean and lower atmosphere of Santa Monica Bay, California, *Geophys. Res. Lett.*, 36, L08601, doi:10.1029/2008GL037018, 2009.
- Lenton, A., Bopp, L., and Matear, R. J.: Strategies for high-latitude northern hemisphere CO₂ sampling now and in the future, *Deep-Sea Res. Pt. II*, 56, 523–532, 2009.
- Merlivat, L. and Brault, P.: CARIOCA buoy – carbon dioxide monitor, *Sea Technol.*, 36, 23–30, 1995.
- Merlivat, L., González Dávila, M., Caniaux, G., Boutin, J., and Reverdin, G.: Mesoscale and diel to monthly variability of CO₂ and carbon fluxes at the ocean surface in the northeastern Atlantic, *J. Geophys. Res. Oceans*, 114, C03010, doi:10.1029/2007JC004657, 2009.
- Olsen, A. and Metzl, N.: SOCAT QC cookbook for SOCAT participants; available at: www.socat.info/publications.html (last access: 11 March 2012), 2009.
- Olsen, A. and Pierrot, D.: Matlab routines to aid QC of SOCAT data; available at: www.socat.info/publications.html (last access: 2 March 2012), 2010.
- Olsen, A., Brown, K. R., Chierici, M., Johannessen, T., and Neill, C.: Sea-surface CO₂ fugacity in the subpolar North Atlantic, *Biogeosciences*, 5, 535–547, doi:10.5194/bg-5-535-2008, 2008.
- Parard, G., Lefèvre, N., and Boutin, J.: Sea water fugacity of CO₂ at the PIRATA mooring at 6° S, 10° W, *Tellus B*, 62, 636–648, 2010.
- Pfeil, B. and Olsen, A.: Uniform format surface *f*CO₂ database; available at: www.socat.info/publications.html (last access: 11 March 2012), 2009.
- Pierrot, D., Neill, C., Sullivan, K., Castle, R., Wanninkhof, R., Lüger, H., Johannessen, T., Olsen, A., Feely, R. A., and Cosca, C. E.: Recommendations for autonomous underway *p*CO₂ measuring systems and data reduction routines, *Deep-Sea Res. Pt. II*, 56, 512–522, doi:10.1016/j.dsr2.2008.12.005, 2009.
- Riebesell, U., Körtzinger, A., and Oschlies, A.: Sensitivities of marine carbon fluxes to ocean change, *P. Natl. Acad. Sci. USA*, 106, 20602–20609, 2009.
- Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R., Wong, C. S., Wallace, D. W. R., Tilbrook, B., Millero, F. J., Peng, T. H., Kozyr, A., Ono, T., and

- Rios A. F.: The oceanic sink for anthropogenic CO₂, *Science*, 305, 367–371, 2004.
- Sabine, C. L., Ducklow, H., and Hood, M.: International carbon coordination. Roger Revelle's legacy in the Intergovernmental Oceanographic Commission, *Oceanography*, 23, 48–61, 2010.
- Sabine, C. L., Hankin, S., Koyuk, H., Bakker, D. C. E., Pfeil, B., Olsen, A., Metzl, N., Kozyr, A., Fassbender, A., Manke, A., Malczyk, J., Akl, J., Alin, S. R., Bellerby, R. G. J., Borges, A., Boutin, J., Brown, P. J., Cai, W.-J., Chavez, F. P., Chen, A., Cosca, C., Feely, R. A., González-Dávila, M., Goyet, C., Hardman-Mountford, N., Heinze, C., Hoppema, M., Hunt, C. W., Hydes, D., Ishii, M., Johannessen, T., Key, R. M., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lourantou, A., Merlivat, L., Midorikawa, T., Mintrop, L., Miyazaki, C., Murata, A., Nakadate, A., Nakano, Y., Nakaoka, S., Nojiri, Y., Omar, A. M., Padin, X. A., Park, G.-H., Paterson, K., Perez, F. F., Pierrot, D., Poisson, A., Ríos, A. F., Salisbury, J., Santana-Casiano, J. M., Sarma, V. V. S. S., Schlitzer, R., Schneider, B., Schuster, U., Sieger, R., Skjelvan, I., Steinhoff, T., Suzuki, T., Takahashi, T., Tedesco, K., Telszewski, M., Thomas, H., Tilbrook, B., Vandemark, D., Veness, T., Watson, A. J., Weiss, R., Wong, C. S., and Yoshikawa-Inoue, H.: Surface Ocean CO₂ Atlas (SOCAT) gridded data products, *Earth Syst. Sci. Data*, 1, 145–153, doi:10.5194/essd-1-145-2013, 2013.
- Skjelvan, I., Johannessen, T., and Miller, L. A.: Interannual variability of *f*CO₂ in the Greenland and Norwegian Seas, *Tellus*, 51B, 477–489, 1999.
- Swift, J.: A guide to submitting CTD/hydrographic/tracer data and associated documentation to the CLIVAR and Carbon Hydrographic Data Office, version of 22 April 2008, UCSD Scripps Institution of Oceanography, 37 pp., 2008.
- Takahashi, T.: Carbon dioxide in the atmosphere and in Atlantic Ocean water, *J. Geophys. Res.*, 66, 477–494, 1961.
- Takahashi, T., Olafsson, J., Goddard, J. G., Chipman, D. W., and Sutherland, S. C.: Seasonal variation of CO₂ and nutrients in the high-latitude surface oceans: a comparative study, *Global Biogeochem. Cy.*, 7, 843–878, 1993.
- Takahashi, T., Feely, R. A., Weiss, R., Wanninkhof, R., Chipman, D. W., Sutherland, S. C., and Takahashi, T. T.: Global air-sea flux of CO₂: An estimate based on measurements of sea-air *p*CO₂ difference, *P. Natl. Acad. Sci. USA*, 94, 8292–8299, 1997.
- Takahashi, T., Sutherland, S. G., Sweeney, C., Poisson, A. P., Metzl, N., Tilbrook, B., Bates, N. R., Wanninkhof, R., Feely, R. A., Sabine, C. L., Olafsson, J., and Nojiri, Y.: Global sea-air CO₂ flux based on climatological surface ocean *p*CO₂, and seasonal biological and temperature effects, *Deep-Sea Res. Pt. II*, 49, 1601–1622, 2002.
- Takahashi, T., Sutherland, S. C., and Kozyr, A.: Global Ocean Surface Water Partial Pressure of CO₂ Database: Measurements Performed During 1968–2008 (Version 2008), ORNL/CDIAC-152, NDP-088r, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Dept. of Energy, Oak Ridge, Tenn., doi:10.3334/CDIAC/otg.ndp088r, 2009a.
- Takahashi, T., Sutherland, S. C., Wanninkhof, R., Sweeney, C., Feely, R. A., Chipman, D. W., Hales, B., Friederich, G., Chavez, F., Sabine, C. L., Watson, A. J., Bakker, D. C. E., Schuster, U., Metzl, N., Inoue, H. Y., Ishii, M., Midorikawa, T., Nojiri, Y., Körtzinger, A., Steinhoff, T., Hoppema, J. M. J., Olafsson, J., Arnarson, T. S., Tilbrook, B., Johannessen, T., Olsen, A., Bellerby, R. G. J., Wong, C. S., Delille, B., Bates, N. R., and De Baar, H. J. W.: Climatological mean and decadal change in surface ocean *p*CO₂, and net sea-air CO₂ flux over the global oceans, *Deep-Sea Res. Pt. II*, 56, 544–577, doi:10.1016/j.dsr2.2008.12.009, 2009b.
- Takahashi, T., Sutherland, S. C., and Kozyr, A.: Global Ocean Surface Water Partial Pressure of CO₂ Database: Measurements Performed During 1957–2010 (Version 2010), ORNL/CDIAC-159, NDP-088(V2010), Carbon Dioxide Information Analysis Center, Oak Ridge Nat. Lab., US Dept. of Energy, Oak Ridge, Tenn., doi:10.3334/CDIAC/otg.ndp088(V2010), 2011.
- Tanhua, T., van Heuven, S., Key, R. M., Velo, A., Olsen, A., and Schirnick, C.: Quality control procedures and methods of the CARINA database, *Earth Syst. Sci. Data*, 2, 35–49, doi:10.5194/essd-2-35-2010, 2010.
- Telszewski, M., Chazottes, A., Schuster, U., Watson, A. J., Moulin, C., Bakker, D. C. E., González-Dávila, M., Johannessen, T., Körtzinger, A., Lüger, H., Olsen, A., Omar, A., Padin, X. A., Ríos, A. F., Steinhoff, T., Santana-Casiano, M., Wallace, D. W. R., and Wanninkhof, R.: Estimating the monthly *p*CO₂ distribution in the North Atlantic using a self-organizing neural network, *Biogeosciences*, 6, 1405–1421, doi:10.5194/bg-6-1405-2009, 2009.
- Wada, A., Midorikawa, T., Ishii, M., and Motoi, T.: Carbon system changes in the East China Sea induced by typhoons Tina and Winnie in 1997, *J. Geophys. Res.*, 116, C07014, doi:10.1029/2010JC006701, 2011.
- Wanninkhof, R. and Thoning, K.: Measurement of fugacity of CO₂ in surface water using continuous and discrete sampling methods, *Mar. Chem.*, 44, 189–204, 1993.
- Watson, A. J., Schuster, U., Bakker, D. C. E., Bates, N. R., Corbiere, A., González-Dávila, M., Friedrich, T., Heinze, C., Johannessen, T., Körtzinger, A., Metzl, N., Olafsson, J., Olsen, A., Oschlies, A., Padin, X. A., Pfeil, B., Santana-Casiano, J. M., Steinhoff, T., Telszewski, M., Rios, A. F., Wallace, D. W. R., and Wanninkhof, R.: A network to accurately estimate the North Atlantic sink for CO₂, *Science*, 326, 1391–1393, 2009.
- Weiss, R. F.: Determinations of carbon dioxide and methane by dual catalyst flame ionization chromatography and nitrous oxide by electron capture chromatography, *J. Chromatogr. Sci.*, 19611–19616, 1981.
- Weiss, R. F., Jahnke, R. A., and Keeling, C. D.: Seasonal effects of temperature and salinity on the partial pressure of CO₂ in seawater, *Nature*, 300, 511–513, 1982.