

Intensification of the Amazon hydrological cycle over the last two decades

M. Gloor,¹ R. J. W. Brienen,¹ D. Galbraith,¹ T. R. Feldpausch,¹ J. Schöngart,² J.-L. Guyot,³ J. C. Espinoza,⁴ J. Lloyd,⁵ and O. L. Phillips¹


Received 18 December 2012; revised 14 March 2013; accepted 15 March 2013; published 14 May 2013.

[1] The Amazon basin hosts half the planet's remaining moist tropical forests, but they may be threatened in a warming world. Nevertheless, climate model predictions vary from rapid drying to modest wetting. Here we report that the catchment of the world's largest river is experiencing a substantial wetting trend since approximately 1990. This intensification of the hydrological cycle is concentrated overwhelmingly in the wet season driving progressively greater differences in Amazon peak and minimum flows. The onset of the trend coincides with the onset of an upward trend in tropical Atlantic sea surface temperatures (SST). This positive longer-term correlation contrasts with the short-term, negative response of basin-wide precipitation to positive anomalies in tropical North Atlantic SST, which are driven by temporary shifts in the intertropical convergence zone position. We propose that the Amazon precipitation changes since 1990 are instead related to increasing atmospheric water vapor import from the warming tropical Atlantic. **Citation:** Gloor, M., R. J. W. Brienen, D. Galbraith, T. R. Feldpausch, J. Schöngart, J.-L. Guyot, J. C. Espinoza, J. Lloyd, and O. L. Phillips (2013), Intensification of the Amazon hydrological cycle over the last two decades, *Geophys. Res. Lett.*, 40, 1729–1733, doi:10.1002/grl.50377.

1. Introduction

[2] The Amazon basin is an important component of the global climate system, being one of the main locations of deep convection and with a river discharge contributing ~17% of total global freshwater input to the oceans [Callède *et al.* 2010]. It hosts the largest contiguous area of tropical forest (~ 5 mio. km²), modulating its own hydrological regime through the recirculation of precipitation and the

need thus to be interpreted with some caution. The second record is Amazon basin catchment integrated precipitation based here primarily on the 0.5×0.5 longitude by latitude Climate Research Unit (CRU) climatology [



decades (Figure 1). The upward trend is clearly evident in the monthly records and is even more pronounced in daily records indicating also a shift toward more severe events. The annual mean discharge record indicates changes from the middle of the last century onward. There was a temporary fast intensification around 1970 which had dissipated by 1980, with a second intensification starting around 1990 and continuing until today ($\sim 8\%$ from 1990 to 2010, $p=0.26$ for annual means, $p=0.08$ for monthly maxima). This increase over the last two decades is also evident in the Amazon catchment integrated precipitation, which has increased at a similar relative rate ($\sim 10\%$ over the same period, $p=0.24$; for another more stringent measure of significance, see Figure 3) with results similar when using the GPCP precipitation climatology. The decadal-scale event in the 1970s is well known and is generally associated with the Pacific Decadal Oscillation and a shift to more intense ENSO variability around 1970 [Miller, 1994]. The general upward trend in river discharge over the last century had already been reported by Callède *et al.* [2004].

[7] Where in the basin has precipitation increased the most? We find no consistent trend across the basin but northwestern Amazonia shows the strongest increase in precipitation (up to 80 mm month^{-1} when comparing the 2000–2009 mean with the 1981–1990 mean; Figure 2). This is also the region, which is already the wettest part of the basin. Southwestern Amazonia and the Gran Chaco region toward the southern periphery of the basin show some drying (up to $-20 \text{ mm month}^{-1}$). Possible reasons for this trend are changes in the land surface cover due to deforestation, and thus, water recirculation to the atmosphere shifts in the intertropical convergence zone (ITCZ) due to changes in albedo caused by deforestation [Doughty *et al.* 2012] or changes in water vapor export out of the basin to the southeast [Espinoza *et al.* 2009]. The southern drying trend

may also be due to decadal scale variation as it disappears when choosing a different reference period (e.g., 1961–1980). We also note that the results of Chou *et al.* [2013] based on the GPCP (Global Precipitation Climatology Project) reveal a picture quite similar to Figure 2 of our study.

[8] To avoid the risk of pattern overinterpretation, we indicate in Figures 3a and 3b the locations of the station records on which the maps are based as well as time trend significance levels calculated from CRU monthly mean station records (available at <http://www.metoffice.gov.uk/hadobs/crutem4/data/download.html>). We also verified that the stronger trend in the Western basin is not an interpolation or “site switching” artifact by estimating the station time series mean time trend directly from the monthly mean observational records (Figures S2a and S2b).

[9] The increasing trend in annual precipitation over the last two decades occurs overwhelmingly during the wet season (the wettest 9 months; Figure 3b), again with the exceptions of the southwestern and Gran Chaco regions which show slight decreases. One consequence is a strong increasing trend in the difference in annual monthly maxima minus annual monthly minima in river discharge (Figure 1a and Figure S1) exhibiting a change of approximately 10% over the period from 1990 to 2009 ($p=0.18$ for period 1990–2010, $p=0.004$ for 1980–2010). Although not included in our analysis, the year 2012 is in line with the general wetting trend during the wet season with Amazon stage measurements in Manaus being the highest recorded in time since 1903.

4. Discussion of Possible Causes

[10] Finally, we attempt attribution by comparing the Amazon River discharge and basin integrated precipitation

records with tropical Atlantic sea surface temperature (Figure 1). There are two notable features. First, interannual variability in the hydrology record tends to be anticorrelated with interannual variability of sea surface temperature, i.e., negative precipitation anomalies tend to be associated with positive sea surface temperature anomalies. This anticorrelation has been attributed to variations in the location of the ITCZ, which is steered northward by the positive equatorial north Atlantic SST anomalies [e.g., Yoon and Zeng 2010; Marengo et al. 2008]. Examples for this anticorrelation are the most recent four severe drought events in 1995, 1998, 2005, 2010 (Figures 1a and 1c), which were particularly intense over western Amazonia [Espinoza Villar et al., 2011] and the decade long positive Amazon precipitation anomaly in the 1970s. Atlantic tropical sea surface temperature anomalies are, however, not the only influence on Amazon precipitation patterns as El Niño Southern Oscillation (ENSO) events exert the most dominant control on interannual variability. El Niño phases are associated with negative precipitation anomalies in the north of the basin leading to lower than usual Amazon River discharge. Likewise La Niña phases are associated with positive precipitation anomalies and higher river discharge [e.g., Costa and Foley, 1999; Marengo et al., 2011; Espinoza et al., 2012].

[11] The second notable feature is that the Tropical Atlantic SST record (17°S–20°N) exhibits a strong upward trend ($p=0.002$, Figure 1c). This is similar to the hydrology record and commences at a similar time, thus suggesting that the two phenomena might be linked. The positive correlation between the upward trends in hydrological records and Tropical Atlantic SSTs over the last two decades also suggests a different mechanism from the ITCZ shifts that is associated with a negative correlation between the two variables.

[12] One possible cause for the changes of the hydrological cycle over the last two decades is an intensification of the monsoonal circulation during austral summer and an associated increase in the sea-land temperature contrast. However, this explanation is not supported by sea surface and land temperature data. The coincidence of the onset of the trends suggests that the intensification of the hydrological cycle might simply be due to the increase of water vapor input by the airstream entering the basin from the tropical Atlantic and “feeding” the Amazon hydrological cycle [Stohl and James, 2005].

[13] An increase in sea surface temperature of $\sim 0.7^{\circ}\text{C}$ as observed for the 1990–2010 period would be expected to cause an increase in the saturation water vapor pressure of approximately 5% (Clausius-Clapeyron equation) [e.g., Held and Soden, 2006], an expectation confirmed by remote sensing estimates of tropical Atlantic air column vapor content [Santer et al. 2007]. All else being equal then, the increased water vapor input can explain around half of the observed Amazon River discharge increase. An additional possibly linked mechanism is therefore likely to be involved, like changes in water vapor outflow or changes in atmospheric stability and cloud convection. A more detailed analysis of the causes behind the observed trends of the hydrological records will likely only be possible with the help of Earth system model simulations.

[14] We note that a link between positive (southern) tropical Atlantic sea surface temperatures and increased precipitation particularly in the northwestern region of the Amazon during the wet season is consistent with the analyses of Yoon and Zeng [2010, Figures 8e and 8f] and Marengo et al. [2011], who analyzed the causes of the exceptional 2009 flood

(their Figure 1f). Furthermore, trends in sea surface salinity also reveal an intensifying trend of the hydrological cycle globally and particularly in the tropics [Durack et al., 2012]. We finally would like to stress that, while the link to a warming tropical Atlantic sea surface might suggest a link to global warming and thus may indicate a longer-term regime shift of the Amazon hydrological regime, the changes from 1990 to present may of course instead be due to multidecadal natural variability.

5. Summary

[15] There is thus good evidence that the catchment of the world’s largest river has become substantially wetter since the late twentieth century starting approximately in 1990 with a simultaneous strong trend toward an increase in the annual amplitude of river discharge and an increase of the severity of events. Together, these observations do not support the recent perception of a general imminent drying of the Amazon basin, although there are indications of some drying of the drier southern margins of the basin. The onset of the increasing trend coincides with the onset of a similar increasing trend in tropical Atlantic sea surface temperatures, which suggests that the two phenomena might be related, possibly through an increase in water vapor import from the tropical Atlantic Ocean. It is not immediately clear how an intensification of the hydrological cycle affects forest productivity and the Amazon carbon balance. It may however not be entirely coincidental that the onset of the intensification starts at approximately the same time as an increase of the global land carbon sink [Sarmiento et al. 2010]. The observed progressively greater differences between annual peak to trough flows may have large implications for both the functioning of the natural ecosystems and the people living along the Amazon floodplain and depending upon them for their livelihood.

[16] **Acknowledgments.** M. G. was financially supported by the NERC consortium grant AMAZONICA (NE/F005806/1) and the EU grant 7th framework GEOCARBON project (grant number agreement 283080). R. J. W. B. was supported by the Gordon and Betty Moore Foundation and a research Fellowship from the U.K. Natural Environment Research Council (NERC). O. P. and T. F. were supported by AMAZONICA (NERC) and the Gordon and Betty Moore Foundation and O. P. by an ERC Advanced Grant (T-FORCES). We thank Peter Knippertz, Piers Forster, Paul Palmer, and Yadvinder Malhi for comments on the manuscript.

References

- Beck, C., J. Grieser, and B. Rudolf (2005) A new monthly precipitation climatology for the global land areas for the period 1951 to 2000, DWD, Klimastatusbericht 2004, 181–190.
- Booth, B. B. B., C. D. Jones, M. Collins, I. J. Totterdell, P. M. Cox, S. Sitch, C. Huntingford, R. A. Betts, G. R. Harris, and J. Lloyd (2012), High sensitivity of future global warming to land carbon cycle processes, *Environ. Res. Lett.*, 7, 1–8.
- Callède, J., J. L. Guyot, J. Ronchail, M. Molinier, and E. de Oliveira (2002), L’Amazonie à Obidos (Brésil), Etude statistique des débits et bilan hydrologique, *Hydrol. Sci. J.*, 47, 321–333.
- Callède, J., J. L. Guyot, J. Ronchail, Y. L’Hôte, H. Niel, and E. De Oliveira (2004), Evolution du débit de l’Amazonie à Obidos de 1903 à 1999, *Hydrol. Sci. J.*, 49, 85–97.
- Callède, J., G. Cochonneau, F. Vieira Alves, J.-L. Guyot, V. Santos Guimaraes, and E. De Oliveira (2010), The River Amazon water contribution to the Atlantic Ocean, *J. Water Sci.*, 23(3), 247–273.
- Costa, M. H., and J. A. Foley (1999), Trends in the hydrologic cycle of the Amazon basin, *J. Geophys. Res.*, 104(D12), 14,189–14,198.
- Costa, M. H., M. T. Coe, and J.-L. Guyot (2009), Effects of Climatic Variability and Deforestation on surface water regimes, in *Amazonia and Global Change*, Geophys. Mon. Ser. 186, edited by M. Keller,

- M. Bustamante, J. Gash, and P. Silva Dias, 543–553, American Geophysical Union, Washington, DC.
- Dai, A., T. Qian, K. E. Trenberth, and J. D. Milliman (2009), Changes in continental freshwater discharge from 1948 to 2005, *J. Clim.*, 2773–2792, doi:http://dx.doi.org/10.1175/2008JCLI2592.1.
- Durack, P. J., S. E. Wijffels, R. J. Matear (2012), Ocean salinities reveal strong global water cycle intensification during 1950 to 2000, *Science*, 336, 455–458, doi:10.1126/science.1212222.
- Doughty, C. E., S. R. Loarie, and C. B. Field (2012), Theoretical impact of changing albedo on precipitation at the southernmost boundary of the ITCZ in South America, *Earth Interact.*, 16, 1–14, doi:http://dx.doi.org/10.1175/2012EI422.1.
- Espinoza, Villar J. C., J. L. Guyot, J. Ronchail, G. Cochonneau, N. P. Filizola, P. Fraizy, D. Labat, E. De Oliveira, J. J. Ordoñez Gálvez, and P. Vauchel (2009), Contrasting regional discharge evolutions in the Amazon basin (1974–2004), *J. Hydrol.*, 375(3–4), 297–311.
- Espinoza Villar, J. C., J. Ronchail, J. L. Guyot, C. Junquas, P. Vauchel, W. S. Lavado Casimiro, G. Drapeau, and R. Pombosa Loza (2011), Climate variability and extreme drought in the upper Solimões River (Western Amazon basin): Understanding the exceptional 2010 drought, *Geophys. Res. Lett.*, 38L13406.
- Espinoza, J. C., J. Ronchail, J. L. Guyot, C. Junquas, G. Drapeau, and J. M. Martinez (2012), From drought to flooding: understanding the abrupt 2010–11 hydrological annual cycle in the Amazonas River and tributaries, *Environ. Res. Lett.*, 7, doi:10.1088/1748-9326/7/2/024008.
- Gentry, A. H., and J. Lopez-Parodi (1980), Deforestation and increased flooding of the Upper Amazon, *Science*, 210, 1354–1356.
- Haylock, M. R. et al. (2006), Trends in total and extreme South American rainfall in 1960–2000 and links with sea surface temperature, *J. Clim.*, 19, 1490–1512.
- Held, I., and B. Soden (2006), Robust responses of the hydrological cycle to global warming, *J. Clim.*, 19, 5686–5699.
- Lewis, S. L., P. M. Brando, O. L. Phillips, G. M. F. van der Heijden, and D. Nepstad (2011), The 2010 Amazon Drought, *Science*, 331, doi:10.1126/science.1200807.
- Malhi, Y. et al. (2006), The regional variation of aboveground live biomass in old-growth Amazonian forests, *Global Change Biol.*, 12, 1–32, doi:10.1111/j.1365-2486.2006.01120.x.
- Malhi, Y., L. E. O. C. Aragão, D. Galbraith, C. Huntingford, R. Fisher, P. Zelazowski, S. Sitch, C. McSweeney, and P. Meir (2009), Exploring the likelihood and mechanism of a climate-change-induced dieback of the Amazon rain forest, *Proc. Natl. Acad. Sci. U. S. A.*, 106, 20610–20615.
- Marengo, J. A., C. A. Nobre, J. Tomasella, M. F. Cardoso, and M. D. Oyama (2008), Hydro-climatic and ecological behaviour of the drought of Amazonia in 2005, *Philos. Trans. R. Soc.*, B 2008, 363, 1773–1778, doi:10.1098/rstb.2007.0015.
- Marengo, J. A., J. Tomasella, L. M. Alves, W. R. Soares, and D. A. Rodriguez (2011), The drought of 2010 in the context of historical droughts in the Amazon region, *Geophys. Res. Lett.*, 38, L12703 doi:10.1029/2011GL047436.
- Miller, A. J. (1994), The 1976/1977 climate shift of the Pacific Ocean, *Oceanogr.*, 7, 21–26.
- Mitchell, T. D., and P. D. Jones (2005), An improved method of constructing a database of monthly climate observations and associated high-resolution grids, *Int. J. Climatol.*, 25, 693–712.
- Nordin, C. F., and R. H. Meade (1982), Deforestation and increased flooding of the upper Amazon, *Science*, 215, 426.
- Salati, E., A. Dall'Olio, E. Matsui, and J. R. Gat (1979), Recycling of water in the Amazon basin: An isotopic study, *Water Res. Res.*, 15, 1250–1258.
- Santer, B. D. et al. (2007), Identification of human-induced changes in atmospheric moisture content, *Proc. Nat. Acad. Sci.*, 104, 15248–15253.
- Sarmiento, J. L., M. Gloor, N. Gruber, C. Beaulieu, A. R. Jacobsen, S. Mikaloff Fletcher, S. Pacala, and K. Rodgers (2010), Trends and regional distributions of land and ocean carbon sinks, *Biogeoscience*, 7, 351–2367. doi:10.5194/bg-7-2351-2010.
- Smith, T. M., R. W. Reynolds, T. C. Peterson, and J. Lawrimote (2008), Improvements to NOAA's historical merged land-ocean surface temperature analysis (1880–2006), *J. Clim.*, 21, 2283–2296.
- Stohl, A., and P. James (2005), Lagrangian analysis of the atmospheric branch of the global water cycle. Part II: Moisture transports between Earth's ocean basins and river catchments, *J. Hydromet.*, 961–984.
- Trenberth, K. E. (2011), Changes in precipitation with climate change, *Clim. Res.*, 47, 123–138.
- Yoon, J.-H., and N. Zeng (2010), An Atlantic influence on Amazon rainfall, *Clim. Dyn.*, 34, 249–264. doi:10.1007/s00382-009-0551-6.