

A Set of 100 Chloroplast DNA Primer Pairs to Study Population Genetics and Phylogeny in Monocotyledons

Nora Scarcelli^{1*}, Adeline Barnaud¹, Wolf Eiserhardt², Urs A. Treier², Marie Seveno¹, Amélie d'Anfray¹, Yves Vigouroux¹, Jean-Christophe Pintaud¹

1 UMR Diversité, Adaptation et Développement des Plantes (DIADE), Institut de Recherche pour le Développement (IRD), Montpellier, France, **2** Ecoinformatics and Biodiversity Group, Department of Biological Science, Aarhus University, Aarhus, Denmark

Abstract

Chloroplast DNA sequences are of great interest for population genetics and phylogenetic studies. However, only a small set of markers are commonly used. Most of them have been designed for amplification in a large range of Angiosperms and are located in the Large Single Copy (LSC). Here we developed a new set of 100 primer pairs optimized for amplification in Monocotyledons. Primer pairs amplify coding (exon) and non-coding regions (intron and intergenic spacer). They span the different chloroplast regions: 72 are located in the LSC, 13 in the Small Single Copy (SSC) and 15 in the Inverted Repeat region (IR). Amplification and sequencing were tested in 13 species of Monocotyledons: *Dioscorea abyssinica*, *D. praehensilis*, *D. rotundata*, *D. dumetorum*, *D. bulbifera*, *Trichopus sempervirens* (Dioscoreaceae), *Phoenix canariensis*, *P. dactylifera*, *Astrocaryum scopatum*, *A. murumuru*, *Ceroxylon echinulatum* (Arecaceae), *Digitaria excilis* and *Pennisetum glaucum* (Poaceae). The diversity found in *Dioscorea*, *Digitaria* and *Pennisetum* mainly corresponded to Single Nucleotide Polymorphism (SNP) while the diversity found in Arecaceae also comprises Variable Number Tandem Repeat (VNTR). We observed that the most variable loci (*rps15-ycf1*, *rpl32-ccsA*, *ndhF-rpl32*, *ndhG-ndhI* and *ccsA*) are located in the SSC. Through the analysis of the genetic structure of a wild-cultivated species complex in *Dioscorea*, we demonstrated that this new set of primers is of great interest for population genetics and we anticipate that it will also be useful for phylogeny and bar-coding studies.

Citation: Scarcelli N, Barnaud A, Eiserhardt W, Treier UA, Seveno M, et al. (2011) A Set of 100 Chloroplast DNA Primer Pairs to Study Population Genetics and Phylogeny in Monocotyledons. PLoS ONE 6(5): e19954. doi:10.1371/journal.pone.0019954

Editor: Jonathan H. Badger, J. Craig Venter Institute, United States of America

Received: January 17, 2011; **Accepted:** April 7, 2011; **Published:** May 26, 2011

Copyright: © 2011 Scarcelli et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study was funded by EU Seventh Framework Programme as part of the project FP7-PALMS (grant agreement 212631) and by annual research budget of IRD. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: nora.scarcelli@ird.fr

Introduction

The knowledge of the chloroplast genome structure and sequence variation in Monocotyledons is still partial and unbalanced. There are currently 25 completely sequenced chloroplast genomes of Monocotyledons available in GenBank [1] but 17 of them are of Poales, and important orders like Liliales, Commelinales and Zingiberales lack complete chloroplast sequences. Comparative genomic analyses of the chloroplast DNA (cpDNA) relevant to Monocotyledons are scarce [2,3,4] and mostly focused on grasses and allied groups [5,6,7,8,9,10,11]. A few monocotyledonous species are documented for many genes [12] while numerous species are documented for a few genes only and non-coding regions, including the most commonly used markers for phylogenetic inference and genetic bar-coding like *rbcL* [13], *atpB* [14], *trnL-F* [15], *matK*, *psbA-trnH*, *rpoC1*, *rpoB-trnC*, *psbK-psbI*, *atpF-atpH*, *atpH-atpI* [16,17,18,19,20,21]. Most other regions of the Large Single Copy (LSC) have been investigated in particular taxa, for example *clpP* intron2 in *Yucca* [22], *rps2* in *Tiphonium* [23], *petN-psbM* in *Elaeocharis* [24], *psbD-trnT* in *Arum* [25], *psbB* and *psbC* in *Vanilla* [26], *psbZ-trnM* in *Livistona* [27], or *accD* in *Hexalectris* [28], to mention a few studies. Variation within the slowly evolving Inverted Repeat region (IR) [29] has received little attention in Monocotyledons [30,31], a large part of it being represented only by the complete chloroplast sequences. Within

the Small Single Copy (SSC), there is limited information outside the extensively used *ndhF* gene [32], with only few studies using *ycf1*, *rpl32-trnL* and *ndhA* [33,34,35,36].

Moreover, available sets of primers for direct sequencing of chloroplast regions in Angiosperms mostly focus on non-coding regions of the LSC [37,38,39] while published information on primers for genes is very dispersed [40].

The possibility of screening a large number of loci is useful to detect polymorphic Small Inversions, microsatellites and minisatellites (i.e. Variable Number Tandem Repeats, VNTR) in species complexes and at the population level [1,37]. Minute and medium size inversions are frequent features of the non-coding cpDNA [41,42], detectable only through sequencing and showing intraspecific variability [1,43]. Microsatellites are also widespread structures in non-coding cpDNA that became important population genetics markers [44]. The most common and most widely used microsatellites are mononucleotide repeats [45]. Longer motifs, in particular minisatellites, are comparatively rare, but also proved to be valuable markers [46,47,48].

Here we propose a large set of primer pairs optimized for PCR amplification and overlapping sequencing in Monocotyledons. Primer pairs are distributed throughout the whole chloroplast genome and include exons, introns and Intergenic Spacers (IGS) with contrasted mutation rates and evolutionary patterns. They are thus suitable for a wide range of studies from higher-level

phylogeny to population genetics. As an example, we used the newly defined primer pairs to study intra-specific cpDNA diversity of three different yam species (*Dioscorea* spp.).

Materials and Methods

Primer definition

The complete sequence of six Monocotyledons chloroplast genomes were downloaded from GenBank, namely *Acorus calamus* (NC_007407), *Dioscorea elephantipes* (NC_009601), *Lemna minor* (NC_010109), *Oryza nivara* (NC_005973), *Phalaenopsis aphrodite* (NC_007499) and *Zea mays* (NC_001666).

Segments of these sequences equivalent to two to six genes were aligned using the program GENEIOUS [49]. Consensus primers anchored in exons were designed using Primer3 [50] incorporated in GENEIOUS, in order to amplify IGS, introns or exons. A total of 105 primer pairs were designed, and 100 successfully amplified: 72 in the Large Single Copy region (LSC), 13 in the Small Single Copy (SSC) and 15 in the Inverted Repeat region (IR). Primer sequences, annealing temperature for PCR amplification, and amplification results are summarized in Table S1.

Test for amplification

Amplification was tested in 13 species of Monocotyledons: 6 Dioscoreaceae species (1 individual each of *Dioscorea abyssinica*, *D. praehensilis*, *D. rotundata*, *D. dumetorum*, *D. bulbifera* and *Trichopus sempervirens*), 5 Arecaceae species (1 individual each of *Phoenix canariensis*, *P. dactylifera*, *Astrocaryum scopatum*, *A. murumuru* and 2 individuals of *Ceroxylon echinulatum*), *Digitaria excilis* (5 individuals) and *Pennisetum glaucum* (6 individuals). Sequences have been deposited in GenBank under accession number JF705257-JF705858, JF745569-JF745769 and JF758190-JF758233.

Amplification was done according to the recommended protocols using either GoTaq (Promega) in its buffer with 5 mM of dNTPs for *D. excilis* and *P. glaucum* or Failsafe enzyme mix (Epicentre) in premix E for Dioscoreaceae and Arecaceae species. Reaction was done in 25 μ L with 25 ng of DNA. The initial denaturation (94°C, 3 min) was followed by 35 cycles of denaturation (94°C, 30 s), annealing (T_m , 30 s) and elongation (72°C, 1 min) and by a final elongation step (72°C, 10 min). Amplification was checked on agarose gel.

Sequencing

PCR products were purified using Ampure (Agencourt) following the recommended protocol. The sequencing PCRs were done using the BigDye terminator kit (Applied Biosystems). PCR products were purified using CleanSeq (Agencourt) and were run on ABI prism 3130 (Applied Biosystems). Note that for *D. excilis* and *P. glaucum*, only a subset of the PCR products was sequenced. Dioscoreaceae and Arecaceae species were sequenced in forward and reverse direction while *D. excilis* and *P. glaucum* were sequenced in forward direction only.

Data analysis

Sequences were aligned with the program GENEIOUS [49]. Intra-generic diversity was estimated within species or between closely-related species as the number of SNP and the number of Variable Number Tandem Repeats (VNTR). The number of SNP was standardised to 1 kb but length variable parts (e.g., gaps or VNTR) were subtracted from the total length of the alignment. For *D. excilis* and *P. glaucum*, intra-generic diversity was estimated within the analysed species. For Dioscoreaceae, it was estimated between closely-related species (*D. rotundata*, *D. abyssinica* and *D. praehensilis* of subgenus *Entantophyllum*). For Arecaceae it was

estimated as the mean of the diversity found within each species pair in *Phoenix*, *Astrocaryum* and *Ceroxylon*.

Due to high inter-generic divergence in Dioscoreaceae [51] causing alignment difficulties in non-coding regions with *T. sempervirens*, nucleotide diversity was only estimated between two distant species of *Dioscorea*, *D. abyssinica* and *D. elephantipes*. For Arecaceae, an average of three inter-generic comparison, between *Phoenix* (subfamily Coryphoideae) and *Astrocaryum* (subfamily Arecaceae), *Phoenix* and *Ceroxylon* (subfamily Ceroxyloideae) and *Ceroxylon* and *Astrocaryum* was calculated. Within Poaceae, inter-generic diversity was not estimated for *D. excilis* and *P. glaucum* because only a part of the loci were sequenced. Instead, inter-generic diversity was estimated between *Oryza sativa* and *Zea mays* using the GenBank sequences but restricted to those parts theoretically amplified by the primer pairs tested in the present study.

Comparisons of genetic diversity between SSC, LSC and IR, and between introns, exons and IGS were performed with Kruskal-Wallis tests using the R environment [52], function `kruskal.test`.

Example of use for population genetic analysis

We analysed the genetic structure of three yam species (*Dioscorea* spp.) forming a crop-wild relatives complex in Western Africa. The main cultivated yam species in West-Africa is *D. rotundata*. In this region, yam is a staple food but is also culturally extremely important [53]. The wild relatives of *D. rotundata* are *D. abyssinica* and *D. praehensilis* [54,55]. The three species are genetically different but can hybridize [56].

One sample of each species has been previously sequenced (see above). Based on these sequences, 19 polymorphic loci were identified showing a total of 21 SNP. These 19 loci have been tested on eight additional individuals (four *D. abyssinica* and four *D. praehensilis*) to selected those loci for which polymorphisms were specific to either *D. abyssinica* or *D. praehensilis*; namely *ccsA*-Exon, *ccsA-ndhD*, *ndhH*-Exon, *psbD*-Exon and *rm4,5-trnN*.

Finally, a total of 160 *Dioscorea* samples have been amplified using the selected five primers pairs. The sampling included 66 *D. abyssinica*, 39 *D. praehensilis* and 55 *D. rotundata* collected in Benin. A list of individuals and sampling locations is given in the supplementary data file (Table S2). Sequences have been deposited in GenBank under accession number JF757240-JF758189. The five loci revealed six SNP (two for *rm4,5-trnN* and one each for the other loci). A chlorotype is defined as a combination of SNP located on the chloroplast, i.e. a haplotype based on chloroplast SNP. Here, the combinations of the six SNP revealed five chlorotypes. The repartition of chlorotype frequencies among species was compared with a chi-squared test. A MSN, Minimum Spanning Network [57], with chlorotypes was constructed using Haplophyle [58]. MSN illustrates the evolutionary relationships between chlorotypes as a network where the branches represent the differences between sequences data.

Results and Discussion

Development of new chloroplast primers

Of the 105 primer pairs designed to sequence the chloroplast genome, 100 amplified consistently and produced good quality sequences. Primers were designed to amplify a wide range of monocotyledons species and we tested them on various species of different genera (*D. abyssinica*, *D. praehensilis*, *D. rotundata*, *D. dumetorum*, *D. bulbifera*, *T. sempervirens*, *P. canariensis*, *P. dactylifera*, *A. scopatum*, *A. murumuru*, *C. echinulatum*, *D. excilis*, *P. glaucum*). Amplification success was 85% (Table S1) which was very similar

to the expected mean amplification of 88% derived from the sequences deposited in GenBank used to design the primers (95% for *A. calamus*, 95% for *D. elephantipes*, 97% for *L. minor*, 80% for *O. nivara*, 88% for *P. aphrodite* and 80% for *Z. mays*). Indeed, due to structural changes (inversions, gene loss, etc.) some primers pairs are expected not to amplify in some species. For example, because of the loss of *ycf2* and *accD* in *O. nivara* and *Z. mays*, we do not expect amplification with primers pairs *rpl23-ycf2*, *ycf2-ndhB*, *accD-psaI*, *rbcL-accD* and *accD-Exon* on these two species.

Primers amplified coding regions (exon 20%), non-coding regions (IGS 35%, intron 9%) and mixed regions (exon+intron 10%, IGS+genes 25%). 75% of these regions were located in the LSC, 12% in the SSC and 15% in the IR.

Sequence diversity

We obtained a total of 1174 kb sequence data. The analysis covered 78 kb of the chloroplast genome for Dioscoreaceae (51% of the *D. elephantipes* cpDNA), 70 kb for Arecaceae (44% of the *P. dactylifera* cpDNA), 34 kb for *Digitaria* (25% of the *O. nivara* cpDNA) and 20 kb for *Pennisetum* (15% of the *O. nivara* cpDNA).

A summary of intra- and inter-generic diversity results are presented in Table 1. Detailed results are given in supplementary data file (Table S3).

Intra-generic diversity

We found on average a SNP each 1700 bp within the three Arecaceae genera, each 2800 bp between the three *Dioscorea* species *D. abyssinica*, *D. prahensis* and *D. rotundata*, each 8900 bp among the six *P. glaucum* samples and each 9600 bp among the five *D. excilis* samples. These very low levels of intra-generic diversity in the studied Poaceae suggest a strong bottleneck effects in such cultivated populations. There were few polymorphic microsatellites in *Dioscorea*, *D. excilis* and *P. glaucum*, all mononucleotide, while the Arecaceae exhibit a high number of mono-, di- and 4–8-nucleotide microsatellites as well as minisatellites (Table 2). A total of 66 VNTR were found in palms, 77% of them located in IGS, 23% in intron and none in exon (Table 3). The 51 polymorphic mononucleotide microsatellites encountered within genera and species of palms can be compared with the 342 homopolymers of 7 bp or longer found in the complete chloroplast genome of *Phoenix dactylifera* [1].

Interestingly, Arecaceae species exhibit a much higher number of VNTR than *Dioscorea* species. Similar levels of mono- and dinucleotide microsatellites in *Dioscorea* as in closely related palm species could only be found if two distant species (*D. abyssinica* and *D. elephantipes*) were compared (data not shown). This result suggests different evolutionary histories with higher mutation rates

Table 2. Comparison of the number and motifs of polymorphic VNTR observed at the intra-generic level.

	<i>Dioscorea</i>	<i>Digitaria</i>	<i>Pennisetum</i>	Arecaceae
n.o. bp sequenced	78134	33862	19746	69422
n.o. mononucleotide	9	0	7	51
n.o. dinucleotide	0	0	0	2
n.o. VNTR with motif >3 bp	0	0	0	13

doi:10.1371/journal.pone.0019954.t002

and/or larger effective population sizes in Arecaceae than in *Dioscorea* species.

Inter-generic diversity

Between *O. sativa* and *Z. mays* we found a SNP each 21 bp, between *D. abyssinica* and *D. elephantipes* each 75 bp and for the three inter-generic comparisons in Arecaceae on average each 113 bp. Since *Oryza sativa* and *Zea mays* diverged about 52 MY ago [59] and the compared palm subfamilies diverged about 68–98 MY ago [60], our result confirmed a 5–6 fold faster substitution rate for cpDNA in Poaceae than in Arecaceae [61]. The genus concept in Dioscoreaceae is very different from that of Poaceae and Arecaceae. Levels of divergence between two distant species of *Dioscorea* was in the range of the inter-generic differentiation in Poaceae and Arecaceae, while different Dioscoreaceae genera, namely *Dioscorea* and *Trichopus* are so divergent that they are not even alignable for some IGS.

Interestingly, we did not find significant differences in number of SNP in introns *vs.* exons and in introns *vs.* IGS, neither for *Dioscorea*, nor for Arecaceae and Poaceae ($p > 0.05$ for *Dioscorea*, Arecaceae and Poaceae). We observed a significantly higher number of SNP in IGS *vs.* exons only for *Dioscorea* ($p < 0.05$) and in Poaceae ($p < 0.01$). This finding highlights the very peculiar dynamics of SNP in the chloroplast genome. It can be compared with the result of Yang et al. [1] who identified 62 out of 78 SNP within the cultivar ‘Khalass’ of the date palm occurring in exons, with an unusual synonymous/non synonymous ratio of 0.94. They suggested a lack of purifying selection within heterogeneous intra-individual chloroplast populations as a possible explanation (Yang et al. 2010b).

The occurrence of SNP among the three regions of the chloroplast (LSC, SSC and IR) varies (Figure 1). LSC and SSC exhibit similar levels of diversity while IR exhibits significantly lower numbers of SNP. The difference in number of SNP is

Table 1. Observed intra- and inter-generic diversity.

Location	Intra-generic diversity				Inter-generic diversity						
	SNP				VNTR				SNP		
	<i>Dioscorea</i>	<i>Digitaria</i>	<i>Pennisetum</i>	Arecaceae	<i>Dioscorea</i>	<i>Digitaria</i>	<i>Pennisetum</i>	Arecaceae	<i>Dioscorea</i>	Poaceae	Arecaceae
LSC	0.18	0	0.34	0.59	0.13	0	0.33	1.02	14.26	62.82	9.07
IR	0.06	0	0	0.16	0	0.13	0.20	0	2.11	11.16	1.44
SSC	0.75	0.31	0	1.07	0.09	0	0	1	23.37	68.84	10.10
Mean	0.24	0.10	0.11	0.57	0.07	0.04	0.18	0.87	13.25	47.61	8.07

Intra-generic diversity was estimated between closely-related species for *Dioscorea* and Arecaceae or within species (*Digitaria excilis* and *Pennisetum glaucum*). Inter-generic diversity was estimated between different genera (Arecaceae and Poaceae) or distant species (*Dioscorea*). The number of SNP was standardised to 1 kb.

doi:10.1371/journal.pone.0019954.t001

Table 3. Polynucleotide VNTRs with repeat number >2 and polymorphic within genera in palms.

Locus	Motif length	Motif sequence	Number of repeats		
			<i>Phoenix</i>	<i>Ceroxylon</i>	<i>Astrocaryum</i>
trnL intron	2 bp	AT	6	6	8–10
trnL-ndhJ	2 bp	AT	5	6	6–7
trnQ-rps16	4 bp	GATA	3	2	3–4
ndhG-ndhI	5 bp	AAATA	3	3	2–3
trnQ-rps16	6 bp	AATATT	2	2	2–3
rbcl-accD	8 bp	TTACTTAT	1	1	2–3
psbZ-trnfM	12 bp	ACTACTATACTA	2–6	2	3
rpl16-rps3	20 bp	CTCGTTTACAAATCCAAA	2–3	1	1–2

doi:10.1371/journal.pone.0019954.t003

significant for LSC *vs.* IR and SSC *vs.* IR ($p < 0.001$ for *Dioscorea*, *Arecaceae* and *Poaceae*) but is not significant for LSC *vs.* SSC ($p > 0.05$ for *Dioscorea*, *Arecaceae* and *Poaceae*). Variation in SNP number in the SSC region is, however, mostly driven by the *ndhF-rpl32* locus. This locus exhibits a very high genetic diversity: 92 and 118 SNP per 1 kb for *Dioscorea* and *Poaceae*, compared to the mean of 16, 10 and 62 SNP per 1 kb for *Dioscorea*, *Arecaceae* and *Poaceae*, respectively and of 11, 7 and 30 SNP per 1 kb in the whole chloroplast.

Most published primer pairs focus on non-coding regions of the LSC [37,38,39]. This region is commonly used for phylogeny and bar-coding [13,14,15,16,17,18,19,20]. In the present study, we observed some of the most variable loci in the SSC, namely *rps15-ycf1*, *rpl32-ccsA* and *ndhF-rpl32* for *Dioscorea* and *rps15-ycf1*, *ndhG-ndhI* and *ccsA* for *Arecaceae*.

Polynucleotide VNTR in palms

Polynucleotide VNTR are apparently rare in Monocotyledons. They are virtually absent in *Dioscorea* and *Poaceae*, although a 22 bp minisatellite located in the *tmD-trnT* region, with 1–3 repeats, has been reported in *Elymus* [47]. A complex evolution of minisatellites was also detected in an orchid, *Anacamptis*, within the *trnL* intron [62].

Palms are outstanding for the frequency of such structures in the chloroplast genome. In this study, 12 VNTR were recorded in the

genus *Astrocaryum*, with motif length varying from 2 to 26 bp. There was, however, considerable variation in VNTRs abundance among genera of palms (Table 3). In *Phoenix*, only two polynucleotide VNTRs were detected, namely 2 minisatellites of 12 and 20 bp. Within *C. echinulatum*, there was no polymorphism at the level of the polynucleotide VNTR, and only 9 of the 51 mononucleotide microsatellites were polymorphic. We note, however, that only two individuals have been compared and VNTRs occurrence might be higher. Differences between *Astrocaryum* and *Ceroxylon* might be explained by differences in divergence time between the pairs of individuals compared (less than 2 MY in *Ceroxylon*, about 7 MY in *Astrocaryum*) and also by a higher sequence variability in *Astrocaryum* and other Bactridinae compared with *Ceroxyleae* and *Phoeniceae* [63,64,65,66].

Thus, polynucleotide VNTRs have a great potential in palms for population genetic studies and species delimitation. They have already been used with success in several studies. For example, the dodecanucleotide minisatellite of the *psbZ-trnfM* locus showed fixed private haplotypes that allowed the separation of closely related *Phoenix* species and tracking interspecific hybridization [48]. The tetranucleotide microsatellite of the *trnQ-rps16* locus allowed tracing seed flow between the wild and cultivated compartments of the peach palm (*Bactris gasipaes*) in western Ecuador and proved to be much more informative than a mononucleotide microsatellite present in the same locus [67].




Figure 1. Inter-generic diversity found in *Dioscorea*, *Arecaceae* and *Poaceae*. Inter-generic diversity was estimated as the number of SNP in the LSC, SSC and IR. Numbers of SNP have been standardised to 1 kb. Bars represent the 95% confidence intervals.

doi:10.1371/journal.pone.0019954.g001

We note, however, that among the 13 polynucleotide VNTRs with motifs longer than 3 bp found in the palm sampling (Table 2), five are direct repeats, i.e. incipient VNTR with 1–2 units of the motif. Direct repeats are common in non-coding cpDNA [68], and are probably the starting point of more repeated polynucleotide VNTR, although few loci undergo this evolution. For example, a sequence of 8 bp in the *rbcL-accD* spacer was found unrepeated in *Phoenix* and *Ceroxylon* but showed 2–3 tandem repeats in *Astrocaryum* (Table 3). Some minisatellites also originate from inversions [62].

As already noted above, the comparison of a limited number of individuals per family, as in the present study, might considerably underestimate the actual number of VNTR in a given taxa. Indeed, an alignment of the 174 palm sequences deposited in GenBank of the locus *tmQ-rps16* alone (1.1 kb) revealed 16 intra-

generic direct repeat polymorphisms 5–22 bp long, a mononucleotide microsatellite with 8–17 repeats, a dinucleotide microsatellite with 3–6 repeats, a tetranucleotide microsatellite with 2–6 repeats and a 26 bp minisatellite with 1–4 repeats. The last structure is polymorphic in a single group, the subtribe *Linospadicinae*, restricted to the south-west Pacific [69].

For detailed studies of VNTR variation in a particular group, it is therefore advisable to begin with the sequencing of a significant number of samples, in order to evaluate accurately the existing polymorphism in the target locus.

Example of use for population genetic analysis

CpDNA is generally inherited by only one parent (usually the mother in angiosperms). It is haploid and it generally lacks




Figure 2. Chlorotypes observed within *Dioscorea* species. (A) Distribution of observed chlorotypes and (B) minimum spanning network (MSN) representing the relationship among chlorotypes. In (B), the size of the circle is proportional to the chlorotype occurrence and each line represents a SNP between the two connected chlorotypes. Each SNP has been labelled with the name of the primer pairs needed for its amplification and the corresponding base change. doi:10.1371/journal.pone.0019954.g002

recombination [70]. CpDNA is therefore of great interest for population genetics studies, including parentage analysis, hybridization, population structure and phylogeography [44].

Here we used the new primer set to study the genetic structure of a yam crop-wild relatives' complex (cultivated: *D. rotundata*, wild: *D. abyssinica* and *D. praeensis*) in Benin, Western Africa. After screening more than half of the chloroplast genome the set of informative loci retained to study this species complex included four (out of five) loci from the SSC and IR regions, among which three were exons. This emphasizes again how interesting the rarely studied SSC and IR regions are and confirms that exons are not less variable than introns or IGS, as far as it concerns SNP.

We found five chlorotypes among the 160 sequenced *Dioscorea* individuals that showed significantly different frequencies among the three species (Fig. 2a, $p < 0.001$ for all pairwise comparisons, chi-squared tests). Chlorotypes 2 and 3 were specific to *D. abyssinica*; chlorotypes 4 and 5 are specific to *D. praeensis*; while the most common chlorotype 1 was found in all three species. Chlorotypes 1, 2 and 3 as well as chlorotypes 4 and 5 were closely related with only one SNP separating them (Fig. 2b).

The cultivated species *D. rotundata* harboured only chlorotype 1 and thus was less diverse than its wild relatives *D. abyssinica* and *D. praeensis*. However, because chlorotype 1 was shared by all three species, we cannot conclude on the maternal origin of *D. rotundata*.

Our results showed that SNP revealed by sequencing can successfully be used to study the diversity of the crop-wild relatives' complex of *Dioscorea*. Furthermore, the genetic diversity revealed by sequencing with five primer pairs was more informative than the genetic diversity observed using five universal chloroplast mononucleotide microsatellites [71].

We thus showed that the new primer set can reveal diversity even when microsatellites might not show polymorphism, as it was the case in the *Dioscorea* species complex studied. We anticipate that the use of sequencing and SNP genotyping for population genetic analysis will be even more interesting for species or species complexes showing higher genetic diversity, as in some groups of Arecaceae like Bactridinae.

Conclusion

In this paper, we present a large set of newly developed chloroplast DNA primer pairs. Compared to the previously

published primer pairs [37,38,39,40], this new set covers a wider range of the chloroplast genome (e.g. up to 51% of the *Dioscorea* cpDNA) and has been designed to optimally amplify in Monocotyledons. This new set of primer pairs spans the Large Single Copy as well as the Small Single Copy and the Inverted Repeats, and has been designed to amplify both coding (exon) and non-coding (intron, intergenic spacer) regions. This new set could be of great interest for phylogeny and bar-coding studies but also for population genetics studies.

Supporting Information

Table S1 Primer sequences and amplification range. Primers were designed using genes alignment of *Dioscorea elephantipes*, *Zea mays*, *Oryza nivara*, *Lemna minor*, *Acorus calamus* and *Phalaenopsis aphrodite*. Amplifications were tested on different species of Dioscoreaceae, *Digitaria*, *Pennisetum* and Arecaceae. (DOC)

Table S2 List of *Dioscorea* individuals used to test the use of the new primers pairs for population genetics studies. Table includes the chlorotype (1 to 5) corresponding to each sample. (DOC)

Table S3 Observed Intra- and inter-generic diversity. Intra-generic diversity was estimated between closely-related species for *Dioscorea* and Arecaceae or within species (*Digitaria excilis* and *Pennisetum glaucum*). Inter-generic diversity was estimated between different genera (Arecaceae and Poaceae) or distant species (*Dioscorea*). The number of SNP was standardised to 1 kb. (DOC)

Acknowledgments

We greatly acknowledge S. De Mita for help in GenBank submission. We thank the editor and an anonymous reviewer for their useful comments.

Author Contributions

Conceived and designed the experiments: NS AB YV JCP. Performed the experiments: NS AB MS AdA. Analyzed the data: NS AB WE UAT. Contributed reagents/materials/analysis tools: NS AB YV JCP. Wrote the paper: NS AB YV JCP.

References

- Yang M, Zhang X, Liu G, Yin Y, Chen K, et al. (2010) The complete chloroplast genome sequence of date palm (*Phoenix dactylifera* L.). PLoS ONE 5: e12762.
- Hansen D, Dastidar S, Cai Z, Penafior C, Kuehl J, et al. (2007) Phylogenetic and evolutionary implications of complete chloroplast genome sequences of four early-diverging angiosperms: *Buxus* (Buxaceae), *Chloranthus* (Chloranthaceae), *Dioscorea* (Dioscoreaceae), and *Illicium* (Schisandraceae). Molecular Phylogenetics and Evolution 45: 547–563.
- Mardanov A, Ravin N, Kuznetsov B, Samigullin T, Antonov A, et al. (2008) Complete sequence of the duckweed (*Lemna minor*) chloroplast genome: structural organization and phylogenetic relationships to other Angiosperms. Journal of Molecular Evolution 66: 555–564.
- Chang C, Lin H, Lin I, Chow T, Chen H, et al. (2006) The chloroplast genome of *Phalaenopsis aphrodite* (Orchidaceae): Comparative analysis of evolutionary rate with that of grasses and its phylogenetic implications. Molecular Biology and Evolution 23: 279–291.
- Matsuoka Y, Yamazaki Y, Ogihara Y, Tsunewaki K (2002) Whole chloroplast genome comparison of rice, maize, and wheat: implications for chloroplast gene diversification and phylogeny of cereals. Molecular Biology and Evolution 19: 2084–2091.
- Saski C, Lee S, Fjellheim S, Guda C, Jansen R, et al. (2007) Complete chloroplast genome sequences of *Hordeum vulgare*, *Sorghum bicolor* and *Agrostis stolonifera*, and comparative analyses with other grass genomes. Theoretical and applied genetics 115: 571–590.
- Leseberg C, Duvall M (2009) The complete chloroplast genome of *Coix lacrym-jobi* and a comparative molecular evolutionary analysis of plastomes in cereals. Journal of Molecular Evolution 69: 311–318.
- Guisinger M, Chumley T, Kuehl J, Boore J, Jansen R (2010) Implications of the plastid genome sequence of *Typha* (Typhaceae, Poales) for understanding genome evolution in Poaceae. Journal of Molecular Evolution 70: 149–166.
- Morris L, Duvall M (2010) The chloroplast genome of *Anomochloa marantoides* (Anomochloideae; Poaceae) comprises a mixture of grass-like and unique features. American Journal of Botany 97: 620–627.
- Bortiri E, Coleman-Derr D, Lazo G, Anderson O, Gu Y (2008) The complete chloroplast genome sequence of *Brachypodium distachyon*: sequence comparison and phylogenetic analysis of eight grass plastomes. BMC Research Notes 1: 61.
- Cahoon A, Sharpe R, Mysayphonh C, Thompson E, Ward A, et al. (2010) The complete chloroplast genome of tall fescue (*Lolium arundinaceum*; Poaceae) and comparison of whole plastomes from the family Poaceae. American Journal of Botany 97: 49–58.
- Jansen R, Cai Z, Raubeson L, Daniell H, Depamphilis C, et al. (2007) Analysis of 81 genes from 64 plastid genomes resolves relationships in angiosperms and identifies genome-scale evolutionary patterns. Proceedings of the National Academy of Sciences of the United States of America 104: 19369–19374.
- Duvall M, Clegg M, Chase M, Clark W, Kress W, et al. (1993) Phylogenetic hypothesis for the Monocotyledons constructed from *rbcl* sequence data. Annals of the Missouri Botanical Garden 80: 607–619.
- Soltis D, Soltis P, Chase M, Mort M, Albach D, et al. (2000) Angiosperm phylogeny inferred from 18S rDNA, *rbcl*, and *atpB* sequences. Botanical Journal of the Linnean Society 133: 381–461.
- Taberlet P, Gicly L, Pautou G, Bouvet J (1991) Universal primers for amplification of three non-coding regions of chloroplast DNA. Plant Molecular Biology 17: 1105–1109.

16. Tamura M, Yamashita J, Fuse S, Haraguchi M (2004) Molecular phylogeny of monocotyledons inferred from combined analysis of plastid *matK* and *rbcL* gene sequences. *Journal of Plant Research* 117: 109–120.
17. Kress J, Wurdack K, Zimmer E, Weigt L, Janzen D (2005) Use of DNA barcodes to identify flowering plants. *Proceedings of the National Academy of Sciences of the United States of America* 102: 8369–8374.
18. Chase M, Cowan R, Hollingsworth P, van den Berg C, Madriñán S, et al. (2007) A proposal for a standardised protocol to barcode all land plants. *Taxon* 56: 295–299.
19. Zeng C, Zhang Y, Triplett J, Yang J, Li D (2010) Large multi-locus plastid phylogeny of the tribe Arundinarieae (Poaceae: Bambusoideae) reveals ten major lineages and low rate of molecular divergence. *Molecular Phylogenetics and Evolution* 56: 821–839.
20. Hollingsworth P, Forrest L, Spouge J, Hajibabaei M, Ratmsingham S, et al. (2009) A DNA barcode for land plants. *Proceedings of the National Academy of Sciences of the United States of America* 106.
21. Seberg O, Petersen G (2009) How many loci does it take to DNA barcode a *Crocus*? *PLoS ONE*. pp e4598.
22. Smith C, Pellmyr O, Althoff D, Balcazar-Lara M, Leebens-Mack J, et al. (2008) Pattern and timing of diversification in *Yucca* (Agavaceae): specialized pollination does not escalate rates of diversification. *Proceeding of the Royal Society London Series B* 275: 249–258.
23. Ohi-Toma T, Wu S, Yadav S, Murata H, Murata J (2010) Molecular phylogeny of *Typhonium sensu lato* and its allied genera in the tribe Aroideae of the subfamily Aroideae (Araceae) based on sequences of six chloroplast regions. *Systematic Botany* 35: 244–251.
24. Hinchliff C, Roalson E (2009) Stem architecture in *Eleocharis* subgenus *Linnochloa* (Cyperaceae): evidence of dynamic morphological evolution in a group of pantropical sedges. *American Journal of Botany* 96: 1487–1499.
25. Espindola A, Buerki S, Bedalov M, Kupfer P, Alvarez N (2010) New insights into the phylogenetics and biogeography of *Arum* (Araceae): unravelling its evolutionary history. *Botanical Journal of the Linnean Society* 163: 14–32.
26. Cameron K, Molina M (2006) Photosystem II gene sequences of *psbB* and *psbC* clarify the phylogenetic position of *Vanilla* (Vanilloideae, Orchidaceae). *Cladistics* 22: 239–248.
27. Crisp M, Isagi Y, Kato Y, Cook L, Bowman D (2009) *Livistona* palms in Australia: ancient relics or opportunistic immigrants? *Molecular Phylogenetics and Evolution* 54: 512–523.
28. Kennedy A, Watson L (2010) Species delimitations and phylogenetic relationships within the fully myco-heterotrophic *Hexaletris* (Orchidaceae). *Systematic Botany* 35: 64–76.
29. Goremykin V, Bobrova V, Pahnke J, Troitsky A, Antonov A, et al. (1996) Noncoding sequences from the slowly evolving chloroplast Inverted Repeat in addition to *rbcL* data do not support Gnetalean affinities of Angiosperms. *Molecular Biology and Evolution* 13: 383–396.
30. Graham S, Zgurski J, McPherson M, Cherniawsky D, Saarela J, et al. (2006) Robust inference of Monocots deep phylogeny using an expanded multigene plastid data set. *Aliso* 22: 3–21.
31. Wang R, Cheng C, Chang C, Wu C, Su T, et al. (2008) Dynamics and evolution of the inverted repeat-large single copy junctions in the chloroplast genomes of monocots. *BMC Evolutionary Biology* 8: 36.
32. Olmstead R, Sweere J, Wolfe K (1993) Ninety extra nucleotide in *ndhF* gene of tobacco chloroplast DNA: a summary of revisions to the 1986 genome. *Plant Molecular Biology* 22: 1191–1193.
33. Yang J-B, Yang H-Q, Li D-Z, Wong K-M, Yang Y-M (2010) Phylogeny of *Bambusa* and its allies (Poaceae: Bambusoideae) inferred from nuclear GBSSI gene and plastid *psbA-trnH*, *rpl32-trnL* and *rps16* intron DNA sequences. *Taxon* 59: 1102–1110.
34. Chase M, Williams N, de Faria A, Neubig K, Amaral Mdo C, et al. (2009) Floral convergence in Oncidiinae (Cymbidiaceae; Orchidaceae): an expanded concept of *Gomesa* and a new genus *Nohavilliamsia*. *Annals of Botany* 104: 387–402.
35. Peterson P, Romaschenko K, Johnson G (2010) A classification of the Chloridoideae (Poaceae) based on multi-gene phylogenetic trees. *Molecular Phylogenetics and Evolution* 55: 580–598.
36. Barrett C, Freudenstein J (2009) Patterns of Morphological and Plastid DNA Variation in the Corallorhiza striata Species Complex (Orchidaceae). *Systematic Botany* 34: 496–504.
37. Ebert D, Peakall R (2009) A new set of universal de novo sequencing primers for extensive coverage of noncoding chloroplast DNA: new opportunities for phylogenetic studies and cpSSR discovery. *Molecular Ecology Resources* 9: 777–1075.
38. Shaw J, Lickey E, Beck J, Farmer S, Liu W, et al. (2005) The tortoise and the hare II: relatively utility of 21 noncoding chloroplast DNA sequences for phylogenetic analysis. *American Journal of Botany* 92: 142–166.
39. Shaw J, Lickey E, Schilling E, Small R (2007) Comparison of whole chloroplast genome sequences to choose noncoding regions for phylogenetic studies in Angiosperms: The tortoise and the hare III. *American Journal of Botany* 94: 275–288.
40. Heinze B (2007) A database of PCR primers for the chloroplast genomes of higher plants. *Plant Methods* 3: 4.
41. Kelchner S, Wendel J (1996) Hairpins create minute inversions in non-coding regions of chloroplast DNA. *Current Genetics* 30: 259–262.
42. Kim K, Lee H-L (2005) Widespread occurrence of small inversions in the chloroplast genomes of land plants. *Molecules and Cells* 16: 104–113.
43. Whitlock B, Hale A, Groff P (2010) Intraspecific inversions pose a challenge for the trnH-psbA plant DNA barcode. *PLoS ONE* 5: e11533.
44. Provan J, Powell W, Hollingsworth P (2001) Chloroplast microsatellites: new tools for studies in plant ecology and evolution. *Trends in Ecology and Evolution* 16: 142–147.
45. Provan J, Biss P, McMeel D, Mathews S (2004) Universal primers for the amplification of chloroplast microsatellites in grasses (Poaceae). *Molecular Ecology Notes* 4: 262–264.
46. Cozzolino S, Cafasso D, Pellegrino G, Musacchio A, Widmer A (2003) Fine-scale phylogeographical analysis of Mediterranean *Anacamptis palustris* (Orchidaceae) populations based on chloroplast minisatellite and microsatellite variation. *Molecular Ecology* 12: 2783–2792.
47. Sun G, Ma X (2009) Nucleotide diversity and minisatellite in chloroplast Asp(GUC)-Thr(GGU) region in *Elymus trachycalus* complex, *Elymus alaskanus* and *Elymus caninus*. *Biochemical Systematics and Ecology* 37: 67–75.
48. Pintaud J-C, Zehdi S, Couvreur T, Barrow S, Henderson S, et al. (2010) Species delimitation in the genus *Phoenix* (Arecaceae) based on SSR markers, with emphasis on the identity of the Date Palm (*Phoenix dactylifera* L.). In: Seberg O, Petersen G, Barfod A, Davis J, eds. Diversity, phylogeny, and evolution in the Monocotyledons. , Denmark: Aarhus University Press. pp 267–286.
49. Drummond A, Ashton B, Buxton S, Cheung M, Cooper A, et al. (2010) Geneious v4.7.6 Available: <http://www.geneious.com>. Accessed 2011 April 26.
50. Rozen S, Skaletsky H (2000) Primer3 for general users and for biologist programmers. In: Krawetz S, Misener S, eds. Bioinformatics methods and protocols: Methods in molecular biology. Totowa: Humana Press.
51. Caddick LR, Rudall PJ, Wilkin P, Hedderson T, Chase M (2002) Phylogenetics of Dioscoreales based on combined analyses of morphological and molecular data. *Botanical Journal of the Linnean Society* 138: 123–144.
52. R Development Core Team (2010) R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing.
53. Coursey DG (1967) Social and cultural importance. Yams, an account of the nature, origins, cultivation and utilisation of the useful members of the Dioscoreaceae. London, UK: Longmans. pp 197–205.
54. Hamon P (1987) Structure, origine génétique des ignames cultivées du complexe *Dioscorea cayenensis-rotundata* et domestication des ignames en Afrique de l'Ouest. Thèse de doctorat, Université Paris XI, France.
55. Terauchi R, Chikaleke V, Thottappilly G, Hahn S (1992) Origin and phylogeny of Guinea yams as revealed by RFLP analysis of chloroplast DNA and nuclear ribosomal DNA. *Theoretical and Applied Genetics* 83: 743–751.
56. Scarcelli N, Tostain S, Vigouroux Y, Agbangla C, Dainou O, et al. (2006) Farmers' use of wild relative and sexual reproduction in a vegetatively propagated crop. The case of yam in Benin. *Molecular Ecology* 15: 2421–2431.
57. Excoffier L, Smouse PE (1994) Using allele frequencies and geographic subdivision to reconstruct gene trees within a species: Molecular variance parsimony. *Genetics* 136: 343–359.
58. Sarah G, Ruiz M, Perrier X, Billot C Available: <http://haplophyle.cirad.fr/index.jsp>. Accessed 2011 April 26.
59. Vicentini A, Barber JC, Alisiani SS, Giussani LM, Kellogg EA (2008) The age of the grasses and clusters of origins of C4 photosynthesis. *Global Change Biology* 14: 2963–2977.
60. Roncal J, Borchsenius F, Asmussen-Lange CB, Balslev H (2010) Divergence times in the tribe Geonomateae (Arecaceae) coincide with Tertiary geological events. In: Seberg O, Petersen G, Barfod A, Davis J, eds. Diversity, phylogeny, and evolution in the Monocotyledons: Aarhus University Press. pp 245–264.
61. Gaut BS, Morton BR, MacCaig BC, Clegg MT (1996) Substitution rate comparisons between grasses and palms: synonymous rate differences at the nuclear gene *Adh* parallel rate differences at the plastid gene *rbcL*. *Proceedings of the National Academy of Sciences of the United States of America* 93: 10274–10279.
62. Cozzolino S, Cafasso D, Pellegrino G, Musacchio A, Wildmer A (2003) Molecular evolution of a plastid tandem repeat locus in an orchid lineage. *Journal of Molecular Evolution* 57: S41–S49.
63. Asmussen CB, Dransfield J, Deickmann V, Barfod AS, Pintaud J-C, et al. (2006) A new subfamily classification of the palm family (Arecaceae): evidence from plastid DNA phylogeny. *Botanical Journal of the Linnean Society* 151: 15–38.
64. Couvreur TLP, Hahn WJ, de Granville J-J, Pham J-L, Ludena B, et al. (2007) Phylogenetic relationships of the cultivated Neotropical palm *Bactris gasipaes* (Arecaceae) with its wild relatives inferred from chloroplast and nuclear DNA polymorphisms. *Systematic Botany* 32: 519–530.
65. Eiserhardt W, Pintaud J, Asmussen-Lange C, Hahn W, Bernal R, et al. (In press) Phylogeny and divergence times of Bactridiinae (Arecaceae, Palmae) based on plastid and nuclear DNA sequences. *Taxon*.
66. Trénel P, Gustafsson MHG, Baker WJ, Asmussen-Lange CB, Dransfield J, et al. (2007) Mid-Tertiary dispersal, not Gondwanan vicariance explains distribution patterns in the wax palm subfamily (Ceroxylodeae: Arecaceae). *Molecular Phylogenetics and Evolution* 45: 272–288.
67. Pintaud J-C, Couvreur TLP, Lara C, Ludeña B, Pham J-L (2008) Reciprocal introgression between wild and cultivated peach palm (*Bactris gasipaes* Kunth, Arecaceae) in western Ecuador. In: Maxted N, Ford-Lloyd BV, Kell SP, Iriondo JM, Dulloo ME, et al. (2008) Crop wild relatives conservation and use. , UK: CAB International. pp 296–308.
68. Kelchner S (2000) The evolution of non-coding chloroplast DNA and its implication in plant systematics. *Annals of the Missouri Botanical Garden* 87: 482–498.

69. Savolainen V, Anstett M-C, Lexer C, Hutton I, Clarkson JJ, et al. (2006) Sympatric speciation in palms on an oceanic island. *Nature* 441: 210–213.
70. Birky CW, Jr. (1995) Uniparental inheritance of mitochondrial and chloroplast genes: Mechanisms and evolution. *Proceedings of the National Academy of Sciences of the United States of America* 92: 11331–11338.
71. Chair H, Perrier X, Agbangla C, Marchand JL, Dainou O, et al. (2005) Use of cpSSRs for the characterisation of yam phylogeny in Benin. *Genome* 48: 674–684.