

New Insights into the Population Structure of *Anopheles gambiae* s.s. in the Gulf of Guinea Islands Revealed by *Herves* Transposable Elements

Patrícia Salgueiro^{1*}, Marta Moreno^{2‡}, Frédéric Simard³, David O'Brochta⁴, João Pinto¹

1 Centro de Malaria e outras Doenças Tropicais/UEI Parasitologia Médica, Instituto de Higiene e Medicina Tropical, Universidade Nova de Lisboa, Lisboa, Portugal, **2** Centro Nacional de Medicina Tropical, Instituto de Salud Carlos III, Madrid, Spain, **3** MIVEGEC (Maladies Infectieuses et Vecteurs: Ecologie, Genétique, Evolution et Contrôle), UMR IRD224-CNRS5290-UM1-UM2, Institut de Recherche pour le Développement, Montpellier, France, **4** Department of Entomology and The Institute for Bioscience and Biotechnology Research, University of Maryland, College Park, Rockville, Maryland, United States of America

Abstract

Transposable elements (TEs) are mobile portions of DNA that are able to replicate and spread in the genome of many organisms. TEs can be used as a means to insert transgenes in insects, being stably inherited throughout generations. *Anopheles gambiae* is the main vector of human malaria in Sub-Saharan Africa. Given the extraordinary burden this disease imposes, the mosquito became a choice target for genetic control approaches with the purpose of reducing malaria transmission. In this study, we investigated the abundance and distribution of *Herves* TE in *An. gambiae* s.s. from Cameroon and four islands in the Gulf of Guinea, in order to determine their genetic structure. We have detected a population subdivision between Equatorial Guinea islands and the islands of Sao Tome, Principe and mainland. This partitioning associates more with political rather than geographic boundaries, possibly reflecting different mainland source populations colonizing the islands.

Citation: Salgueiro P, Moreno M, Simard F, O'Brochta D, Pinto J (2013) New Insights into the Population Structure of *Anopheles gambiae* s.s. in the Gulf of Guinea Islands Revealed by *Herves* Transposable Elements. PLoS ONE 8(4): e62964. doi:10.1371/journal.pone.0062964

Editor: Zhijian Tu, Virginia Tech, United States of America

Received: December 4, 2012; **Accepted:** March 27, 2013; **Published:** April 26, 2013

Copyright: © 2013 Salgueiro et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work received financial support from the UNICEF/UNDP/World Bank/WHO Special Program for Research and Training in Tropical Diseases (TDR, A50239) and CMDT/IHMT Plurianual Program funds, as well as the US National Institutes of Health grant GM48102. PS was funded by a post-doc fellowship (SFRH/BPD/72532/2010) from Fundação para a Ciência e Tecnologia (FCT) through the European Program FEDER. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: Senior author Joao Pinto is part of the academic editorial board of PLOS ONE. There are no other competing interests to declare.

* E-mail: psalgueiro@ihmt.unl.pt

Current address: Division of Infectious Diseases, School of Medicine, University of California San Diego, La Jolla, California, United States of America

Introduction

Over the past recent years, transgenic technologies have been proposed for malaria vector control [1,2,3]. One of the approaches in development consists in altering the vectorial capacity of wild populations of *Anopheles* mosquitoes by delivering and spreading transmission-blocking transgenes (e.g. refractoriness to *Plasmodium* infection) [4,5]. This approach requires gene drive systems and transposable elements (TEs) were one of the first candidate tools to be considered for population replacement [6,7].

Transposons or TEs are mobile genetic sequences that can replicate and change their relative position within the genome [8]. Most TEs used for insect germline transformation are Class II elements that transpose by a direct DNA cut-and-paste mechanism [7,9,10,11]. Within the genome of the main Afrotropical malaria vector *Anopheles gambiae sensu stricto* several TEs have been catalogued [12,13]. Among these, the *Herves* element is a transpositionally active Class II TE isolated from *An. gambiae* s.s. and found in the sibling species *Anopheles arabiensis* and *Anopheles merus* [14]. *Herves* was inferred to be transpositionally active in natural populations of *An. gambiae* based on the high frequency of intact, transposase-encoding *Herves* elements observed and highly polymorphic insertion sites [15].

The dynamics of TEs within populations is determined by the transpositional activity of the element, the strength of self- and host-regulatory mechanisms and the genetic structure of host populations [16]. Sequence and insertion-site polymorphisms have allowed TEs to be used as markers for population genetic studies [17], including in *An. gambiae* [18,19,20].

Islands have been considered as candidate sites for experimental releases of transgenic mosquitoes for vector control [21,22,23], because the effects of migration from neighboring regions not under control are expected to be minimal. In this context, the islands of the Gulf of Guinea (West Africa) are potential sites that might benefit from malaria control strategies based on transgenic vectors. These islands are volcanic in origin and extend from the coast of Cameroon (Bioko island, 32 km off shore) to 350 km west of the coast of Gabon (Annobon island, Figure 1). Both Bioko and Annobon are part of Equatorial Guinea. Between these two islands, the archipelago of Sao Tome and Principe is found, which comprises two islands located 140 km apart and about 250 and 225 km, respectively, off the northwestern coast of Gabon. The main malaria vector on these islands is *An. gambiae* s.s. [24,25]. In the island of Bioko, as well as in mainland (Equatorial Guinea and Cameroon), both M and S molecular forms, considered as incipient species within *An. gambiae* s.s. [26], occur in sympatry




Figure 1. Map of the Gulf of Guinea displaying the islands where collections were conducted and the geographic linear distances among them.

doi:10.1371/journal.pone.0062964.g001

[19,25]. In São Tomé, Príncipe and Annobón only the M form is present [24,27].

Populations of *An. gambiae* s.s. on Bioko island were found to be genetically differentiated from mainland populations, using microsatellite DNA markers [28]. However, Moreno et al. [27] found a higher degree of genetic isolation of Annobón island when compared with Bioko, which is closest to mainland. Similarly, it has been shown that populations of *An. gambiae* s.s. from the islands of São Tomé and Príncipe are genetically isolated from continental populations [29,30]. The later authors suggested two main colonization events for *An. gambiae* s.s. into these islands, coincident with episodes of intense human migration [30].

In the islands of the Gulf of Guinea, malaria remains a major cause of morbidity and child mortality. However, in recent years the situation has progressively changed. In São Tomé and Príncipe, a malaria control program has reduced the number of cases by half over the past decade [31]. This program included the use of Indoor Residual Spraying (IRS), Insecticide Treated Nets (ITNs) and artemisinin-combination therapies [32,33]. Similarly, the prevalence of childhood infection in the island of Bioko has been dropping considerably since the combined implementation of ITNs and IRS [34,35]. Under these circumstances, the implementation of a transgenic control component might be a promising complement to current malaria control measures, especially when insecticide-based measures are known to be difficult to sustain on the long-term due to issues such as financial constraints or insecticide resistance. However, there are still many issues related with the applicability of transgenic technologies to vector control that require further attention. One of these issues relates with the accurate estimation of the degree of isolation of island populations from mainland [30,36]. In addition, understanding the dynamics of TE elements in insular populations of *An.*

gambiae is critical for developing predictive tools to be used in planning and managing release of transgenic mosquitoes.

In this study, we measured the abundance and site-occupancy frequency distribution of the *Herves* transposon in *An. gambiae* s.s. from Cameroon and the four islands in the Gulf of Guinea, in order to determine the genetic structure of this TE among physically isolated populations and relate with the potential applicability of transposons as gene drive systems for island vector populations.

Materials and Methods

Mosquito DNA samples

Mosquito specimens were collected from one continental population of Cameroon, and four island populations of the Gulf of Guinea: Bioko, Príncipe, São Tomé and Annobón (Table 1, Figure 1). Details on the localities and sampling methods have been described elsewhere [27,29]. Table 1 recalls collection dates and GPS position of collection sites, and Figure 1 displays the geographic linear distances between collection sites.

No specific permits were required for the described field studies. Mosquito collections were performed in villages either inside houses or outdoors. Informed verbal consent was obtained from household owners to perform the indoor sampling. This type of consent was chosen to avoid potential conflicts due to mistrust in signing official forms by the community. No specific ethics clearance was requested, since all mosquito collections were performed as part of routine mosquito surveillance implemented by the Malaria Control Programs of the regions sampled. The field studies did not involve endangered or protected species.

Extraction of DNA from single mosquitoes was performed by a phenol-chloroform protocol [37]. Species identification and determination of molecular forms were performed by PCR-RFLP

Table 1. Collection data, site occupancy and genetic diversity of the sampled *Anopheles gambiae* populations.

Country	island/ mainland	Locality	Geographical coordinates	Year of collection	N	k	dun	hs	95% credibility interval of hs	
									lower	upper
S. Tomé and Príncipe	S. Tome	Riboque	0°19'N/6°43'E	1997	14	15	7.1	0.299	0.238	0.356
				2004	12	15	5.6	0.315	0.241	0.376
	Príncipe	Rua dos Trabalhadores	1°38'N/7°25'E	1998	13	12	4.5	0.302	0.234	0.363
Equatorial Guinea	Annobon	Annobon	1°24'S/5°57'E	2004	14	21	7.5	0.291	0.232	0.344
	Bioko	Malabo	3°45'N/8°46'E	2003	14	12	6.0	0.301	0.238	0.358
Cameroon	mainland	Tiko	4°05'N/9°21'E	1999	14	15	4.9	0.273	0.210	0.331
					$\Sigma = 81$			Hs = 0.297	0.243	0.340

N- number of individuals analyzed by TE display per locality; k Number of unique chromosomal sites containing *Herves*; dcn Diploid copy number of *Herves* [39]; hs gene diversity with credibility intervals calculated by a Bayesian approach as implemented in HICKORY; Hs the mean within-population expected heterozygosity (= Neis gene diversity within populations).
doi:10.1371/journal.pone.0062964.t001

according to a previously described protocol [38]. A total of 81 *An. gambiae* s.s. of the M molecular form were analyzed in this study. Sample sizes for each locality are shown in Table 1.

Transposable element display

PCR-based transposable element display was performed as described by Subramanian et al. [15]. Images of the scanned TE display gels were printed and bands were assigned to a molecular weight based on their mobility relative to the size markers using a standard curve developed for each gel. All samples from each site were analyzed on the same gel, allowing a single standard molecular weight curve to be used to score each dataset. On the basis of the combined results of three TE display experiments, a band was called as present or absent if it was unambiguously present in at least two of the three replicates. Determining the presence of bands in this way resulted in a single scoring matrix that was then used in subsequent analyses.

Data Analysis

Site occupancy frequency distributions were estimated using the TE display data. To obtain estimates of element frequencies and copy numbers per haploid genome we assumed HardyWeinberg equilibrium and followed Wright et al. [39], where the mean number of elements per haploid genome was calculated as described in [40]. In order to discard any locus under selection, we looked for outlier loci using the Dfdist approach with 100,000 simulations [41,42] in MCHEZA program [43].

Expected heterozygosity *hs*, mean within-population expected heterozygosity *Hs* and their 95% credibility values, were calculated using a Bayesian approach with 250,000 generations after a burn-in of 50,000 generations implemented in HICKORY 1.1 [44].

We calculated pairwise *Fst* estimates as a measure of genetic differentiation based on [45] and tested isolation by distance through the analysis of the correlation coefficient of *Fst*/(1-*Fst*) over pairwise geographical distances using Mantel tests [46]. Analysis of Molecular Variance (AMOVA) was used to investigate the genetic structure of the mosquito populations as described in Excoffier et al. [47]. For all these analyses, we used ARLEQUIN 3.1.1 [48]. Permutation-based statistical tests were performed with 50,000 permutations.

For a visual representation of genetic differentiation patterns, we performed a factorial correspondence analysis (FCA) on the

multilocus genotype of each individual, using the option that takes into account the population of origin (FCA over populations), as implemented in GENETIX v. 4.05.2 [49].

Results

Site occupancy

We detected *Herves* elements in all specimens analyzed by TE display (N= 81). A total of 35 sites were detected. Within each sample, element copy number (dun) ranged from 4.5 to 7.5 per diploid genome (Table 1). The maximum number of elements per individual ranged from six in Príncipe to nine in Annobon.

The TEs with the highest site-occupancy frequency were: *Herves* element with 160 bp (Hv160, showing 613 occurrences per sample) and Hv130 (614 occurrences per sample) (Table S1, supporting information file). We have found a single element (Hv157) common to all six samples. The sample with the highest number of exclusive sites was Annobon (7, Table 2). The two temporal samples from S. Tomé shared 13 out of 15 possible sites, and these had 10 sites in common with the continental sample (Cameroon, Table 2).

Table 2. Distribution of shared *Herves* sites among populations of *An. gambiae* in the Gulf of Guinea.

	Came roon	Bio ko	Prín cipe	S. Tom é 97	S. Tom é04	Anno bón
Cameroon	<u>2</u>					
Bioko	5	<u>1</u>				
Príncipe	6	4	<u>3</u>			
S. Tome 1997	10	5	8	<u>1</u>		
S. Tome 2004	10	5	9	13	<u>0</u>	
Annobon	7	7	7	9	9	<u>7</u>

Values in the diagonal represent the number of sites exclusive to the correspondent population.

doi:10.1371/journal.pone.0062964.t002

Population structure of Herves element

The analysis with Dfdist showed no evidence of sites under




Figure 2. Projection of 81 individual *Herves* genotypes of *Anopheles gambiae* on the principal axes of a factorial component analysis. Each colour corresponds to a sampled island population as indicated in the legend. Inertia percentage values are presented for each factorial component (FC1 and FC2).
doi:10.1371/journal.pone.0062964.g002

populations. In this context, the analysis of additional mainland samples would be required in order to further clarify the source populations of *An. gambiae* s.s. from Annobon and Bioko.

Finally, we cannot exclude the possibility that differences in the mode of evolution and inheritance may also contribute for the apparent discrepancy between patterns of population structure obtained by the *Herves* TE and by previous microsatellite based analyses. However, little is known about the mechanisms of regulation of *Herves* TE activity, which poses difficulties for comparing these markers.

The low differentiation observed between the two temporal samples from S. Tome provides no evidence for significant demographic changes in the mosquito population between 1997 and 2004 or for high rates of *Herves* transposition during this period. This result is consistent with the large estimates of current effective population size obtained for *An. gambiae* M-form in S

Tomeisland [29,50], comparable to those observed in mainland for the members of *An. gambiae* complex [55,56]. The relative demographic stability of *An. gambiae* suggested by this and previous studies in S

and Principe is based on the analysis of samples for which collections pre-date the recent implementation of vector control measures that resulted in a significant malaria decrease on the islands [32,33]. In this context, it would be interesting to analyze the genetic variation of *Herves* TE in samples collected post-intervention as a means to assess the usefulness of TE in determining the impact of vector control in the genetic structure of *An. gambiae*.

In this study, locally fixed *Herves*-occupied sites were rare and sites occupied in only a few individuals were very common in the sampled populations. This suggests that *Herves* remains active within these island mosquito populations, as was also observed in previous analyses of other mainland African *An. gambiae* popula-

Table 4. Apportionment of molecular variance measured among populations of *An. Gambiae*.

Groups Tested	Total variance	% among groups	P	% among populations within groups	% within populations
A	0.30	56.3	<0.001	-2.7	46.5
B	0.21	-11.8	n.s.	44.5	67.4
C	0.26	48.3	<0.05	-3.1	54.8
D	0.23	41.8	n.s.	-2.74	60.9

P represents the significance of the variation among groups (random > observed). Variation was partitioned in the tested structures.

A: group 1: Cameroon, Principe, S. Tome (1997, 2004), group 2: Annobon and Bioko;

B: group 1: Cameroon, group 2: all other populations;

C: group 1: Cameroon, group 2: S. Tome and Principe, group 3: Annobon and Bioko.

D: group 1: Cameroon, group 2: Principe, group3: S. Tome, group 4: Annobon and group 5: Bioko.

doi:10.1371/journal.pone.0062964.t004

tions [15,40]. Following WHO's recommendations [22], transgenic mosquitoes use should involve several phases of field release under conditions that limit spread into the environment. In this context, remote islands are considered as suitable candidate sites. Accordingly, the most recent open-field trials to attempt transgenic-based vector control involved island mosquito populations [23]. In agreement with previous studies based on other genetic markers [27], the analysis of *Herves* TE suggests Annobon as probably the most adequate of the four islands of the Gulf of Guinea for experimental release of transgenic vectors: it is the most remote island with *An. gambiae* displaying the highest degree of genetic isolation and the highest amount of *Herves* insertion site polymorphism. However, the patterns of population structure on the islands of the Guinean Gulf disclosed in this study by *Herves* TE are more likely to reflect different origins of mosquito colonization rather than levels of contemporary gene flow. These results should therefore be interpreted with caution. Further analyses involving comparisons with more mainland samples and combining different genetic markers are required to clarify the degree of isolation of these islands.

References

- Collins F, Sakai R, Vernick K, Paskewitz S, Seeley D, et al. (1986) Genetic selection of a Plasmodium-refractory strain of the malaria vector *Anopheles gambiae*. *Science* 234: 607610.
- Miller LH, Sakai RK, Romans P, Gwadz RW, Kantoff P, et al. (1987) Stable integration and expression of a bacterial gene in the mosquito *Anopheles gambiae*. *Science* 237: 779781.
- Marshall JM, Taylor CE (2009) Malaria Control with Transgenic Mosquitoes. *PLoS Med* 6: e20.
- Ito J, Ghosh A, Moreira LA, Wimmer EA, Jacobs-Lorena M (2002) Transgenic anopheline mosquitoes impaired in transmission of a malaria parasite. *Nature* 417: 452455.
- Kim W, Koo H, Richman AM, Seeley D, Vizioli J, et al. (2004) Ectopic expression of a Cecropin transgene in the human malaria vector mosquito *Anopheles gambiae* (Diptera: Culicidae): effects on susceptibility to Plasmodium. *J Med Entomol* 41: 447455.
- Ribeiro JMC, Kidwell MG (1994) Transposable Elements as population drive mechanisms: specification of critical parameter values. *J Med Entomol* 31: 1016.
- Sinkins SP, Gould F (2006) Gene drive systems for insect disease vectors. *Nat Rev Genet* 7: 427435.
- Craig NL, Craigie R, Gellert M, Lambowitz A, editors (2002) Mobile DNA II. Washington D.C.: American Society for Microbiology Press. 1204 p.
- Finnegan DJ (1989) Eukaryotic transposable elements and genome evolution. *Trends Genet* 5: 103107.
- Fraser MJ (2012) Insect Transgenesis: Current Applications and Future Prospects. *Annu Rev Entomol* 57: 267289.
- Handler AM, OBrochta DA (2012) Transposable elements for insect transformation. In: Gilbert LI, editor. *Insect Biochemistry and Molecular Biology*. London, United Kingdom: Academic Press. 90133.
- Struchiner CJ, Massad E, Tu Z, Ribeiro JMC (2009) The tempo and mode of evolution of transposable elements as revealed by molecular phylogenies reconstructed from mosquito genomes. *Evolution* 63: 31363146.
- Fernández-Medina RD, Struchiner CJ, Ribeiro JMC (2011) Novel transposable elements from *Anopheles gambiae*. *BMC Genomics* 12: 260260.
- Arensburger P, Kim Y-J, Orsetti J, Aluvihare C, OBrochta DA, et al. (2005) An active transposable element, Herves, from the African malaria mosquito *Anopheles gambiae*. *Genetics* 169: 697708.
- Subramanian RA, Arensburger P, Atkinson PW, OBrochta DA (2007) Transposable element dynamics of the hAT element Herves in the human malaria vector *Anopheles gambiae* s.s. *Genetics* 176: 24772487.
- Deceliere G, Charles S, Biemont C (2005) The dynamics of transposable elements in structured populations. *Genetics* 169: 467474.
- Lepetit D, Brehm A, Fouillet P, Biemont C (2002) Insertion polymorphism of retrotransposable elements in populations of the insular, endemic species *Drosophila madeirensis*. *Mol Ecol* 11: 347354.
- Barnes MJ, Lobo NF, Coulibaly MB, Sagnon NF, Costantini C, et al. (2005) SINE insertion polymorphism on the X chromosome differentiates *Anopheles gambiae* molecular forms. *Insect Mol Biol* 14: 353363.
- Boulesteix M, Simard F, Antonio-Nkondjio C, wono-Ambene H, Fontenille D, et al. (2007) Insertion polymorphism of transposable elements and population structure of *Anopheles gambiae* M and S molecular forms in Cameroon. *Mol Ecol* 16: 441452.
- Esnault C, Boulesteix M, Duchemin JB, Koffi AA, Chandre F, et al. (2008) High genetic differentiation between the M and S molecular forms of *Anopheles gambiae* in Africa. *PLoS ONE* 3: e1968.
- Alphey L, Beard CB, Billingsley P, Coetzee M, Crisanti A, et al. (2002) Malaria control with genetically manipulated insect vectors. *Science* 298: 119121.
- WHO (2009) Progress and prospects for the use of genetically modified mosquitoes to inhibit disease transmission. Geneva: WHO. 68 p.
- Harris AF, Nimmo D, McKemey AR, Kelly N, Scaife S, et al. (2012) Field performance of engineered male mosquitoes. *Nat Biotechnol* 29: 10341037.
- Pinto J, Sousa CA, Gil V, Ferreira C, Goncalves L, et al. (2000) Malaria in São Tomé and Príncipe: parasite prevalences and vector densities. *Acta Trop* 76: 185193.
- Berzosa P, Cano J, Roche J, Rubio J, García L, et al. (2002) Malaria vectors in Bioko Island (Equatorial Guinea): PCR determination of the members of *Anopheles gambiae* Giles complex (Diptera: Culicidae) and pyrethroid knockdown resistance (kdr) in *An. gambiae sensu stricto*. *J Vector Ecol* 27: 102106.
- della Torre A, Fanello C, Akogbeto M, Dossou-yovo J, Favia G, et al. (2001) Molecular evidence of incipient speciation within *Anopheles gambiae* s.s. in West Africa. *Insect Mol Biol* 10: 918.
- Moreno M, Salgueiro P, Vicente J, Cano J, Berzosa P, et al. (2007) Genetic population structure of *Anopheles gambiae* in Equatorial Guinea. *Malaria J* 6: 137.
- Reimer LJ, Tripet F, Slotman M, Spielman A, Fondjo E, et al. (2005) An unusual distribution of the kdr gene among populations of *Anopheles gambiae* on the island of Bioko, Equatorial Guinea. *Insect Mol Biol* 14: 683688.
- Pinto J, Donnelly MJ, Sousa CA, Malta-Vacas J, Gil V, et al. (2003) An island within an island: genetic differentiation of *Anopheles gambiae* in São Tomé, West Africa, and its relevance to malaria vector control. *Heredity* 91: 407414.
- Marshall JC, Pinto J, Charwood JD, Gentile G, Santolamazza F, et al. (2008) Exploring the origin and degree of genetic isolation of *Anopheles gambiae* from the islands of São Tomé and Príncipe, potential sites for testing transgenic-based vector control. *Evol Appl* 1: 631644.
- WHO (2011) World Malaria Report.
- Lee P-W, Liu C-T, Rampao HS, do Rosario V, Shaio M-F (2010) Pre-elimination of malaria on the island of Príncipe. *Malaria J* 9: 26.
- Teklehaimanot HD, Teklehaimanot A, Kiszewski A, Rampao HS, Sachs JD (2009) Malaria in São Tomé and Príncipe: on the brink of elimination after three years of effective antimalarial measures. *Am J Trop Med Hyg* 80: 133140.
- Kleinschmidt I, Sharp B, Benavente LE, Schwabe C, Torrez M, et al. (2006) Reduction in infection with *Plasmodium falciparum* one year after the introduction of malaria control interventions on Bioko Island, Equatorial Guinea. *Am J Trop Med Hyg* 74: 972978.
- Pardo G, Descalzo MA, Molina L, Custodio E, Lwanga M, et al. (2006) Impact of different strategies to control Plasmodium infection and anaemia on the island of Bioko (Equatorial Guinea). *Malaria J* 5: 10.
- Lanzaro G, Nuzhdin S, Tripet F (2006) Tools for monitoring the genetic structure and stability of mosquito populations. Bridging laboratory and field research for genetic control of disease vectors. 157164.
- Donnelly MJ, Cuamba N, Charwood JD, Collins FH, Townson H (1999) Population structure in the malaria vector, *Anopheles arabiensis* Patton, in East Africa. *Heredity* 83: 408417.
- Favia G, Lanfrancotti A, Spanos L, Siden-Kiamos I, Louis C (2001) Molecular characterization of ribosomal DNA polymorphisms discriminating among chromosomal forms of *Anopheles gambiae* s.s. *Insect Mol Biol* 10: 1923.

Supporting Information

Table S1 Distribution of Herves sites detected over the six sampled populations.

(DOCX)

Acknowledgments

We thank Barbara Ngudiankama and Kristina Pilitt (University of Maryland, USA) for valuable technical advice. We also acknowledge Dr. Conceição Ferreira (Centro Nacional de Endemias) for logistic support to mosquito collections carried out in São Tomé and Príncipe.

Author Contributions

Conceived and designed the experiments: DOB JP PS. Performed the experiments: PS. Analyzed the data: PS DOB. Contributed reagents/materials/analysis tools: DOB FS JP MM. Wrote the paper: PS JP.

39. Wright SI, Le QH, Schoen DJ, Bureau TE (2001) Population dynamics of an ac-like transposable element in self- and cross-pollinating *Arabidopsis*. *Genetics* 158: 12791288.
40. OBrochta DA, Subramanian R, Orsetti J, Peckham E, Nolan N, et al. (2006) *hAT* element population genetics in *Anopheles gambiae s.l.* in Mozambique. *Genetica* 127: 185198.
41. Beaumont MA, Balding DJ (2004) Identifying adaptive genetic divergence among populations from genome scans. *Mol Ecol* 13: 969980.
42. Beaumont MA, Nichols RA (1996) Evaluating Loci for Use in the Genetic Analysis of Population Structure. *Proc R Soc B-Biol Sci* 263: 16191626.
43. Antao T, Beaumont MA (2011) Mchaza: A Workbench to Detect Selection Using Dominant Markers. *Bioinformatics*.
44. Holsinger KE, Lewis PO (2005) Hickory: A Package for Analysis of Population Genetic Data v1. Storrs: Department of Ecology and Evolutionary Biology, University of Connecticut.
45. Weir B, Cockerham C (1984) Estimating F-statistics for the analysis of population structure. *Evolution* 38: 13581370.
46. Rousset F (1997) Genetic differentiation and estimation of gene flow from F-statistics under isolation by distance. *Genetics* 145: 12191228.
47. Excoffier L, Smouse P, Quattro J (1992) Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics* 131: 479491.
48. Excoffier L, Laval G, Schneider S (2005) Arlequin ver. 3.0: An integrated software package for population genetics data analysis. *Evol Bioinform Online* 1: 4750.
49. Belkhir K, Borsa P, Chikhi L, Raufaste N, Bonhomme F (1996) GENETIX 4.05, computer program, logiciel sous Windows TM pour la génétique des populations.
50. Pinto J, Donnelly MJ, Sousa CA, Gil V, Ferreira C, et al. (2002) Genetic structure of *Anopheles gambiae* (Diptera: Culicidae) in São Tome and Príncipe (West Africa): implications for malaria control. *Mol Ecol* 11: 21832187.
51. Ribeiro H, Ramos HC, Pires CA (1990) Sobre os vectores da malaria em São Tomee Príncipe. *Garcia de Orta Serie Zoologia* 15: 135152.
52. Molina R, Benito A, Roche J, Blanca F, Amela C, et al. (1993) Baseline entomological data for a pilot malaria control program in Equatorial Guinea. *J Med Entomol* 30: 622624.
53. Jones PJ (1994) Biodiversity in the Gulf of Guinea: an overview. *Biodiversity & Conservation* 3: 772784.
54. Mateu E, Comas D, Calafell F, Pérez-Lezaun A, Abade A, et al. (1997) A tale of two islands: population history and mitochondrial DNA sequence variation of Bioko and São Tome, Gulf of Guinea. *Annals of Human Genetics* 61: 507518.
55. Lehmann T, Hawley WA, Grebert H, Collins FH (1998) The effective population size of *Anopheles gambiae* in Kenya: implications for population structure. *Mol Biol Evol* 15: 264276.
56. Simard F, Lehmann T, Lemasson J, Diatta M, Fontenille D (2000) Persistence of *Anopheles arabiensis* during the severe dry season conditions in Senegal: an indirect approach using microsatellite loci. *Insect Mol Biol* 9: 467479.