Conserving Biodiversity in a Human-Dominated World: Degradation of Marine Sessile Communities within a Protected Area with Conflicting Human Uses

Valeriano Parravicini^{1,2}*, Fiorenza Micheli³, Monica Montefalcone⁴, Carla Morri⁴, Elisa Villa⁴, Michela Castellano⁴, Paolo Povero⁴, Carlo Nike Bianchi⁴

1 CESAB-FRB, Immeuble Henri Poincaré - Domaine du Petit Arbois, Aix-en-Provence, France, 2 IRD – UR 227 Coreus – Laboratoire Arago, Banyuls/mer, France, 3 Hopkins Marine Station, Stanford University, Pacific Grove, California, United States of America, 4 DiSTAV, Dipartimento di Scienze della Terra, dell'Ambiente e della Vita, Genoa University, Genoa, Italy

Abstract

Conservation research aims at understanding whether present protection schemes are adequate for the maintenance of ecosystems structure and function across time. We evaluated long-term variation in rocky reef communities by comparing sites surveyed in 1993 and again in 2008. This research took place in Tigullio Gulf, an emblematic case study where various conservation measures, including a marine protected area, have been implemented to manage multiple human uses. Contrary to our prediction that protection should have favored ecosystem stability, we found that communities subjected to conservation measures (especially within the marine protected area) exhibited the greatest variation toward architectural complexity loss. Between 1993 and 2008, chronic anthropogenic pressures (especially organic load) that had already altered unprotected sites in 1993 expanded their influence into protected areas. This expansion of human pressure likely explains our observed changes in the benthic communities. Our results suggest that adaptive ecosystem-based management (EBM), that is management taking into account human interactions, informed by continuous monitoring, is needed in order to attempt reversing the current trend towards less architecturally complex communities. Protected areas are not sufficient to stop ecosystem alteration by pressures coming from outside. Monitoring, and consequent management actions, should therefore extend to cover the relevant scales of those pressures.

Citation: Parravicini V, Micheli F, Montefalcone M, Morri C, Villa E, et al. (2013) Conserving Biodiversity in a Human-Dominated World: Degradation of Marine Sessile Communities within a Protected Area with Conflicting Human Uses. PLoS ONE 8(10): e75767. doi:10.1371/journal.pone.0075767

Editor: Arga Chandrashekar Anil, CSIR- National institute of oceanography, India

Received April 3, 2013; Accepted August 18, 2013; Published October 15, 2013

Copyright: © 2013 Parravicini et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The present work is done under the aegis of the project 'The impacts of biological invasions and climate change on the biodiversity of the Mediterranean Sea' (Italy–Israel co-operation) funded by the Italian Ministry of the Environment. FM was partly supported by a Pew Fellowship in Marine Conservation. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: valeriano.parravicini@ird.fr

Introduction

Most marine ecosystems are challenged worldwide by a vast set of human pressures ranging from pollution to habitat modification, overfishing and climate change [1]. The recognition of the profound influence humans exert on marine ecosystems has motivated worldwide conservation efforts mainly aimed at the establishment of Marine Protected Areas (MPAs). Yet, MPAs alone are not sufficient [2,3]. They rarely cover an adequate extent of ecosystem types being typically quite small (ranging approx. 10^{-3} to 10^3 km²) and are thereby ineffective in halting the effects of pressures acting at scales larger than those encompassed by the protection schemes.

Anthropogenic pressures acting over large scales are common [1], especially along intensely populated coastal areas, where marine ecosystems are challenged by the cumulative effect of multiple pressures and conflicting human uses. The need for complementary measures to the protection by MPAs has called for the development of ecosystem-based management (EBM) to be implemented at regional scales. EBM is an integrated approach that considers the entire ecosystem, including humans [4]. EBM differs from other approaches that focus on a single species or

sector because it includes consideration of the interactions among ecosystem components and the cumulative impacts of multiple activities [5]. Some examples of EBM implementation are the zoning of the Great Barrier Reef Marine Park [6] and the establishment of marine sanctuaries in the USA [7]. Similar instruments have been developed in the European Union, where legislation mandates the establishment of a network of *Sites of Community Importance* (SCIs) to preserve habitats and species [8] and *Integrated Coastal Zone Management* (ICZM), whose goal is the sustainable development of coastal areas [9]. Despite the application of EBM approaches to complement MPAs, few studies have assessed whether such plans effectively sustain the conservation of marine coastal ecosystems. Ecosystem-based approaches have recently been advocated also to understand ecosystem shifts due to climate change [10].

Thanks to historical data from 1993 onwards ([11] and unpublished information), we were able to evaluate the results of the management plan applied in an emblematic study case (the Tigullio Gulf) by examining the decadal-scale change of rocky reef communities. The Tigullio Gulf is a densely populated area where conservation measures included the establishment of one MPA in 1998 (designed mainly to regulate fishing) and of six SCIs in 2000 (designed to regulate multiple human pressures on protected habitat and species). The area has been, and still is, subjected to various human uses and associated possible impacts: organic load from sewage outfalls, input of sediments and nutrients from a river, aquaculture, periodic beach replenishments, and small marinas.

Coastal development and related pressures have been shown to affect Mediterranean rocky reef communities by causing the replacement of canopy-forming algae with algal turfs, with consequent habitat homogenization and loss of architectural complexity [12]. In addition to removing or preventing these disturbances directly, through a regulation of development within SCIs, MPAs are expected to maintain or recover algal canopies through top-down control as protection from fishing allows predatory fishes to exert control upon sea urchins, which otherwise may proliferate and overgraze benthic algae [13,14]. Yet, at the same time, protection in MPAs may increase the abundance of herbivorous fish, which may also imply increased grazing pressure on canopy-forming algae [15].

To date, few studies have addressed how concomitant human pressures [16] impact reef communities within the Mediterranean Sea and whether EMB effectively manages non-fishing human uses that are not reduced by MPAs.

Here, we compared surveys of reef communities conducted prior to protection in 1993 and in 2008, 10 years after MPA and 8 years after SCIs establishment. We assessed relevant pressures operating in the study area and we related their intensities and trends to the observed change in reef communities in order to clarify whether the existing protection effort was effective. Results represent the first assessment of the efficacy of current management plans, and provide guidelines for EBM, with particular attention to the design of monitoring plans.

Methods

Study area and pressure regime

The Tigullio Gulf is located in the Ligurian Sea, NW Mediterranean Sea (Figure 1). The coastal zone is partially included within the Portofino MPA and six SCIs [3]. According to the standards of the European Water Framework Directive, water quality in the study area is classified as 'good' [17]. However, the area is densely populated by residents and tourists, resulting in comparatively high nutrient inputs into coastal waters.

Since the late 1950s, many studies have been carried out in this area [18]. Historical information on planktonic production [19,20] evidenced that chlorophyll-*a* concentration was significantly lower in the period 2000–2008 than in 1985–1994 (Figure S1a; t-test, df = 8, P<0.017). On the contrary, water transparency did not show any significant variation of the annual mean for the same periods (Figure S1b).

Potential anthropogenic pressures on coastal marine ecosystems in the last two decades include the presence of four outfalls discharging above the summer thermocline (i.e. about 25 m depth), the Entella River mouth (with a number of waste water pipes along its way), a fish farm (established in 2000), marinas and tourist harbors (built during the 1970s), and beaches frequently replenished (Table S1). Human population remained virtually stable: residents passed from 52 inhabitants km⁻² in 1993 to 50 inhabitants km⁻² in 2008, tourists from 1.34 million in 1998 to 1.36 million in 2007 (ISTAT, the Italian Statistical Institute). Due to change in local industrial activity, the potential organic load computed in term of equivalent individuals [21] showed a slight increase from 1991 to 2001 (Figure S1c). In the same decades, global warming has been showing a large influence on the hydrology and ecology of the Mediterranean basin [22]; the Ligurian Sea, in particular, underwent a regime shift in the late 1990s, two major heat waves in 1999 and 2003 having caused mass mortalities of gorgonians, sponges and other sessile organisms [23–25].

Sampling

Reef communities were surveyed in summer 1993 [11], before the establishment of the MPA and SCIs, in 10 sites showing similar environmental characteristics (Figure 1). Four sites (G, H, I and J) have subsequently been included within the Portofino MPA in 1998, the remaining six sites within SCIs since 2000. When necessary, all permits needed for performing fieldwork were obtained by the Genoa University from the Portofino MPA Authority. At each site, the percent cover of individual epibenthic taxa was visually estimated within four 1 m² quadrats randomly placed on vertical rocky walls at 5 m depth. Although the sampling effort employed may be considered rather small, Mediterranean rocky reef communities are 'miniaturized' compared to those of other temperate regions, and the surface of epibenthic communities surveyed at each site is one order of magnitude larger than the recommended minimum area of $2,170 \text{ cm}^2$ to be sampled in the Mediterranean [26]. Thus, our sampling units encompassed small-scale heterogeneity, while the four replicates were placed within a habitat (shallow vertical and sub-vertical walls) that exhibits fairly homogeneous assemblages over scales of 10 s m.

Percent cover was quantified by dividing the sampling quadrat into 25 sub-squares through a nylon line and giving each taxon a score from 0 to 4 within each square and then adding up scores for all squares within the quadrat [27]. Species identification was conducted visually at the lower possible taxonomic level; voucher specimens were collected only when in doubt and never involved protected or endangered species. In the same quadrats used for the assessment of epibenthic communities, the total density of seaurchins was estimated. In 2008, sites were precisely relocated by two of the authors (CNB and CM) that sampled communities in 1993 [11]. The sites were then surveyed by relocating 4 sampling quadrats at 5 m depth and following the same procedure. Vertical rocky walls were about 10 m height and the tidal height in the area is about 30 cm.

Community descriptors

Architectural aspects were considered in addition to taxonomic composition. For architectural complexity [28], the three-dimensionality of the assemblages was assessed. Based on the ratio between height (h) and radius (r), species were grouped into five categories: 'high' (h>r, e.g. *Cystoseira zosteroides*), 'medium-high' (h>r, e.g. *Dictyopteris polypodioides*), 'medium' (h~r, e.g. *Reteporella grimaldii*), 'medium-low' (h<r, e.g. *Petrosia ficiformis*), and 'low' (h «r, e.g. *Lithophyllum incrustans*).

Data analysis

In order to characterize the degree of human pressure upon individual sites, the distances from the MPA (protection) and the main potential sources of impact (i.e. river mouth, outfalls, fish farm, marinas, and replenished beaches) were calculated for each site. As raw explanatory variables were collinear, the matrix of distances from potential impact sources was treated with principal component analysis (PCA) [29]: the 1st and the 2nd axes of the PCA (PCA1 and PCA2) were found to be a good synthesis of the pressure gradient from protection by the MPA and closeness to potential sources of organic load (i.e. outfalls, river mouth, and fish farm), whereas the 3rd axis was correlated with the distance from the marinas and the replenished beaches (Table S2).



Figure 1. Position of sampling sites and of the main human-induced pressures in the study area. On the bottom, the position of individual sites on PCA axes is shown. The first axis represents closeness to sources of organic loading, the third closeness to replenished beaches and marinas.

doi:10.1371/journal.pone.0075767.g001

Two distinct datasets were obtained from communities surveys, the first using species as variables, the second using categories of three-dimensional structure. In order to test whether a significant change occurred in the time considered, permutational multivariate analysis of variance (PERMANOVA) [30] was used on each dataset considering sampling year and sampling site as random, crossed factors. PCA1 and PCA2 were included in the analysis as covariates and their interaction with sampling year were considered in order to test whether different pressures lead to different variation through time. The two sampling years (i.e. 1993 vs. 2008) were considered as random. Protection in the MPA was not included as a categorical factor to avoid redundancy since its potential effect was already considered in the PCA1 as distance from the MPA.

Permutational analysis of multivariate dispersion (PERMDISP) was used to test potential variation in β diversity [31]. Multivariate patterns were visualized through non-metric multidimensional scaling ordination (nMDS). Similarity percentage analysis (SIM-PER) [32] was used to identify the taxa and community descriptors mainly responsible for significant differences.

Sea-urchins had low abundance in both sampling occasions, and potential differences through time were assessed using descriptive statistics.

Results

A total of 88 conspicuous sessile species was found (Table 1). The most speciose groups were algae (35 species, of which 17 rhodophytes, 7 ochrophytes, and 11 chlorophytes) and sponges (22 species); other invertebrate groups were less represented (9 cnidarians, 4 mollusks, 5 serpuloideans, 2 barnacles, 6 bryozoans, and 5 ascidians). Sessile communities had changed significantly between 1993 and 2008 (Table 2). Total species richness passed from 68 species (17 ± 0.78 SE per quadrat) in 1993, to 81 (21 ± 0.63

SE per quadrat) in 2008. Dominant species in 1993 included tall algae such as *Dictyopteris polypodioides* and *Sphaerococcus coronopifolius*, which had nearly disappeared in 2008; amongst the invertebrates, sponges decreased, while hydroids increased.

However, differences between the two sampling years were not consistent among sampling sites (significant Year × Sites) and were affected by the spatial distribution of human pressures, both for taxonomic composition and three-dimensional structure. In particular, the pressures considered by the PCA1 (i.e. protection by the MPA and distance from potential source of organic loading) had a significant role in driving the community change through time (significant Year×PCA1) whereas beach replenishments and tourist harbors played a negligible role (Year×PCA3 not significant). Correspondingly, the MDS plots highlighted marked differences in the direction of change between years (i.e. 1993 vs 2008) among individual sites using both species (Figure 2a) and categories of species three-dimensionality (Figure 2b) as community descriptors. MDS further suggested that sites far from potential sources of organic load or included within the MPA showed a higher degree of change between the two years than the remaining sites.

Although varying significantly among sites, the overall direction of change implied a decreased multivariate dispersion in the pattern of species occurrence and abundance among sites and hence a reduction in β diversity (PERMDISP: P = 0.04), indicative of homogenization of the communities through time.

Even within the MPA, the between-year change greatly varied among individual sites (SIMPER results, Table S3). *Dictyopteriss polypodioides*, a potential canopy forming algal species, disappeared from site J (the furthest from potential impact sources) while *Dictyota dichotoma*, an erect generalist algal species, increased. The perennial algae *Halimeda tuna*, *Peyssonnelia squamaria* and *Flabellia petiolata* decreased at sites I, H and G, respectively, while the **Table 1.** Total list of the sessile taxa found, in systematic

 order by major taxa.

Ochrophyta	Crambe crambe		
Cystoseira zosteroides	Dysidea avara		
Dictyopteris polypodioides	Haliclona cratera		
Dictyota dichotoma	Haliclona fulva		
Dictyota fasciola	Hemimycale columella		
Dictyota implexa	Hymeniacidon perlevis		
Padina pavonica	Ircinia oros		
Stypocaulon scoparium	Ircinia variabilis		
Rhodophyta	Leucosolenia variabilis		
Acrothamnion preissii	Petrosia ficiformis		
algal turf	Phorbas tenacior		
Amphiroa rigida	Spongia lamella		
Amphiroa cryptarthrodia	Cnidaria		
Asparagopsis armata	Aiptasia mutabilis		
Bonnemaisonia asparagoides	Balanophyllia europaea		
Ellisolandia elongata	Caryophyllia inornata		
Jania rubens	Cladocora caespitosa		
Laurencia obtusa	Clavularia crassa		
Lithophyllum incrustans	Eudendrium racemosum		
Lithophyllum stictaeforme	Hydrozoa indet.		
Mesophyllum lichenoides	Parazoanthus axinellae		
Peyssonnelia rubra	Pennaria disticha		
Peyssonnelia squamaria	Mollusca		
Sphaerococcus coronopifolius	Arca noae		
Tricleocarpa fragilis	Chama gryphoides		
Wranaelia penicillata	Thylacodes arenarius		
Chlorophyta	Vermetus triquetrus		
Acetabularia acetabulum	Serpuloidea		
Caulerpa racemosa	Protula tubularia		
Cladophora prolifera	Salmacina dysteri		
Cladophora sp.	Serpula vermicularis		
Codium bursa	Serpulidae indet.		
Codium coralloides	Spirorbidae indet.		
Codium fraaile	Cirripedia		
Flahellia netiolata	Balanus triaonus		
Halimeda tuna	Perforatus perforatus		
Pseudochlorodesmis furcellata	Bryozoa		
Valonia utricularis	Maraaretta cereoides		
Porifera	Rentadeonella violacea		
Acanthella acuta	Retenorella arimaldii		
Agelas oroides	Schizonorella dunkeri		
Avinella damicornis	Schizoporella errata		
Avinella nolynoides			
Avinella vertucosa			
Chondrilla pucula	Clauding langdiformic		
Clothring, clothring			
Clathring contents	Diplosoma listerianum		
Ciumnina contorta	naiocyntnia papillosa		
Cliona viridis	Microcosmus polymorphus		

Within major taxa, species or species groups are ordered alphabetically. Species are named according to the World Register of Marine Species (www. marinespecies.org/).

doi:10.1371/journal.pone.0075767.t001

s category 'medium' three-dimensionality (see Figure 3). Sea urchins were represented in both years by *Paracentrotus lividus* and, especially, *Arbacia lixula*. There were no obvious changes in

and, especially, *Arbacia lixula*. There were no obvious changes in their densities between 1993 and 2008. Sea-urchins were absent from most quadrats in both years, with few exceptions. The highest sea urchin density in 1993 was recorded at site J (4 ind. $m^{-2}\pm 2.5$ SE) and in 2008 at site E (3 ind. $m^{-2}\pm 1.1$ SE).

seasonal algae Stypocaulon scoparium and Dictyota dichotoma increased.

Algal turf greatly increased at all sampling sites. The invasive algae *Caulerpa racemosa* and *Acrothamnion preissii* were recorded in 2008, but not in 1993. In particular, *Caulerpa racemosa* appeared only at

Considering the change in the percentage cover of threedimensionality categories, all sites within the MPA experienced a reduction of the cover by the categories of 'high' or 'medium-high' three-dimensionality and a parallel increase of the cover of the

sites close to potential sources of organic load.

Discussion

In this study we found that an MPA and a network of six SCIs, two major tools used in the Mediterranean Sea for EBM implementation, were ineffective in halting degradation on rocky reef sessile communities. Species composition and architectural complexity changed significantly between 1993 and 2008, before and after the implementation of these conservation measures. Although we compared only two points in time, too few to assess long-term change and to elucidate cause-effect relationships, the strong association of the observed changes in the communities with the documented pressure regime allows generating hypotheses about possible drivers of change. In particular, our analyses indicate that organic loading from sewage outfalls may be a key driver of the observed change in these communities.

The taxonomic composition of rocky reef communities within the MPA became more similar, in 2008, to that of the communities close to sewage outfalls than in the past, thus implying a reduction of β diversity over the whole area. Threedimensional structure decreased owing to the reduction of erect forms (e.g. tall macroalgae) and a parallel increase of flatter forms (e.g. turf-forming or encrusting organisms). In 1993, assemblages within the MPA and far from sources of impact were dominated by erect algae, most of which were canopy-formers. Earlier descriptive information from the 1950s through the 1980s [18,33-38] reported the diffuse presence of dense canopies of the brown algae Dictyopteris polypodioides and Sargassum vulgare (Figure 4), already rare in 1993 and virtually absent in 2008. The loss of habitat complexity is a sign of degradation that has been highlighted as one of the major threats to marine biodiversity worldwide [8,12]. Notwithstanding sessile species diversity showed slightly higher in 2008 than in 1993, reduced habitat complexity is likely to have negatively affected numerous canopy-associated epiphytic and motile organisms, whose diversity and abundance is known to be facilitated by habitat cascades via habitat provision and ecosystem engineering [39,40]. Moreover, we cannot discard the possibility that the occurrence of algal canopies in 1993 hid species that showed obvious only to the visual census of 2008, when canopies were reduced.

In temperate reefs, sea water warming and associated extreme climate events are known to be able to cause extirpation of canopy-forming macroalgae [41–43]. However, lack of erect macroalgae from unprotected sites was already noticed in 1993, i.e. before the heat waves of 1999 and 2003. Similarly, the arrival in the Ligurian Sea of the invasive species *Acrothamnion preissii* and *Caulerpa racemosa* predates the heat waves [44,45]; establishment of invasive species is said to be favored by already stressed ecosystems

Table 2. Results of PERMANOVA analyses on individual taxa and three-dimensional categories.

Source of variation	df	SS	MS	pseudo-F	<i>P</i> (perm)		
Year	1	20266	20266	16.59	0.001		
Site	9	53914	6739	5.51	0.001		
PCA1	1	21050	21050	17.23	0.001		
PCA3	1	2116	2116	1.73	0.075		
Year×Site	9	27053	3381	2.76	0.001		
Year×PCA1	1	5642	5641	4.61	0.001		
Year × PCA3	1	580	580	0.47	0.902		
Residuals	60	70853	1221				
Total	79	201440					
Three-dimensional structure	•						
Source of variation	df	SS	MS	pseudo-F	P (perm)		
Year	1	1118	1118	4.66	0.008		
Site	9	20121	2515	10.48	0.001		
PCA1	1	11193	11193	46.63	0.001		
PCA3	1	526	526	2.19	0.090		
Year×Site	9	3788	473	1.97	0.017		
Year × PCA1	1	2313	2313	9.64	0.001		
Year × PCA3	1	214	214	0.89	0.424		
Residuals	60	13920	240				
Total	79	53197					

Bold numbers indicate significant results.

doi:10.1371/journal.pone.0075767.t002

[46,47]. A shift from erect-dominated to turf-dominated algal communities can be produced by nutrient enrichment from sewage outfalls [48]. Our sampling sites were relatively far from sewage outlet; therefore, no indicator species of organic pollution was detected. However, algal turf and many of the algal species first appeared or found more abundant in 2008 are known to be comparatively tolerant to lowered water quality with respect to those dominating in 1993 [49].

In the Tigullio Gulf, the undiminished input of organic load has possibly expanded its effects further away from its sources in recent vears. The assemblages close to these chronic sources already showed a bi-dimensional structure in 1993 and have likely changed less as compared to the furthest sites, still architecturally complex in 1993. The sequence of disturbance is known to affect the final structure of communities whose response depends also on their previous history [50]. The spatio-temporal variation of stressors intensities is likely to be the main driver of the delay in degradation observed in the MPA. However, the establishment of the MPA in 1998 and the implementation of SCIs were ineffective in reversing the trend toward the deterioration of communities. Finding the alien alga Caulerpa racemosa in 2008 constituted a further signal of alteration. C. racemosa has been recognized as a strong invader [51] and its fast spread is favored by the presence of algal turf [52]. Turf development in the protected sites anticipated the settlement of C. racemosa, which has been subsequently recorded also in the Portofino MPA [53].

Our results suggest sequential community change under the combined influence of human pressure and invasive species. Reef communities are undergoing a shift from canopy-dominated to turf-dominated, and then to a *C. racemosa*-dominated state. This

trajectory towards ecosystem deterioration has not been hampered by protection measures. Paradoxically, protection itself might even have contributed to the observed community shift. Fishing restriction through MPA enforcement may have a dual effect on reef communities: on one hand, it recovers the abundance of large carnivorous fish, which have the desirable effect of controlling seaurchin overgrazing [54]; on the other hand, it also favors large herbivorous fishes such as Sarpa salpa. Sea urchin densities did not change between 1993 and 2008 and, although published quantitative data about their densities prior to 1993 are not available, field observations in the 1980s by two of us (CNB and CM) suggest no obvious change with time. On the contrary, the abundance of S. salpa was shown to have increased after the creation of the Portofino MPA [15]. Increased grazing by Sarpa salpa is capable of driving algal assemblages towards a turfdominated state [55].

Anthropogenic stressors and climate change may act in synergy on temperate reef communities, lowering their resilience to perturbations [56]. While climate and other global impacts require international actions, regional management practices may help reducing local impacts. Managing temperate reefs requires efforts to reduce nutrient and organic inputs from land sources and pollution by boat traffic. Best coastal management practices may promote favorable conditions to confer canopyforming species with resistance and resilience against pressures that cannot be managed locally, such as sea water warming, storms, and introduced species.

MPAs, which are designed mainly to regulate fishing activities, are largely ineffective in managing land-based pressures (e.g. outfalls, river discharge) and the spread of invasive species. In this



Figure 2. MDS plots calculated on species (a) and threedimensional categories (b). Black characters and arrows refer to sampling sites inside the MPA, grey ones to sampling sites outside the MPA.

doi:10.1371/journal.pone.0075767.g002

context, SCIs or other complementary measures need to play a significant role in addressing diverse and escalating threats. Yet, SCIs implementation regulated a limited set of pressures (i.e. anchoring and the building of new coastal structures) that may not be the major drivers of change, especially in cases as the Tigullio Gulf where multiple and chronic pressures act together. At present, SCIs represent a 'passive' application of the legislative constraints in the coastal zones. Legislation, of course, plays a crucial role for conservation. However, legislative constraints alone are not enough to achieve effective EBM [3]: managers and scientists should apply and use the legislative instruments in a scientifically sound way. For SCIs to become effective EBM tools, the selection of the crucial pressures to be managed should not be driven by existing laws, but should rather be dictated by the actual pressure regime and local ecological characteristic of different coastal zones (Figure 5).

Small-scale protection measures and SCIs will not be effective if key large scale pressures, such as those coming from adjacent unprotected zones or linked to global change, are not identified. Comprehensive management plans should address multiple pressures through a suite of policy and management tools: carefully designed monitoring is critical for the early detection and eradication of invasive species, for the upgrade of water treatment plants, and for the regulation of coastal development in proximity to sensitive reef and seagrass habitats.



Figure 3. Cover of three dimensional categories for each sites in 1993 (white) and 2008 (grey). a): high; b) medium-high; c) medium; d) medium-low; e) low. Note that scales on Y axis are different. The vertical dotted line separates the sampling sites within the MPA (black characters) from those outside (grey characters). doi:10.1371/journal.pone.0075767.g003



Figure 4. Change in reef community at Portofino, in the vicinity of sampling site J: a) *Sargassum vulgare* and *Dictyopteris* polypodioides canopy in 1981; b) *Sarpa salpa* grazing in a turf-dominated environment in 2009. doi:10.1371/journal.pone.0075767.q004

Monitoring programs that are too narrow in scope, e.g., involving only ecosystems within MPAs, may fail in the identification of relevant pressures that often act at larger scales [57] and in the quantification of the cumulative impacts of multiple pressures [1,7]. Only when pressures are exclusively local, monitoring based on hierarchical designs within the protected area and according to enforcement levels may be effective in providing managers with valuable solutions [58]. When multiple pressures originate outside the boundaries of the protected area and have wider impact, monitoring programs should be designed across gradients at the appropriate scales to address relevant pressures and resulting cumulative impacts (Figure 5). This will help identifying the ultimate causes of deterioration and setting management priorities to reverse ecosystem trajectory.

Supporting Information

Table S1Characteristics of the main pressures in the study area.Information on beach replenishment was available from 2003.(DOCX)



Figure 5. Decision flow-chart for implementing EBM and choosing effective monitoring design in coastal zones to be protected. White boxes: steps required for ecosystem based evaluation of relevant pressures; grey boxes: policy consequences of ecosystem-based analysis. doi:10.1371/journal.pone.0075767.g005

Table S2Significant correlations of the individual pressures withthe first three axes of the Principal Component Analysis (PCA).(DOCX)

Table S3 Results of SIMilarity PERcentage (SIMPER) analysis identifying taxa major contributing to differences between Years and among Sites. Grey letters indicate sampling sites outside the MPA while black letters indicate sampling sites within the MPA boundaries.

(DOCX)

Figure S1 a) Annual and seasonal mean (+SE) of chlorophyll-a concentration in the period 1985–1994 (grey bars) and in the period 2002–2008 (black bars). **b**) Annual and seasonal mean (+SE) of water transparency (measured with Secchi disk) in the period 1985–1994 (grey bars) and in the period 2002–2008 (black bars). **c**) Potential organic load (expressed in term of equivalent inhabitants) calculated for the 7 administrations facing the coastal zone of the study area following the method described by Lopez y Royo et al. (2010); C: Camogli; L: Lavagna; P: Portofino; R:

References

- Halpern BS, Walbridge S, Selkoe KA, Kappel CV, Micheli F (2008) A global map of human impact on marine ecosystems. Science 319: 948–952.
- Allison GW, Lubchenco J, Carr MH (1998) Marine reserves are necessary but not sufficient for marine conservation. Ecol Appl 8: S79–S92.
- Montefalcone M, Albertelli G, Morri C, Parravicini V, Bianchi CN (2009) Legal protection is not enough: *Posidonia oceanica* meadows in marine protected areas are not healthier than those in unprotected areas of the northwest Mediterranean Sea. Mar Pollut Bull 58: 515–519.
- McLeod KL, Lubchenco J, Palumbi SR, Rosenberg AA (200). Scientific consensus statement on marine ecosystem-based management. Communication Partnership for Science and the Sea, www.compassonline.org/pdf_files/EBM_ Consensus_Statement_v12.pdf.
- Leslie HM, McLeod KL (2007) Confronting the challenges of implementing marine ecosystem-based management. Front Ecol Environ 5: 540–548.
- Olsson P, Folke C, Hughes TP (2008) Navigating the transition to ecosystembased management of the Great Barrier Reef, Australia. Proc Natl Acad Sci USA 105: 9489–9494.
- Lester SE, McLeod KL, Tallis H, Ruckelshaus M, Halpern BS, et al. (2010) Science in support of ecosystem-based management for the US West Coast and beyond. Biol Conserv 143: 576–587.
- Airoldi L, Beck MW (2007) Loss, status and trends for coastal marine habitats of Europe. Ann Rev Mar Sci 45: 345–405.
- Sano M, Gonzalez-Riancho P, Areizaga J, Medina R (2010) The strategy for coastal sustainability: a Spanish initiative for ICZM. Coast Manag 38: 76–96.
- Russell BD, Harley CDG, Wernberg T, Mieszkowska N, Widdicombe S, et al. (2012) Predicting ecosystem shifts requires new approaches that integrate the effects of climate change across entire systems. Biol Lett 8: 164–166.
- Bianchi CN, Cocito S, Morgigni M, Morri C, Peirano A, et al. (1994) Benthos del Golfo Tigullio: fondi duri e prati di *Cymodocea*. ENEA CRAM Santa Teresa, La Spezia.
- Airoldi L, Balata D, Beck MW (2008) The Gray Zone: relationships between habitat loss and marine diversity and their applications in conservation. J Exp Mar Biol Ecol 366: 8–15.
- Micheli F, Benedetti-Cecchi L, Gambaccini S, Bertocci I, Borsini C, et al. (2005) Cascading human impacts, marine protected areas, and the structure of Mediterranean reef assemblages. Ecol Monog 75: 81–102.
- Parravicini V, Thrush SF, Chiantore M, Morri C, Croci C, et al. (2010) The legacy of past disturbance: chronic angling impairs long-term recovery of marine epibenthic communities from acute date-mussel harvesting. Biol Conserv 143: 2435–2440.
- Guidetti P, Milazzo M, Bussotti S, Molinari A, Murenu M, et al. (2008) Italian marine reserve effectiveness: does enforcement matter? Biol Conserv 141: 699– 709.
- Crain CM, Kroeker K, Halpern BS (2008) Interactive and cumulative effects of multiple human stressors in marine systems. Ecol Lett 11: 1304–1315.
- Asnaghi V, Chiantore M, Bertolotto RM, Parravicini V, Cattaneo-Vietti R, et al. (2009) Implementation of the European Water Framework Directive: Natural variability associated with the CARLIT method on the rocky shores of the Ligurian Sea (Italy). Mar Ecol 30: 505–513.
- Morri C, Bianchi CN, Damiani V, Peirano A, Romeo G, et al. (1988) The marine environment between Punta della Chiappa and Sestri Levante (Ligurian Sea): ecotipological outline and proposal of a biocenotic map. Boll Mus Ist Biol Univ Genova 52: 213–231.
- Danovaro R, Fabiano M (1997) Seasonal changes in quality and quantity of food available for benthic suspension-feeders in the Golfo Marconi (North-western Mediterranean). Estuar Coast Shelf Sci 44: 723–736.

Rapallo; Sa: Santa Margherita; Se: Sestri Levante; Z: Zoagli; grey bars: data relative to 1991; black bars: data relative to 2001. (TIF)

Acknowledgments

Data of 1993 were collected in the frame of an investigation coordinated by the ENEA Marine Environmental Research Centre of La Spezia: thanks are due to S. Cocito, F. Degl'Innocenti, M. Morgigni, A. Peirano and S. Sgorbini, who participated in the field activities. Data used in this study are available from the authors upon request.

Author Contributions

Conceived and designed the experiments: VP FM CNB MM CM EV MC PP. Performed the experiments: VP MM EV MC CNB CM PP. Analyzed the data: VP FM MM EV CNB. Contributed reagents/materials/analysis tools: CNB FM CM PP MC. Wrote the paper: VP FM MM CM EV MC PP CNB.

- Misic C, Covazzi Harriague A (2008) Organic matter recycling in a shallow coastal zone (NW Mediterranean): The influence of local and global climatic forcing and organic matter lability on hydrolytic enzyme activity. Cont Shelf Res 28: 2725–2735.
- Lopez y Royo C, Silvestri C, Pergent G, Casazza G (2009) Assessing humaninduced pressures on coastal areas with publicly available data. J Env Manag 90: 1494–1501.
- Bianchi CN, Morri C (2003) Global sea warming and "tropicalization" of the Mediterranean Sea: biogeographic and ecological aspects. Biogeographia 24: 319–327.
- Cerrano C, Bavestrello G, Bianchi CN, Cattaneo-Vietti R, Bava S, et al. (2000) A catastrophic mass-mortality episode of gorgonians and other organisms in the Ligurian Sea (North-western Mediterranean), summer 1999. Ecol Lett 3: 284– 293.
- Garrabou J, Coma R, Bensoussan N, Bally M, Chevaldonné P, et al. (2009) Mass mortality in Northwestern Mediterranean rocky benthic communities: effects of the 2003 heat wave. Glob Change Biol 15: 1090–1103.
- Parravicini V, Guidetti P, Morri C, Montefalcone M, Donato M, et al. (2010) Consequences of sea water temperature anomalies on a Mediterranean submarine cave ecosystem. Estuar Coast Shelf Sci 86: 276–282.
- Martí R, Uriz MJ, Ballesteros E, Turon X (2005) Seasonal variation in the structure of three Mediterranean algal communities in various light conditions. Estuar Coast Shelf Sci 64: 613–622.
- Bianchi CN, Pronzato R, Cattaneo-Vietti R, Benedetti-Cecchi L, et al. (2004) Mediterranean marine benthos: a manual of methods for its sampling and study. Hard bottoms. Biol Mar Medit 11: 185–215.
- Alvarez-Filip L, Gill JA, Dulvy NK, Perry AL, Watkinson AR, et al. (2011) Drivers of region-wide declines in architectural complexity on Caribbean reefs. Coral Reefs 30: 1051–1060.
- Mardia KV, Kent JT, Bibby JM (1980) Multivariate Analysis. Academic Press, London.
- Anderson MJ, Ter Braak CJF (2003) Permutation tests for multi-factorial analysis of variance. J Stat Comp Sim 73: 85–113.
- Anderson MJ, Ellingsen KE, McArdle BH (2006) Multivariate dispersion as a measure of beta diversity. Ecol Lett 9: 683–693.
- Clarke KR, Gorley RN (2006) PRIMER v6: User Manual/Tutorial. PRIMER-E Ltd, Plymouth.
- Tortonese E (1958) Bionomia marina della regione costiera fra Punta della Chiappa e Portofino (Riviera ligure di levante). Archo Oceanogr Limnol 11: 167–210.
- Tortonese E (1961) Nuovo contributo alla conoscenza del bentos della scogliera ligure. Archo Oceanogr Limnol 12: 163–183.
- Tortonese E (1962) Recenti ricerche sul bentos in ambienti litorali del mare Ligure. Pubbl Staz Zool Napoli 32: 99–116.
- Melegari GE (1973) Portofino sub: guida alla conoscenza dei fondali del promontorio di Portofino. ERGA, Genova.
- Pisano E, Bianchi CN, Matricardi G, Relini G (1982) Accumulo della biomassa su substrati artificiali immersi lungo la falesia di Portofino (Mar Ligure). Atti del Convegno delle Unità Operative afferenti ai sottoprogetti Risorse Biologiche ed Inquinamento Marino. CNR, Roma: 93–105.
 Pisano E, Bianchi CN, Relini G. (1980) Insediamento su substrati artificiali
- Pisano E, Bianchi CN, Relini G. (1980) Insediamento su substrati artificiali immersi lungo la falesia di Portofino (Mar Ligure): metodologie e dati preliminari. Mem Biol Mar Oceanogr 10: 269–274.
- Thomsen MS, Wernberg T, Altieri A, Tuya F, Gulbransen D, et al. (2010) Habitat cascades: the conceptual context and global relevance of facilitation cascades via habitat formation and modification. Integr Comp Biol 50: 158–175.

- Wernberg T, Thomsen MS, Tuya F, Kendrick GA (2011a) Biogenic habitat structure of seaweeds change along a latitudinal gradient in ocean temperature. J Exp Mar Biol Ecol 400: 264–271.
- 41. Smale DA, Wernberg T (2013) Extreme climatic event drives range contraction of a habitat-forming species. Proc R Soc B 280: 20122829.
- Wernberg T, Smale DA, Tuya F, Thomsen MS, Langlois TJ, et al. (2013) An extreme climatic event alters marine ecosystem structure in a global biodiversity hotspot. Nature Clim Change 3: 78–82.
- 43. Duarte L, Viejo RM, Martínez B, de Castro M, Gómez-Gesteira M, et al. (2013) Recent and historical range shifts of two canopy-forming seaweeds in North Spain and the link with trends in sea surface temperature. Acta Oecol 51: 1–10.
- Bianchi CN, Morri C (1994) Southern species in the Ligurian Sea (northern Mediterranean): new records and a review. Boll Mus Ist Biol Univ Genova 58– 59: 181–197.
- Bussotti S, Conti M, Guidetti P, Martini F, Matricardi G (1996) First record of *Caulerpa racemosa* (Forsskål) J. Agardh along the coast of Genoa (North-Western Mediterranean). Doriana 6: 1–5.
- MacDougall AS, Turkington R (2005) Are invasive species the drivers or passengers of change in degraded ecosystems? Ecology 86: 42–55.
 Didham RK, Tylianakis JM, Gemmell NJ, Rand TA, Ewers RM (2007)
- Didham RK, Tylianakis JM, Gemmell NJ, Rand TA, Ewers RM (2007) Interactive effects of habitat modification and species invasion on native species decline. Trends Ecol Evol 22: 489–496.
- Claudet J, Fraschetti S (2010) Human-driven impacts on marine habitats: a regional meta-analysis in the Mediterranean Sea. Biol Conserv 143: 2195–2206.
- Orfanidis S, Panayotidis P, Stamatis N (2001) Ecological evaluation of transitional and coastal waters: a marine benthic macrophytes based model. Med Mar Sci 2: 45–65.

- Fukami T (2001) Sequence effects of disturbance on community structure. Oikos 92: 215–224.
- Montefalcone M, Albertelli G, Morri C, Bianchi CN (2010) Pattern of wide-scale substitution within *Posidonia oceanica* meadows of NW Mediterranean Sea: invaders are stronger than natives. Aquat Conserv 20: 507–515.
- Bulleri F, Benedetti-Cecchi L (2008) Facilitation of the introduced green alga *Caulerpa racemosa* by resident algal turfs: experimental evaluation of underlying mechanisms. Mar Ecol Prog Ser 364: 77–86.
- Cattaneo-Vietti R, Albertelli G, Aliani S, Bava S, Bavestrello G, et al. (2010) The Ligurian Sea: present status, problems and perspectives. Chem Ecol 26: 319– 340.
- Sala E, Boudouresque CF, Harmelin-Vivien M (1998) Fishing, trophic cascades, and the structure of algal assemblages: evaluation of an old but untested paradigm. Oikos 82: 425–439.
- 55. Ruitton S, Francour P, Boudouresque CF (2000) Relationships between algae, benthic herbivorous invertebrates and fishes in rocky sublittoral communities of a temperate sea (Mediterranean). Estuar Coast Shelf Sci 50: 217–230.
- Wernberg T, Russell BD, Moore PJ, Linge SD, Smale DA, et al. (2011) Impacts of climate change in a global hotspot for temperate marine biodiversity and ocean warming. J Exp Mar Biol Ecol 400: 7–16.
- Hewitt JE, Thrush SF, Dayton PK, Bonsdorff E (2007) The effect of spatial and temporal heterogeneity on the design and analysis of empirical studies of scaledependent systems. Am Nat 169: 398–408.
- Greco S, Notarbatolo di Sciara G, Tunesi L (2004) 'Sistema Afrodite': an integrated programme for the inventorying and monitoring of the core zones of the Italian marine protected areas. Aquat Conserv 14: S119–S122.