

RESEARCH ARTICLE

Identification of an Unclassified Paramyxovirus in *Coleura afra*: A Potential Case of Host Specificity

Gael D. Maganga^{1,2*}, Mathieu Bourgarel^{1,3,4}, Judicael Obame Nkoghe^{1,5}, Nadine N'Dilimabaka¹, Christian Drosten⁶, Christophe Paupy^{1,5}, Serge Morand^{3,7}, Jan Felix Drexler⁶, Eric M. Leroy^{1,5}

1. Centre International de Recherches Médicales de Franceville (CIRMF), Franceville, Gabon, 2. Institut National Supérieur d'Agronomie et de Biotechnologies (INSAB), Franceville, Gabon, 3. Centre de Coopération Internationale en Recherche Agronomique pour le Développement (CIRAD), UPR AGIRs, Montpellier, France, 4. Centre de Coopération Internationale en Recherche Agronomique pour le Développement (CIRAD), UPR AGIRs, Harare, Zimbabwe, 5. Institut de Recherche pour le Développement (IRD), UMR 224 (MIVEGEC), IRD/CNRS/UM1/UM2, Montpellier, France, 6. Institute of Virology, University of Bonn Medical Centre, Bonn, Germany, 7. des Sciences de l'Evolution, CNRS-UM2, CC065, Université de Montpellier 2, Montpellier, France

*gael_maganga@yahoo.fr



CrossMark
click for updates

OPEN ACCESS

Citation: Maganga GD, Bourgarel M, Obame Nkoghe J, N'Dilimabaka N, Drosten C, et al. (2014) Identification of an Unclassified Paramyxovirus in *Coleura afra*: A Potential Case of Host Specificity. PLoS ONE 9(12): e115588. doi:10.1371/journal.pone.0115588

Editor: Wanda Markotter, University of Pretoria, South Africa

Received: July 25, 2014

Accepted: December 1, 2014

Published: December 31, 2014

Copyright: © 2014 Maganga et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability: The authors confirm that all data underlying the findings are fully available without restriction. All relevant data are within the paper.

Funding: This work was supported by Global Viral Forecasting, a "Fonds de Solidarité Prioritaire" grant from the Ministère des Affaires Etrangères de la France (FSP n° 2002005700). CIRMF is supported by the government of Gabon, Total-Fina-Elf Gabon, and the Ministère des Affaires Etrangères de la France. This study was also made possible by the generous support of the American people through the United States Agency for International Development (USAID), Emerging Pandemic Threats PREDICT. The contents are the responsibility of the authors and do not necessarily reflect the views of USAID or the United States Government. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: CIRMF (Centre International de Recherche Médicale de Franceville) is partly supported by Total-Fina-Elf

Abstract

Bats are known to harbor multiple paramyxoviruses. Despite the creation of two new genera, *Aquaparamyxovirus* and *Ferlavirus*, to accommodate this increasing diversity, several recently isolated or characterized viruses remain unclassified beyond the subfamily level. In the present study, among 985 bats belonging to 6 species sampled in the Belinga caves of Gabon, RNA of an unclassified paramyxovirus (Belinga bat virus, BelPV) was discovered in 14 African sheath-tailed bats (*Coleura afra*), one of which exhibited several hemorrhagic lesions at necropsy, and viral sequence was obtained in two animals. Phylogenetically, BelPV is related to *J virus* and *Beilong virus* (BeiPV), two other unclassified paramyxoviruses isolated from rodents. In the diseased BelPV-infected *C. afra* individual, high viral load was detected in the heart, and the lesions were consistent with those reported in wild rodents and mice experimentally infected by *J virus*. BelPV was not detected in other tested bat species sharing the same roosting sites and living in very close proximity with *C. afra* in the two caves sampled, suggesting that this virus may be host-specific for *C. afra*. The mode of transmission of this paramyxovirus in bat populations remains to be discovered.

Gabon. There are no patents, products in development, or marketed products to declare. This does not alter the authors' adherence to all the PLOS ONE policies on sharing data and materials, as detailed in the online guide for authors.

Introduction

Members of the Paramyxoviridae family are pleomorphic enveloped viruses [1] divided into two subfamilies, *Paramyxovirinae* and *Pneumovirinae*. *Paramyxovirinae* has recently been subdivided into seven genera: *Aquaparamyxovirus*, *Avulavirus*, *Ferlavirus*, *Henipavirus*, *Morbillivirus*, *Respirovirus*, and *Rubulavirus* (<http://ictvonline.org/virusTaxonomy.asp?version=2012>). Viruses of this family affect a wide range of animals, including primates, birds, carnivores, ungulates, snakes, cetaceans and humans, and cause a wide variety of infections, such as measles, mumps, pneumonia and encephalitis in humans, and distemper, peste des petits ruminants, Newcastle disease and respiratory tract infections in animals. However, several paramyxoviruses (PVs) have not been classified into any of these seven genera, including *Nariva virus* (NarPV) [2], *Mossman virus* (MosPV) [3], *Beilong virus* (BeiPV) [4], *J virus* (JPV) [5,6], *Tupaia paramyxovirus* (TupPV) [7] and *Tailam virus* [8], all of which belong to a group of novel paramyxoviruses isolated from wild animals, as well as *Salem virus* isolated from horses [9]. Among them, only JPV has been shown to be pathogenic, causing extensive haemorrhagic lesions in rodents [6]. Horizontal transmission is the principal mode of intraspecies PV infection, suggesting that contaminated faeces, urine or saliva may be responsible for spillover to other species [10].

Bats have a close evolutionary relationship with several genera of mammalian paramyxoviruses [11]. Otherwise, bat-borne paramyxoviruses are in close relationship to known paramyxoviruses of mammalian. These small mammals are known to harbour a broad diversity of PVs, including emergent henipaviruses (Nipah virus and Hendra virus) and rubulaviruses [Menangle virus, Tioman virus, Mapuera virus, and Tuhoko virus 1, 2 and 3 (ThkPV-1, ThkPV-2 and ThkPV-3)]. A very broad diversity of paramyxoviruses, including Henipa-, Rubula-, Pneumo- and Morbilli-related viruses, have been detected in six of ten tested bat families [11]. Whereas most of the viruses identified in bats do not seem to cause clinical disease in these animals, there have been reports of rabid bats [12,13] and of unusually large numbers of animals succumbing to infection by rabies virus [14].

As part of a large-scale investigation of viral diversity in bats and of associated zoonotic risks, we have previously detected a bat paramyxovirus in one insectivorous African sheath-tailed bat (*Coleura afra*) [11], exhibiting several hemorrhagic lesions at necropsy. We therefore examined occurrence of this bat paramyxovirus in other bats.

Materials and Methods

Ethics statements

The study was conducted in the Belinga mountains (northeast Gabon), where Ebola outbreaks occurred in 1994–1996 and 2001–2002. All the capture events, animal handling, euthanasia and transfer of samples across country borders were

performed in accordance with the guidelines of the American Society of Mammalogists (<http://www.mammalsociety.org/committees/animal-care-and-use>) [15]: bats were captured following recommendations by Kunz and Parsons [16] and identified by trained field biologist. Captured bats were removed carefully from nets as soon as possible to minimize injury, drowning, strangulation, or stress. Safe and humane euthanasia was achieved through the use of inhalant anaesthetic (halothane) prior to autopsy.

All work (capture, euthanasia and autopsy) was carried out with authorization from the Gabonese Ministry of Water and Forestry (Département de la Faune et de la Chasse – Authorization N°003/MEFEPA/SG/DGEF/DFC and N°0021/MEFEPA/SG/DGEF/DFC) and the Gabonese Ministry of Higher Education, Scientific Research and Innovation (Centre National de la Recherche Scientifique et Technique – Authorization N° AR0027/10/MENESRI/CENAREST/CG/CST/CSAR).

Bat sampling

Samples of liver, spleen, kidney, lung, heart, gut, brain and salivary glands were collected, stored in liquid nitrogen and transferred to the CIRMF laboratory (Centre International de Recherches Médicales de Franceville, Gabon), where they were stored at -80°C until analysis. Blood samples were also collected, except from the smallest individuals (body mass <12 g). A total of 985 bats (Table 1) were sampled from the caves of Faucon ($1^{\circ}07'\text{N}$, $13^{\circ}20'\text{E}$), Zadié ($0^{\circ}98'\text{N}$, $13^{\circ}19'\text{E}$) and Batouala ($0^{\circ}82'\text{N}$, $13^{\circ}45'\text{E}$) in July 2009, December 2009 and June 2010.

Interestingly, at autopsy, only one *C. afra* out of the 26 individuals, from Batouala cave, exhibited diarrhea and severe hemorrhagic lesions in both thoracic and abdominal organs, along with lung congestion and pleurisy.

Viral PCR screening

Virological screening of the diseased bat was performed to identify the cause of these lesions. The screening process included the whole Paramyxoviridae family and also filoviruses [Marburg virus (MARV) and Zaire Ebola virus (EBOV)], given the nature of the lesions and the known filovirus tropism for bats. Briefly, approximately 100 mg each of this animal's liver and spleen were pooled and crushed in 600 μl of cold PBS in a ball-mill tissue grinder (Genogrinder 2000, Spex Centriprep). Total RNA was extracted using a Biorobot EZ1 and the EZ1 RNA tissue mini kit (Qiagen) according to the manufacturer's guidelines. The RNA was then tested for paramyxoviruses, using three heminested reverse transcription-PCR (hnRT-PCR) assays targeting the polymerase gene [17], and also for Marburg virus [18] and Zaire Ebola virus [19]. However, the screening was extended to Arenaviruses, Flaviviruses, Alphaviruses, polyomaviruses, Orthopoxviruses, Parapoxviruses, Influenza viruses, Lyssaviruses and Rhabdoviridae family.

Table 1. Overview of specimens collected in Belinga caves and tested by specific BelPV qPCR assay.

Sampling sites	Bats species	Total	Positive RT-PCR	
Batouala cave	<i>Coleura afra</i>	26	6	
	<i>Hipposideros cf. ruber</i>	117	0	
	<i>Hipposideros gigas</i>	59	0	
	<i>Miniopterus inflatus</i>	161	0	
	<i>Rousettus aegyptiacus</i>	18	0	
Faucon cave	<i>Coleura afra</i>	68	8	
	<i>Hipposideros cf. ruber</i>	281	0	
	<i>Hipposideros gigas</i>	121	0	
	<i>Miniopterus inflatus</i>	10	0	
	<i>Rhinolophus alcyone</i>	14	0	
Zadié cave	<i>Rousettus aegyptiacus</i>	110	0	
Total		985	14	
Belinga caves	Diptera: Nycteribiidae			
	<i>Eucampsipoda africana</i>	51	0	
	<i>Nycteribia schmidlii scotti</i>	26	0	
	<i>Penicillidia fulvida</i>	21	0	
	Diptera: Streblidae			
	<i>Brachytarsina allaudi</i>	3	0	
	<i>Raymondia huberi huberi</i>	1	0	
	Diptera: Culicidae			
	<i>Uranotaenia nigromaculata</i>	10	0	
	<i>Culex wigglesworthi</i>	320	0	
	Total		432	0

doi:10.1371/journal.pone.0115588.t001

Paramyxovirus detection in bat populations

To further investigate the presence of the virus in bat populations, a strain-specific real-time RT-PCR assay (primers: GB09-478-F, 5'-GGCGGCTCTTAAAAGT-GAATG-3'; GB09-478-R, 5'-GCGGGGTCAAATTGGTCAT-3'; probe: GB09-478-P, 5'-TCCAGCACAAACATATCCGAGAAGGCTAG-3') was designed within the initial PCR fragment and was used to test total RNA extracted from mixed liver and spleen samples from each of all the other bat species. The amplification was performed in a final volume of 25 µl, containing, 12.5 µl TaqMan^R 2X PCR Master Mix (Applied Biosystems), 0.5 µl each primer and probe (10 µM), 1 µl bovine serum albumin (1 µg/µl) (Invitrogen), 5 µl cDNA and RNase-free water

(Invitrogen). Amplification generally involved 2 min at 55°C, 10 min at 95°C followed by 45 cycles of 15s at 95°C and 1 min at 58°C.

Virus distribution in organs of infected bats

In order to determine the organ distribution of this virus in infected bats, total RNA was extracted from heart, liver, spleen, kidney, lung, intestine and brain samples from all 14 real-time RT-PCR-positive bats, as described previously, and screened, using the same strain-specific real-time RT-PCR assay shown above.

Entomological study

Transmission by hematophagous arthropods was studied. Hematophagous arthropods were collected inside Faucon and Zadié caves. Mosquitoes were sampled using CDC light traps whereas bat-flies were manually collected on bats (*Coleura afra*, *Miniopterus inflatus*, *Hipposideros cf. ruber* and *Rousettus aegyptiacus*) trapped in the Faucon and Zadié caves between November and December 2009, June 2010, and January, March and April 2011. After the morphologic species determination, insects were crushed by monospecific pools (up to 10 specimens for mosquitoes, and between 1 and 5 for bat-flies) in 300 µl PBS. Total RNA was extracted using 100 µl of the supernatant from each pool by using the RNeasy Mini Kit (Qiagen) and then was tested with the specific real-time RT-PCR assay described above.

Blast and phylogenetic analysis

The viral sequences obtained were first compared to those available in the public database using the algorithm "Blastn" from BLAST program [20] and then aligned with homologous sequences of *Paramyxoviridae* reference strains from GenBank, using the MEGA program version 5 [21]. Bayesian inference of phylogeny was done using MrBayes V.3.2 software and the GTR+G+I nucleotide substitution model [22] for two million generations with a burn-in of 25%.

Results

Identification and molecular characterization

One of three hnRT-PCR, Respiro-, Morbilli- and Henipaviruses PCR (RMH-PCR) assays yielded an amplicon of 559 base pairs (bp) from diseased bat, in collaborative work with the Bonn Institute of Virology Bonn (Germany). The PCR product was sequenced with dye terminator chemistry (Applied Biosystems). No other virus was detected in the diseased bat.

Blast and phylogenetic analysis confirmed that the sequence from diseased bat, designated BatPV GB09 478 (Genbank accession number HQ660155), belonged to the Paramyxoviridae family. It showed 65% and 66% nucleotide identity, respectively, with JPV and BeiPV, and had pairwise nucleotide identities of 36–

42% with the Henipavirus group and 37–40% with the Morbillivirus group. Phylogenetic analysis indicated that this bat PV, that we named Belinga bat virus (BelPV), clustered with unclassified paramyxoviruses and was located between the genera *Morbillivirus* and *Henipavirus* (Fig. 1).

In this study, among all animals tested by the strain-specific real-time RT-PCR assay only 14 *C. afra* bats, including the diseased animal, were positive. Bat samples positive by specific real-time RT-PCR were then tested with RMH-hemi-nested-PCR for molecular characterization. Another 439-bp PCR product was amplified from one liver-spleen pool from a second *C. afra* specimen, and was sequenced. These two phylogenetically related sequences (Fig. 1), designated BatPV GB09 478 and BatPV GB09 450, displayed 100% of nucleotide identity.

Prevalence and organ distribution of BelPV in *Coleura afra*

BelPV-specific RNA was detected only in *C. afra*, with a prevalence of 14.9% (14/94 of the *C. afra* individuals sampled from the Faucon and Batouala caves) (Table 1). Interestingly, no bats belonging to the other five species tested positive.

BelPV RNA was detected in the heart (8/12) and liver (5/14), at low and variable loads (Table 2 and Fig. 2). No significant difference in the BelPV RNA detection rate was found between the two organs ($\chi^2=2.476$, $df=1$, $p>0.20$). In the bat exhibiting severe hemorrhagic lesions (GB09 478), both the liver and heart were BelPV-positive, with a higher viral load in the heart (Ct value =28).

Entomological study

In total, 432 arthropods were collected inside Faucon and Zadié caves, including Culicidae (10 *Uranotaenia nigromaculata* and 320 *Culex wigglesworthi*), Nycteribiidae (51 *Eucampsipoda africana*, 26 *Nycteribia schmidlii scotti* and 21 *Penicillidia fulvida*) and Streblidae (*Brachytarsina allaudi* and 1 *Raymondia huberi huberi*) (Table 1). No arthropods tested were positive for Belinga Bat Virus (Table 1).

Discussion

The BelPV nucleotide sequence obtained showed similarity with the JPV and BeiPV sequences. BelPV has previously been reported to hold a phylogenetic position between the genera *Henipavirus* and *Morbillivirus*. The same phylogenetic position had been observed with MosPV and J-V [4].

In this study, organs with high BelPV concentrations are different from those found with high paramyxovirus concentrations in pteropodids and microchiroptera bats. In microchiroptera bats from Brazil, spleen has been found more positive than the others organs with highest viral load, as in *Eidolon helvum* (megachiroptera bat) in Africa [11]. However, in our study majority of spleen were not available.




Fig. 1. Phylogenetic tree based on a 439-basepair fragment of the polymerase gene (L) of members of the *Paramyxoviridae* family. Sequences generated in this study are highlighted in red. Bayesian posterior probabilities are shown; values < 0.80 were removed for clarity. The viruses are designated as follows (virus abbreviation/typical host/accession number of reference sequences in brackets): HeV = Hendra virus, NiV = Nipah virus, BatPV = Bat paramyxovirus, BeiPV = Beilong virus, JPV = J virus, MosPV = Mossman virus, TupPV = Tupaia paramyxovirus, NarPV = Nariva virus, PDV = Phocine distemper virus, CDV = Canine distemper virus, CeMV DMV = Cetacean morbillivirus strain dolphin morbillivirus, MeV = Measles virus, PPRV = Pestivirus, Rinderpest virus, FdIPV = Fer-de-lance virus, PSPV = Pacific salmon paramyxovirus, ASPV = Atlantic salmon paramyxovirus, SeV = Sendai virus, bPIV3 = Bovine parainfluenza virus 3, hPIV1 = Human parainfluenza virus 1, hPIV3 = Human parainfluenza virus 3, SwPIV3 = Swine parainfluenza virus 3, NDV = Newcastle disease virus, PigeonPMV = Pigeon paramyxovirus, AMPV9 = Avian paramyxovirus type 9, AMPV6 = Avian paramyxovirus type 6, AMPV2 = Avian paramyxovirus type 2, AMPV3 = Avian paramyxovirus type 3, AMPV4 = Avian paramyxovirus type 4, PIV5 = parainfluenza virus 5, SV41 = Simian virus 41, MenPV = Menangle paramyxovirus, MprPV = Mapuera virus, MuV = Mumps virus, PorPV = Porcine rubulavirus, TioPV = Tioman paramyxovirus, hPIV2 = Human parainfluenza virus 2, hMPV = Human metapneumovirus, MPV = Murine pneumonia virus, bRSV = Bovine respiratory syncytial virus, hRSV = Human respiratory syncytial virus, APV = Avian Pneumovirus, ThkPV-1 = Tuhoko virus 1, ThkPV-2 = Tuhoko virus 2, ThkPV-3 = Tuhoko virus 3.

doi:10.1371/journal.pone.0115588.g001

The within-host BelPV distribution tended to be organ-specific. BelPV seemed to be restricted to the heart and liver. In contrast, JPV has been isolated from blood, lung, liver, kidney and spleen of experimentally infected laboratory mice [6] but not in heart. The BelPV distribution for the heart and liver, together with the high viral load in heart tissue, could suggest that this virus is likely to be present in the bloodstream and might thus be transmitted during aggressive contacts between bats, or by blood-sucking vectors. Nevertheless, viremia was

Table 2. Results of real-time PCR on organs from *Coleura afra* individuals.

Samples ID	Heart	Liver	Spleen	Kidney	Lung	Intestine	Brain
GB09 408	ND	41	ND	ND	ND	ND	Undet.
GB09 415	Undet.	40	ND	Undet.	Undet.	ND	ND
GB09 450	Undet.	40	ND	Undet.	ND	ND	ND
GB09 478	28	36	ND	Undet.	Undet.	Undet.	ND
GB09 482	ND	40	ND	Undet.	Undet.	Undet.	ND
GB09 516	Undet.	39	ND	Undet.	Undet.	ND	ND
GB10 239	42	Undet.	Undet.	Undet.	Undet.	Undet.	Undet.
GB10 244	36	Undet.	Undet.	Undet.	Undet.	Undet.	Undet.
GB10 262	41	Undet.	ND	Undet.	Undet.	Undet.	Undet.
GB10 265	44	Undet.	ND	ND	ND	ND	ND
GB10 271	38	Undet.	Undet.	Undet.	Undet.	Undet.	Undet.
GB10 289	40	Undet.	Undet.	Undet.	Undet.	Undet.	Undet.
GB10 301	36	Undet.	ND	Undet.	Undet.	Undet.	Undet.
GB10 259	42	Undet.	Undet.	Undet.	Undet.	Undet.	Undet.

Numbers indicate the cycle threshold (Ct). ND, not done because of missing samples. Undet., Ct undetermined.

doi:10.1371/journal.pone.0115588.t002




Fig. 2. Virus distribution in organs from *Coleura afra* individuals. Virus distribution is shown in terms of Ct values on the y-axis for each bat organ tested (x-axis). n represents the number of organs available for the study.

doi:10.1371/journal.pone.0115588.g002

not proven. BelPV RNA was not searched from blood because in these small species of bats blood was difficult to collect in the field.

We detected BelPV only in *Coleura afra* and not in other bat species sharing the same roosting sites and living in very close proximity in the two caves sampled. However, it has been shown that bats of different species occupying the same roosting sites can share the same viruses. Marburg virus had been detected in *Rousettus aegyptiacus* and *Hipposideros* sp. bats living in Kitaka cave in Uganda [23] and *Miniopterus inflatus* and *Rousettus aegyptiacus* bats caught in Goroubwa mine in the Democratic Republic of the Congo [24]. These bat species are known to live in close proximity. Thus, virus transmission between different bat species is possible [25]. Thus, we can speculate that the failure to detect BelPV in other bat species sharing the same caves would suggest that this virus has strong host specificity for *C. afra*, as well as restricted intraspecies transmission. Henipaviruses occur naturally in fruit bats belonging to the genus *Pteropus* [26], and this also appears to be true of severe acute respiratory syndrome-like coronaviruses in *Rhinolophus* bats [27, 11].

In view of our data we can assume that BelPV might have pathogenic potential for its host *C. afra*. Indeed, high viral load was detected in the heart of the diseased bat, and the lesions were consistent with those reported in wild rodents and mice experimentally infected by JPV [6].

Although BelPV RNA was also detected in asymptomatic bats, pathogenicity may appear in long term under some immunological and/or ecological conditions. Indeed, virus must not induce pathology to persist or adapt within its reservoir host. Many authors suggested that persistence in the absence of pathology or disease appears to be a common characteristic of bat viruses in their natural host population [28, 29]. However, a severe immunodepression for instance, may increase the risk of infection with opportunistic pathogens. Under some environmental conditions (cool environments for example), some avirulent pathogens, such as *Geomyces destructans*, causative agent of white-nose syndrome, may become pathogenic in hibernating bats in North America [30, 31]. Nevertheless, infection by BelPV may be mild for bats and thus the pathology observed not directly related. Otherwise, it may also be that this animal had an underlying disease or infection with a different pathogen. Even in this case, we might not draw any conclusions neither establish a link with lesions seen. Therefore, the pathogenicity of the BelPV should be demonstrated by experimental animal infection. Otherwise, viral antigens or RNAs should be detected histologically in the lesions of naturally-infected bats. However, the unavailability of biological tissues from the diseased bat failed to perform these analyzes. Consequently, other captures of *Coleura afra* species are considered in order to find BelPV again for further studies (pathogenicity to its host, isolation and complete genome characterization). However, *Coleura afra* is a migratory species living in colonies of several hundred individuals. In Gabon, this species, which has been recently described, is not present all year round in the caves of the north-east of the country, making the studies on this species difficult and thereof partly explaining the lack of virological studies.

Some viruses appear to cause clinical disease in wild-living bats; these include lyssaviruses and an ebola-like filovirus named Lloviu virus [32, 33, 34].

Bats are the natural reservoirs for many viruses, including emerging zoonotic viruses such as SARS-CoV [27], Hendra and Nipah viruses [26, 35], Ebola virus [36], Marburg virus [23, 37], rabies virus and other Lyssaviruses [11]. In general, humans are infected through an intermediate amplifying host such as palm civets for SARS-CoV, horses for Hendra virus and pigs for Nipah virus [38]. However, in humans Nipah virus outbreaks linked to bats exposure have been reported [39]. It remains to be shown whether the BelPV reported here presents a zoonotic risk. Nonetheless, like most RNA viruses, for example coronaviruses, characterized by high mutation and/or recombination rates [40], PVs may adapt to novel hosts, including humans. A serological test capable of detecting antibodies to this virus in human populations living in the vicinity of these animals is needed to assess zoonotic potential.

All the blood-sucking arthropods collected from bats, as well as mosquitoes collected in the caves where bat sampling took place, were negative for BelPV, in keeping with the lack of known PV vectors [41]. However, BelPV transmission by blood-sucking vectors within the Gabonese population of *C. afra* cannot be ruled out. Indeed, a haemosporidian parasite (*Polychromophilus*) was found in a blood parasite vector (*Penicillidia fulvida*) in Faucon cave in Gabon in 1977 [42] and also in its host *M. inflatus* (greater long-fingered bat) from the same cave in 2010 and 2011 [43]. In addition, the methodology used to collect flying hematophagous insects (based on light traps) possibly introduced a bias by selecting only those attracted by light. Therefore, we can not exclude that additional sampling techniques could increase the number of mosquitoes species or groups known to colonize caves such as sandflies or biting midges. Hence, the natural mode of transmission of this unclassified paramyxovirus in bat populations, through bat-bat aggression for example, remains to be determined.

This association between *C. afra* and BelPV could serve as an interesting model, (i) to evaluate modes of transmission within host populations, (ii) to study host-virus interactions (pathogenesis and host specificity), and (iii) to evaluate the zoonotic risk of a newly identified virus.

Further studies of *C. afra* populations and a broader diversity of arthropod vectors, spanning larger areas and time scales, are needed to confirm this apparent host-virus specificity, and to determine the modes of BelPV transmission. Further studies are needed to characterize complete BelPV genome and demonstrate the pathogenicity of this virus for its host *Coleura afra*.

Acknowledgments

We thank all the persons involved in sample collections, and especially André Délicat, Peggy Motsch, Dieudonné Nkogué, Tabea Binger, Peter Vallo, and Nil Rahola. We acknowledge Heidi Lançon for improving the English. CIRMF is supported by the Government of Gabon, Total-Fina-Elf Gabon, and Ministère de la Coopération Française.

Author Contributions

Conceived and designed the experiments: GDM MB EML. Performed the experiments: GDM JON NN. Analyzed the data: GDM JFD. Contributed reagents/materials/analysis tools: GDM MB CD CP JFD EML. Wrote the paper: GDM CP EML SM.

References

1. Lamb RA, Kolakofsky D (2001) *Paramyxoviridae*: the viruses and their replication. In: Knipe DM, Howley PM, Griffin DE, Lamb RA, Martin MA, Roizman B, Straus SE (Eds.), *Fields Virology*, 4th ed. Lippincott Williams and Wilkins, Philadelphia; pp. 1305–1340.
2. Tikasingh ES, Jonkers AH, Spence L, Aitken TH (1966) Nariva virus, a hitherto undescribed agent isolated from the Trinidadian rat, *Zygodontomys b. brevicauda* (J.A. Allen and Chapman). *Am J Trop Med Hyg* 15: 235–238.
3. Campbell RW, Carley JG, Doherty R, Domrow R, Filippich C, et al. (1977) Mossman virus, a paramyxovirus of rodents isolated in Queensland. *Search* 8: 435–436.
4. Li Z, Yu M, Zhang H, Magoffin DE, Jack PJM, et al. (2006) Beilong virus, a novel paramyxovirus with the largest genome of non-segmented negative-stranded RNA viruses. *Virology* 346: 219–228.
5. Jun MH (1976) Studies on a virus isolated from wild mice (*Mus musculus*). M. Sc. Thesis. James Cook University, Townsville, Queensland.
6. Jun MH, Karabatsos N, Johnson RH (1977) A new mouse paramyxovirus (*J virus*). *Aust J Exp Biol Med Sci* 55: 645–647.
7. Tidona CA, Kurz HW, Gelderblom HR, Darai G (1999) Isolation and molecular characterization of a novel cytopathogenic paramyxovirus from tree shrews. *Virology* 258: 425–434.
8. Woo PC, Lau SK, Wong BH, Wong AY, Poon RW, et al. (2011) Complete genome sequence of a novel paramyxovirus, *Tailam virus*, discovered in Sikkim rats. *J virol* 85(24): 13473–4. doi: 10.1128/JVI.06356-11.
9. Renshaw RW, Glaser AL, Vancampen H, Weiland F, Dubovi EJ (2000) Identification and phylogenetic comparison of *Salem virus*, a novel paramyxovirus of horses. *Virology* 270: 417–429.
10. Plowright RK, Field HE, Smith C, Divljan A, Palmer C, et al. (2008) Reproduction and nutritional stress are risk factors for Hendra virus infection in little red flying foxes (*Pteropus scapulatus*). *Proc R Soc B* 275: 861–869 (doi:10.1098/rspb.2007.1260)
11. Drexler JF, Corman VM, Müller MA, Maganga GD, Vallo P, et al. (2012) Bats host major mammalian paramyxoviruses. *Nat Commun* 3: 796.
12. Fraser GC, Hooper PT, Lunt RA, Gould AR, Gleeson LJ, et al. (1996) Encephalitis caused by a lyssavirus in fruit bats in Australia. *Emerg Infect Dis* 2: 327–331.
13. Johnson N, Phillpotts R, Fooks AR (2006) Airborne transmission of lyssaviruses. *J Med Microbiol* 55: 785–79
14. Baer GM, Smith JS (1991) Rabies in nonhematophagous bats. In “The Natural History of Rabies”, (G. M Baer, , ed.), CRC Press, Boca Raton, FL, USA, pp.341–366.
15. Gannon W, Sikes R (2007) Guidelines of the American Society of Mammalogists for the use of wild mammals in research. *Journal of Mammalogy*, 88: 809–823.
16. Kunz TH, Parsons S (2009) Ecological and behavioral methods for the study of bats. 2nd ed. Kunz TH, Parsons S, , editors Baltimore, Maryland: Johns Hopkins University Press.
17. Tong S, Chern SW, Li Y, Pallansch MA, Anderson LJ (2008) Sensitive and broadly reactive reverse transcription-PCR assays to detect novel paramyxoviruses. *J Clin Microbiol* 46: 2652–2658.
18. Towner JS, Pourrut X, Albarino CG, Nkogue CN, Bird BH, et al. (2007) Marburg virus infection detected in a common African bat. *PLoS ONE* 2: e764.

19. Towner JS, Sealy TK, Khristova ML, Albarino CG, Conlan S, et al. (2008) Newly discovered ebola virus associated with hemorrhagic Fever outbreak in Uganda. *PLoS Pathog* 4: e1000212.
20. Altschul S, Gish W, Miller W, Myers E, Lipman D (1990) "Basic local alignment search tool". *Journal of Molecular Biology* 215 (3): 403–410. doi:10.1016/S0022-2836(05)80360-2
21. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, et al. (2011) MEGA5: Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance and Maximum Parsimony Methods. *Molecular Biology and Evolution* 28: 2731–2739.
22. Huelsenbeck JP, Ronquist F (2001) MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* 17: 754–5.
23. Towner JS, Amman BR, Sealy TK, Carroll SAR, Comer JA, et al. (2009) Isolation of Genetically Diverse Marburg Viruses from Egyptian Fruit Bats. *PLoS Pathog* 5: e1000536. doi:10.1371/journal.ppat.1000536
24. Swanepoel R, Smit SB, Rollin PE, Formenty P, Leman PA, et al. (2007) Studies of Reservoir Hosts for Marburg Virus. *Emerg Infect Dis* 13: 1847–51. (doi: 10.3201/eid1312.071115).
25. Chu DK, Peiris JS, Chen H, Guan Y, Poon LL (2008) Genomic characterizations of bat coronaviruses (1A, 1B and HKU8) and evidence for co-infections in *Miniopterus* bats. *J Gen Virol* 89(5): 1282–1287.
26. Halpin K, Young PL, Field HE, Mackenzie JS (2000) Isolation of Hendra virus from pteropid bats: a natural reservoir of Hendra virus. *J Gen Virol* 81: 1927–1932.
27. Li W, Shi Z, Yu M, Ren W, Smith C, et al. (2005) Bats are natural reservoirs of SARS-like coronaviruses. *Science* 310: 676–9.
28. Calisher CH, Childs JE, Field HE, Holmes KV, Schountz T (2006) Bats: important reservoir hosts of emerging viruses. *Clin Microbiol Rev* 19: 531–45.
29. Chu DK, Poon LL, Guan Y, Peiris JS (2008) Novel astroviruses in insectivorous bats. *J Virol* 82: 9107–9114.
30. Blehert DS, Hicks AC, Behr MJ, Meteyer CU, Berlowski-Zier BM, et al. (2009) Bat white-nose syndrome: an emerging fungal pathogen? *Science* 323: 227. doi: 10.1126/science.1163874.
31. Warnecke L, Turner JM, Bollinger TK, Lorch JM, Misra V, et al. (2012) Inoculation of bats with European *Geomyces destructans* supports the novel pathogen hypothesis for the origin of white-nose syndrome. *Proc Natl Acad Sci USA* 109: 6999–7003. doi: 10.1073/pnas.1200374109.
32. McColl KA, Chamberlain T, Lunt RA, Newberry KM, Middleton D, et al. (2002) Pathogenesis studies with Australian bat lyssavirus in grey-headed flying foxes (*Pteropus poliocephalus*). *Aust Vet J* 80: 636–641.
33. Banyard AC, Hayman D, Johnson N, McElhinney L, Fooks AR (2011) Bats and Lyssaviruses. *Advances in Virus Research*, 79. doi: 10.1016/B978-0-12-387040-7.00012-3
34. Negrodo A, Palacios G, Vázquez-Morón S, González F, Dopazo H, et al. (2011) Discovery of an Ebolavirus-Like Filovirus in Europe. *PLoS Pathog* 7(10): e1002304. doi:10.1371/journal.ppat.1002304
35. Johara MY, Field H, Rashdi AM, Morrissy C, Heide Bvd, et al. (2001) Nipah virus infection in bats (Order Chiroptera) in Peninsular Malaysia. *Emerging Infectious Diseases* 7: 439–441.
36. Leroy EM, Kumulungui B, Pourrut X, Rouquet P, Hassanin A, et al. (2005) Fruit bats as reservoirs of Ebola virus. *Nature* 438: 575–576.
37. Maganga GD, Bourgarel M, Ebang Ella G, Drexler JF, Gonzalez JP, et al. (2011) Is Marburg Virus Enzootic in Gabon? *J Infect Dis* 204 (Suppl 3): S800–3.
38. Wong S, Lau S, Woo P, Yuen KY (2007) Bats as a continuing source of emerging infections in humans. *Rev Med Virol* 17: 67–91.
39. Montgomery JM, Hossain MJ, Gurley E, Carroll DS, Croisier A, et al. (2008) Risk factor for Nipah virus encephalitis in Bangladesh. *Emerg Infect Dis* 14: 1526–32.
40. Lau SK, Woo PC, Li KS, Huang Y, Tsoi HW, et al. (2005) Severe acute respiratory syndrome coronavirus-like virus in Chinese horseshoe bats. *Proc Natl Acad Sci USA* 102: 14040–14045.
41. Frank GH (1981) *Paramyxovirus* and *Pneumovirus* diseases of animals and birds: comparative aspects and diagnosis. In Kurstack E, editors: *Comparative diagnosis of viral diseases*, New York, Academic Press, pp. 187–233.

42. Landau I, Chabaud AG, Miltgen F, Baccam D (1980) *Dionisia bunoi* n. g. n. sp., Haemoproteidae parasite du microchiroptère *Hipposideros cyclops* au Gabon. *Ann Parasitol Hum Comp* 55: 271–280.
43. Duval L, Mejean C, Maganga GD, Makanga BK, Mangama Koumba LB et al. (2012) The chiropteran haemosporidian *Polychromophilus melanipherus*: A worldwide species complex restricted to the family *Miniopteridae*. *Infections Genetics and Evolution* 12: 1558–1566.