

# *Wolbachia* Age-Sex-Specific Density in *Aedes albopictus*: A Host Evolutionary Response to Cytoplasmic Incompatibility?

Pablo Tortosa<sup>1,2\*</sup>, Sylvain Charlat<sup>3</sup>, Pierrick Labbé<sup>1</sup>, Jean-Sébastien Dehecq<sup>4</sup>, Hélène Barré<sup>5</sup>, Mylène Weill<sup>1</sup>

**1** Institut des Sciences de l'Evolution, CNRS UMR 5554, Université Montpellier 2, Montpellier, France, **2** Centre de Recherche et de Veille sur les Maladies Emergentes dans l'Océan Indien, Ste Clotilde, France, **3** Laboratoire de Biométrie et Biologie Evolutive, CNRS UMR 5558, Université Claude Bernard Lyon 1, Villeurbanne, France, **4** Direction Régionale des Affaires Sanitaires et Sociales de La Réunion, Saint-Denis, France, **5** Laboratoire Parasites et Ecosystèmes Méditerranéens, Université de Corse, Corte, France

## Abstract

**Background:** *Wolbachia* bacteria have invaded many arthropod species by inducing Cytoplasmic Incompatibility (CI). These symbionts represent fascinating objects of study for evolutionary biologists, but also powerful potential biocontrol agents. Here, we assess the density dynamics of *Wolbachia* infections in males and females of the mosquito *Aedes albopictus*, an important vector of human pathogens, and interpret the results within an evolutionary framework.

**Methodology/Principal Findings:** *Wolbachia* densities were measured in natural populations and in age controlled mosquitoes using quantitative PCR. We show that the density dynamics of the wAlbA *Wolbachia* strain infecting *Aedes albopictus* drastically differ between males and females, with a very rapid decay of infection in males only.

**Conclusions/Significance:** Theory predicts that *Wolbachia* and its hosts should cooperate to improve the transmission of infection to offspring, because only infected eggs are protected from the effects of CI. However, incompatible matings effectively lower the fertility of infected males, so that selection acting on the host genome should tend to reduce the expression of CI in males, for example, by reducing infection density in males before sexual maturation. The rapid decay of one *Wolbachia* infection in *Aedes albopictus* males, but not in females, is consistent with this prediction. We suggest that the commonly observed reduction in CI intensity with male age reflects a similar evolutionary process. Our results also highlight the importance of monitoring infection density dynamics in both males and females to assess the efficiency of *Wolbachia*-based control strategies.

**Citation:** Tortosa P, Charlat S, Labbé P, Dehecq J-S, Barré H, et al. (2010) *Wolbachia* Age-Sex-Specific Density in *Aedes albopictus*: A Host Evolutionary Response to Cytoplasmic Incompatibility? PLoS ONE 5(3): e9700. doi:10.1371/journal.pone.0009700

**Editor:** Adam K. Chippindale, Queens University, Canada

**Received:** December 5, 2009; **Accepted:** February 23, 2010; **Published:** March 16, 2010

**Copyright:** © 2010 Tortosa et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This work was funded by ANR CHIKVENDOM Santé-Environnement (Ministère délégué à la recherche); Région Réunion (<http://www.regionreunion.com/fr/spip/spip.php?page=accueilv2>); and Région Languedoc Roussillon (<http://www.laregion.fr/>). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: pablo.tortosa@ird.fr

## Introduction

*Wolbachia* are maternally inherited bacteria that optimize their own fitness by manipulating the reproduction of their host [1]. Among the reported manipulations, Cytoplasmic Incompatibility (CI) is found in several mosquito species and is by far the best documented. In the simplest CI picture, uninfected females' embryos rapidly die following fertilization by *Wolbachia* infected males' sperm. In contrast, infected females' eggs are "immune" to this lethal effect and develop normally into infected adults, so that the infection frequency tends to increase. Similar embryonic mortality is seen in crosses between males and females harbouring different, incompatible, *Wolbachia* variants [2].

This pattern produces two consequences of interest for the present study. First, it makes *Wolbachia* a very promising tool for pest species control strategies: *Wolbachia* can be used as a sterilizing factor when present in males, and is also a potential powerful

vector to drive transgenes through host populations [3]. Second, the drastically different effects of infection on males and females have interesting evolutionary consequences that should be taken into consideration when envisaging *Wolbachia*-based control strategies. *Wolbachia* acts as a rescue factor when present in infected females' embryos; for this reason, the bacteria and its host's genome are expected to cooperate and optimize transmission to offspring [4]. However, the lethal effect of infected males' sperm expressed in crosses with incompatible females is detrimental to the transmission of the host nuclear genes. For this reason, theory predicts that selection acting on host nuclear genes should tend to prevent this effect, for example by reducing the density of infection in males before they become sexually mature [4,5].

In the present work, we use natural populations and age-controlled specimens to show that one *Wolbachia* strain infecting the mosquito *Aedes albopictus* follows the above-predicted density dynamics. We show that females are always co-infected with

*Wolbachia* strains *wAlbA* and *wAlbB*, with bacterial densities slightly increasing with aging. Males display a drastically different pattern: they are co-infected at emergence but *wAlbA* density quickly decreases towards complete loss with male aging. In addition, we investigate the putative role of a WO bacteriophage, previously described in *Ae. albopictus*, in this pattern. WO can alternate lytic and lysogenic cycles and may therefore control *Wolbachia* densities as shown for the parasitoid wasp *Nasonnia vitripennis* [6]. Our data show that WO density is driven by *wAlbB* and not *wAlbA*, confirming previously published data [7] and suggesting that the involvement of this phage in the *wAlbA* depletion is unlikely.

*Ae. albopictus* is raising much concern after multiple invasive episodes documented over the last 20 years [8] with serious medical consequences such as the recent Chikungunya epidemics in the Indian Ocean [9] and Italy [10]. Any *Wolbachia*-based control strategy targeting this species would require the introduction of a new bacterial strain incompatible with the resident infections. Our results emphasize that such strategies should take into consideration the possibility of sex-dependent *Wolbachia* dynamics, which could impede the expression of CI.

## Results

### Sex-specific infection patterns

Infection status of wild male and female *Ae. albopictus* collected on La Réunion Island was determined by a standard PCR assay [11]. Surprisingly, all females were found bi-infected while roughly half of the males were singly infected with *wAlbB* (Table 1). The prevalence of the *wAlbA* infection differed significantly between males and females (Fisher's exact test,  $p < 0.001$ ). Mosquitoes collected on Madagascar were also genotyped following this procedure and displayed the same sex-specific infection pattern although sample size was too small to give significant results (Fisher's exact test,  $p = 0.1$ , Table 1).

We used a recently developed quantitative PCR approach [7] to further investigate this result. This alternative method pointed out that most of the specimens previously characterised as *wAlbB* mono-infected males were actually co-infected with scarce densities of *wAlbA*, undetectable by standard PCR. In addition, quantitative PCR revealed highly variable *wAlbB/wAlbA* ratios in males, ranging from 1 to 5000. The previously described PCR [11] and q-PCR [12] protocols use a common downstream primer annealing on *wAlbA* and *wAlbB* *wsp* sequences (*i.e.* primer 691R) which we suspect may have been titrated by the overwhelming *wAlbB* bacteria, leading to the observed false negative results. This hypothesis was substantiated by a standard PCR assay using *wAlbA-wsp* specific primers [7] which revealed that 4 out of 5

specimens characterised as negative for *wAlbA* with the previously described PCR primers [11] were actually positive.

Quantification data (Fig. 1) showed that densities of both *wAlbA* and *wAlbB* *Wolbachia* differed significantly among males and females (Generalized Linear Model (GLM):  $density = sex$ , where  $density$ , the response variable, is the result of the qPCR analyses, square-root transformed to ensure the normality of the residuals, and  $sex$  a factorial explanatory variable with two levels, male or female:  $F = 57.9$ ,  $P < 0.001$  for *wAlbA*,  $F = 6.3$  and  $P = 0.019$  for *wAlbB*). This approach thus confirmed that *Wolbachia* density pattern is strongly dependent upon sex in *Ae. albopictus*.

### Age and sex-specific infection patterns

All wild females tested were co-infected. This pattern suggests a highly efficient vertical transmission of both infections, making it unlikely that singly infected males received only the *wAlbB* strain from their mother. In addition, should singly infected embryos be produced, we expect most of those to be eliminated through CI. We therefore conjectured that *wAlbA* may be lost after fertilization specifically in male individuals and tested this hypothesis. Eggs were obtained with an oviposition trap deposited on La Réunion Island from the site where the wild larvae had been previously collected. Eggs were dried for 2 days at room temperature and hatching was subsequently induced by immersion into water. Larvae were reared until emergence and adult specimens from each sex were tested at different ages (0, 5, 11 and 15 days after emergence). Total DNA was prepared and analysed by qPCR. The results, reported on figure 2A, show that all females were co-infected. The pattern was drastically different in males, with *wAlbA* densities being very low at emergence and decreasing with aging until no *wAlbA* could be detected within 5 days after emergence.


Two similar experiments were carried out using mosquitoes collected in Greece and Corsica. Mosquitoes were reared under laboratory conditions and individually tested at different times after emergence. The results showed that the above described age-sex-specific pattern is geographically widespread, although some quantitative variation is seen. In the Greek lines, both *wAlbA* and *wAlbB* were detected in all tested females and males, but old

**Table 1.** Infection status of wild caught *Ae. albopictus* as determined with standard PCR [11].

Collection site	Sex	N	A-B+	A+B+	<i>p</i> -value
La Réunion Island	Males	39	20	19	
	Females	40	0	40	<i>&lt;0.001</i>
Madagascar	Males	3	3	0	
	Females	2	0	2	0.1


A-B+ and A+B+ refer to *wAlbB* singly infected and *wAlbA/wAlbB* co-infected mosquitoes, respectively. Differences between male and female infection patterns were analysed using Fisher's exact test (*p*-values are reported for each sample, in italics when significant).

doi:10.1371/journal.pone.0009700.t001



**Figure 1.** Mean *Wolbachia* density in wild specimens from La Réunion Island. Each DNA was quantified in triplicate and the average density was calculated for each specimen. Mean density in the population is reported for each sex and is depicted in black for *wAlbA* and grey for *wAlbB*. Standard error is calculated on the mean of all average densities (males  $N = 15$ , females  $N = 10$ ).

doi:10.1371/journal.pone.0009700.g001



**Figure 2. Effect of aging on *Wolbachia* and WO prophage densities.** The age is indicated on the x axis as days (D) after emergence, for males and females separately. Plots report average *wAlbA* (black), *wAlbB* (grey) and WO prophage (white) density for each age and sex in samples from (A) La Réunion Island, (B) Greece (\* the very low but non-zero density seen in 5-days-old males from Greece is not visible on the figure (mean =  $3.4 \times 10^{-3}$ ; standard error =  $4.2 \times 10^{-3}$ )), and (C) Corsica. Numbers into brackets indicate the number of individuals analysed. doi:10.1371/journal.pone.0009700.g002

males exhibited scarce levels of *wAlbA* (Fig 2B). For mosquitoes from Corsica, the pattern turned out to be very sharp: all tested males were devoid of *wAlbA*, regardless of age, suggesting the *wAlbA* decrease might take place earlier or faster in this line (Fig. 2C). *Wolbachia wAlbA* and *wAlbB* densities were variable depending on the mosquitoes origin, a result previously described in the field [13].

Data from these three quantitative experiments were analyzed together using a GLM in order to simultaneously explore the effects on *Wolbachia* density of three different variables (*age*, *sex* and

*population*), acting either independently or in interaction. In a first model taking *wAlbA* as the variable of interest, *Wolbachia* density is found significantly affected by *age* in interaction with both the *sex* and *population* factors ( $F=17.8$  and  $P<0.001$ ;  $F=8.64$  and  $P<0.001$ , respectively). Interestingly, similar results are obtained in a second model, taking *wAlbB* as the variable of interest (*sex* and *age*:  $F=5.13$  and  $P=0.001$ ; *sex* and *population*:  $F=9.7$  and  $P<0.001$ ). This analysis thus confirms that the density of *wAlbA* is affected by age and sex in all populations, but further suggests that these variables also affect the *wAlbB* density.

To further investigate the effect of age and sex on each infection, a direct correlation between density and age was investigated in both sexes by using Spearman correlation tests. Three independent tests corresponding to each population were run sequentially. Multiple testing was taken into account using the Bonferroni sequential procedure according to Hochberg [14]. The *wAlbA* density was significantly positively correlated with age in females from La Réunion Island and Corsica ( $\rho = 0.637$  and  $P = 0.026$ ;  $\rho = 0.93$  and  $P < 0.001$ , respectively) but not in females from Greece. In contrast, the *wAlbA* density was negatively correlated with age in males from La Réunion Island and Greece ( $\rho = -0.89$  and  $P = 0.007$ ;  $\rho = -0.76$  and  $P = 0.004$ , respectively). The test could not be implemented with the Corsican males which are not infected by *wAlbA*. For *wAlbB* density, results were generally less clear: a significant positive correlation was found between age and *wAlbB* density in females from La Réunion Island ( $\rho = 0.86$  and  $P < 0.001$ ) and Corsica ( $\rho = 0.91$  and  $P < 0.001$ ), but none for females from Greece; a positive correlation was found in males from Corsica ( $\rho = 0.69$  and  $P = 0.013$ ) and from La Réunion Island ( $\rho = 0.85$  and  $P = 0.016$ ; not significant after Bonferroni correction), but the correlation was negative in males from Greece ( $\rho = -0.81$  and  $P = 0.001$ ).

Overall, our data show that both *Wolbachia* types increase with aging in females. In contrast, in males, the *wAlbA* density strongly decreases with age, towards complete loss in males from La Réunion Island while males from Corsica are depleted of this *Wolbachia* at emergence. *wAlbB* density correlates either positively or negatively with age depending on the source sample.

### Assessing WO phage involvement in infection patterns

The density of the WO phage previously described in *Ae. albopictus* was measured in the same mosquitoes by qPCR (Fig. 2) as previously described [7]. The involvement of this phage in the observed pattern was investigated by computing data from the three quantitative experiments in a GLM. This analysis allowed exploring simultaneously the effects of *wAlbA* and *wAlbB* densities on WO density, while controlling for the variables *age*, *sex* and *population*. The model was  $\log(\text{WO}) = \text{sex} * \text{age} * \log(w\text{AlbB}) + \text{sex} * \text{age} * \log(w\text{AlbA}) + \text{population}$ , where WO, *wAlbA* and *wAlbB* are the quantification of the corresponding element (quantitative variables); the values were log-transformed to ensure normality of the residuals; \* indicates that the model considers both the main and interaction effects of the variables. We found a significant effect of *age:wAlbB* interaction ( $F = 4.55$ ,  $P = 0.037$ ), a significant effect of population ( $F = 3.58$ ,  $P = 0.034$ ) and a highly significant effect of *wAlbB* ( $P < 0.001$ ), while all other interactions or main effects were non significant ( $P > 0.05$ ). This analysis thus confirms that the density of this phage is driven only by the density of *wAlbB*, as shown in a previous study [7]. From these results, we can conclude that the WO phage previously described in *Ae. albopictus* is not responsible for the decrease of *wAlbA* density in aging males.

### Discussion

CI causes the death of embryos in crosses between males carrying a given *Wolbachia* strain and females devoid of a compatible infection [15]. Females' offspring infected by a compatible *Wolbachia* are immune to this effect so that the infection frequency tends to increase [2]. The presence of *Wolbachia* is crucial to the survival of infected females' offspring, as only these bacteria can restore the functionality of the sperm "modified" by the paternal *Wolbachia*. Therefore, selection acting on the host nuclear genome is expected to optimise the efficiency

of *Wolbachia* maternal transmission [4,16]. Selection on hosts and symbionts should thus converge toward improved transmission. However, if incompatible crosses do occur, then infected males suffer a fertility reduction. In other words, the expression of CI in a male can be detrimental to the transmission of its nuclear genes, so that selection is expected to suppress the expression of CI, for example by eliminating the infection during development [4,5]. This is the exact pattern we observe for the *wAlbA* infection in *Ae. albopictus*. Male embryos receive the *wAlbA* infection from their doubly infected mothers, and are therefore immune to CI-induced embryonic mortality, but they lose this infection as they become older (Fig. 2). Although other interpretations can be envisaged, including non-adaptive ones, it is tempting to propose that the loss of *wAlbA* in adult *Ae. albopictus* males ultimately results from an evolutionary process selecting nuclear countermeasures to *Wolbachia* manipulation. The decrease of *wAlbA* in males might in turn produce a decline of *wAlbA*-induced CI, which is an assumption of this rational. Notably, age has been found to affect the expression of CI in several systems, sometimes in relation with *Wolbachia* density in testes [17,18,19,20,21,22]. In particular, a marked decrease in CI was previously observed in 10-days-old *wAlbA* mono-infected *Ae. albopictus* males grown in laboratory conditions [23]. The decrease of *wAlbA* densities with aging observed in the present work may be the primary cause of this phenomenon. However, this hypothesis remains to be thoroughly tested since decrease in *Wolbachia* density is not always associated with CI reduction [24,25]. Furthermore, the low *wAlbA* densities quantified here in DNA extracts from entire mosquito bodies might not correlate with densities in testes specifically [26].

Although double infection is virtually fixed among females in modern populations [27], suggesting that the loss of *wAlbA* in males is hardly adaptive nowadays, it is plausible that it was adaptive in the recent past, when the *wAlbA* infection was not fixed. Besides, rare events of *wAlbA* loss in females through imperfect transmission could produce a strong enough pressure to select for the loss of *wAlbA* in adult males.

The *wAlbA* and *wAlbB* infections show markedly different density patterns with regard to age and sex. Under the hypothesis that the decrease of *wAlbA* density in males is adaptive, the non-decrease of *wAlbB* can appear puzzling. The following hypotheses can be proposed to account for this contrast. A first possibility is that the spread of *wAlbB* in *Ae. albopictus* is a very recent event, consistent with the observation that two lab-maintained strains, collected in Mauritius and Samui Islands before 1970, carry only *wAlbA* [27]. A second plausible explanation would relate to differences in the vertical transmission efficiencies of the two strains. If *wAlbB* is more efficiently transmitted from females to their offspring than *wAlbA*, then *wAlbB* infected males suffer less fertility reduction due to CI. In such a case, the selective pressure for a reduction of density in males would be reduced for *wAlbB*. The first hypothesis could be tested using museum specimens, collected in the past, an approach used with success in other systems [28]. Testing the second hypothesis would require accurate measures of transmission efficiencies, controlling for the elimination of uninfected offspring through CI.

In addition to the above, potential interactions between *wAlbA*, *wAlbB* and the host genome would be worth investigating. Indeed, the *wAlbA* depletion might rely in part on the presence of the *wAlbB* strain, through competitive interactions. However, the previous observation that CI expression reduces with aging in *wAlbA* mono-infected males suggests that such interaction cannot be the full explanation for the *wAlbA* depletion. The comparison

of *wAlbA* dynamics in mono-infected vs. bi-infected males, in a controlled genomic background, would allow addressing this issue.

We carried out quantitative PCR to test whether the dramatic decline of *wAlbA* in males is caused by entrance in lytic cycle of a WO phage previously described in *Ae. albopictus*. Our data show that the density of this phage does not increase in males and is driven by *wAlbB* and not *wAlbA* in all three populations. This result confirms similar data obtained on a mosquito population from La Réunion [7,25], and further suggests that the WO-*wAlbB* linkage is not restricted to mosquitoes from La Réunion but appears to be geographically widespread. These data further suggest that this phage is not involved in the *wAlbA* disappearance observed specifically in aging males. Notably, our detection method specifically targets the previously described WO phage from *Ae. albopictus*. Hence the possibility remains that other WO variants, possibly integrated in *wAlbA*, might be involved in the observed pattern.

The use of *Wolbachia* in alternative, environment-friendly vector control strategies is currently investigated in several laboratories [29,30]. CI can be used for mass production of sterilizing males in a derivative of standard sterile insect techniques [31]. The present work emphasizes a problem potentially associated with *Wolbachia*-based vector control strategies. The possibility of *Wolbachia* loss associated with CI decrease in aged males needs to be investigated since it may be detrimental to sterilization strategy. Interestingly, it has been recently reported that *Ae. albopictus* males in La Réunion show an unexpectedly high mean life expectancy, ranging from 16.2 to 24.5 days [32], which would strengthen the impact of density reduction on the effective expression of CI in the field. Therefore, any attempt of producing incompatible males should include CI monitoring over the whole male lifespan. If not appropriately controlled, sterilizing properties of the released males may decrease with aging and therefore compromise the success of such an appealing strategy.

## Methods

### Mosquito rearing

*Wolbachia* genotyping in *Ae. albopictus* natural populations was carried out on mosquitoes collected on La Réunion Island and Madagascar. Mosquitoes from La Réunion Island were collected in the field as larvae, brought to adulthood in the laboratory and immersed into ethanol after emergence. The 5 specimens from Madagascar were collected as adults and directly immersed into ethanol.

Age controlled experiments were carried out as follows: *Ae. albopictus* eggs were collected on Réunion Is. and Corfu Is. with ovitraps made of black-painted soda cans containing water and ovipositing brown paper. Corsican mosquitoes eggs derived from a first generation reproduced in laboratory.

Filter papers were dried for 48h after oviposition and subsequently immersed into fresh water to induce hatching. When larvae reached pupal stage, each bowl was placed inside a

mosquito cage (20cm×20cm×20cm). After emergence, adults were maintained at room temperature with daylight exposure and fed with cotton soaked in honey. Adult specimens were eventually frozen and stored in ethanol.

### Wolbachia genotyping

Total DNA was extracted using a hexadecyltrimethylammonium bromide (CTAB) protocol [33]. Infection status was determined following a previously described method allowing specific amplification of either *Wolbachia* A (81F and 691R primers) or *Wolbachia* B (183F and 691R primers) clades [11]. In addition, the same PCR conditions were used to specifically amplify *wAlbA-wsp* locus using the QAdir1 and QArev2 primers, previously developed for qPCR assays [7].

### Quantitative (q)PCR

*wAlbA*, *wAlbB* and WO prophage were quantified in *Ae. albopictus* following a previously described method [7]. About 2ng of genomic DNA was mixed with primers (0.5 μM) amplifying specifically either *wAlbA*, *wAlbB*, WO or *act* locus, 2 μl of anti-*Taq*-containing master mix and complemented to 20 μl with water. Master mix and anti-*Taq* antibody were used according to Roche LightCycler instructions for SYBR technology [34]. PCR was run for 45 cycles (94°C for 4 s, 65°C for 14 s, and 72°C for 19 s). The pQuantAlb plasmid, containing a single copy of *wAlbA*, *wAlbB*, WO and *actin* template from *Ae. albopictus* nuclear DNA, was serially diluted to build a standard curve with all four *loci* present at an equimolar concentration. Thus same dilutions were amplified with the four specific couples in each qPCR run so that signals could be easily standardized with the nuclear *actin* reference. For each mosquito, quantification measurements were triplicated and mean genome number of *wAlbA*, *wAlbB* and WO was obtained per mosquito nuclear *actin* copy number.

### Statistical analysis

All statistical analyses were computed using the free software R (<http://www.r-project.org/>) using methods, tests and model simplification procedures as described in Crawley [35].

### Acknowledgments

We wish to thank Samuel Huet and Abdoul Ruttee for collecting the natural mosquito population from La Réunion and for sharing their impressive knowledge of mosquitoes breeding sites on the island. We are very grateful to Ilias Kioulos (University of Athens) and John Vontas (University of Crete) for providing Greek samples.

### Author Contributions

Conceived and designed the experiments: MW. Performed the experiments: PT SC MW. Analyzed the data: PT PL MW. Contributed reagents/materials/analysis tools: JSD HB. Wrote the paper: PT.

## References

1. Werren JH, Baldo L, Clark ME (2008) Wolbachia: master manipulators of invertebrate biology. *Nat Rev Microbiol* 6: 741–751.
2. Hoffmann A, Turelli M (1997) Cytoplasmic incompatibility in insects. In: O'Neill S, Hoffmann AA, Werren JH, eds. *Influential Passengers: Inherited Microorganisms and Arthropod Reproduction*. Oxford: Oxford University Press. pp 42–80.
3. Sinkins SP, Gould F (2006) Gene drive systems for insect disease vectors. *Nat Rev Genet* 7: 427–435.
4. Turelli M (1994) Evolution of incompatibility-inducing microbes and their hosts. *Evolution* 48: 1500–1513.
5. Koehncke A, Telschow A, Werren JH, Hammerstein P (2009) Life and death of an influential passenger: Wolbachia and the evolution of CI-modifiers by their hosts. *PLoS One* 4: e4425.
6. Bordenstein SR, Marshall ML, Fry AJ, Kim U, Wernegreen JJ (2006) The tripartite associations between bacteriophage, Wolbachia, and arthropods. *PLoS Pathog* 2: e43.
7. Tortosa P, Courtiol A, Moutailler S, Failloux AB, Weill M (2008) Chikungunya-Wolbachia interplay in *Aedes albopictus*. *Insect Mol Biol* 17: 677–684.
8. Gratz NG (2004) Critical review of the vector status of *Aedes albopictus*. *Med Vet Entomol* 18: 215–227.

9. Delatte H, Dehecq JS, Thiria J, Domerg C, Paupy C, et al. (2008) Geographic Distribution and Developmental Sites of *Aedes albopictus* (Diptera: Culicidae) During a Chikungunya Epidemic Event. *Vector Borne Zoonotic Dis* 8: 25–34.
10. Vazeille M, Jeannin C, Martin E, Schaffner F, Failloux AB (2008) Chikungunya: a risk for Mediterranean countries? *Acta Trop* 105: 200–202.
11. Zhou W, Rousset F, O'Neil S (1998) Phylogeny and PCR-based classification of Wolbachia strains using wsp gene sequences. *Proc Biol Sci* 265: 509–515.
12. Dutton TJ, Sinkins SP (2004) Strain-specific quantification of Wolbachia density in *Aedes albopictus* and effects of larval rearing conditions. *Insect Mol Biol* 13: 317–322.
13. Ahantarig A, Trinachartvanit W, Kittayapong P (2008) Relative Wolbachia density of field-collected *Aedes albopictus* mosquitoes in Thailand. *J Vector Ecol* 33: 173–177.
14. Hochberg Y (1988) A sharper Bonferroni procedure for multiple tests of significance. *Biometrika* 75: 800–802.
15. Dobson SL, Marsland EJ, Rattanadechakul W (2002) Mutualistic Wolbachia infection in *Aedes albopictus*: accelerating cytoplasmic drive. *Genetics* 160: 1087–1094.
16. Engelstadter J, Hammerstein P, Hurst GD (2007) The evolution of endosymbiont density in doubly infected host species. *J Evol Biol* 20: 685–695.
17. Jammongluk W, Kittayapong P, Baisley KJ, O'Neill SL (2000) Wolbachia infection and expression of cytoplasmic incompatibility in *Armigeres subalbatus* (Diptera: Culicidae). *J Med Entomol* 37: 53–57.
18. Noda H, Koizumi Y, Zhang Q, Deng K (2001) Infection density of Wolbachia and incompatibility level in two planthopper species, *Laodelphax striatellus* and *Sogatella furcifera*. *Insect Biochem Mol Biol* 31: 727–737.
19. Turelli M, Hoffmann AA (1995) Cytoplasmic incompatibility in *Drosophila simulans*: dynamics and parameter estimates from natural populations. *Genetics* 140: 1319–1338.
20. Hoffmann AA, Turelli M, Simmons GM (1986) Unidirectional incompatibility between populations of *Drosophila simulans*. *Evolution* 40: 692–701.
21. Binnington KC, Hoffmann AA (1989) Wolbachia-like organisms and cytoplasmic incompatibility in *Drosophila simulans*. *J Invertebr Pathol* 54: 344–352.
22. Bressac C, Rousset F (1993) The reproductive incompatibility system in *Drosophila simulans*: DAPI-staining analysis of the Wolbachia symbionts in sperm cysts. *J Invertebr Pathol* 61: 226–230.
23. Kittayapong P, Mongkalagoon P, Baimai V, O'Neill SL (2002) Host age effect and expression of cytoplasmic incompatibility in field populations of Wolbachia-superinfected *Aedes albopictus*. *Heredity* 88: 270–274.
24. Rasgon JL, Scott TW (2003) Wolbachia and cytoplasmic incompatibility in the California *Culex pipiens* mosquito species complex: parameter estimates and infection dynamics in natural populations. *Genetics* 165: 2029–2038.
25. Riparbelli MG, Giordano R, Callaini G (2007) Effects of Wolbachia on sperm maturation and architecture in *Drosophila simulans* Riverside. *Mech Dev* 124: 699–714.
26. Duron O, Fort P, Weill M (2007) Influence of aging on cytoplasmic incompatibility, sperm modification and Wolbachia density in *Culex pipiens* mosquitoes. *Heredity* 98: 368–374.
27. Armbruster P, Damsky WE, Jr., Giordano R, Birungi J, Munstermann LE, et al. (2003) Infection of New- and Old-World *Aedes albopictus* (Diptera: Culicidae) by the intracellular parasite Wolbachia: implications for host mitochondrial DNA evolution. *J Med Entomol* 40: 356–360.
28. Hornett EA, Charlat S, Wedell N, Jiggins CD, Hurst GD (2009) Rapidly shifting sex ratio across a species range. *Curr Biol* 19: 1628–1631.
29. Jeffery JAL, Nguyen TY, Vu SN, Le TN, Hoffman AA, et al. (2009) Characterizing the *Aedes aegypti* Population in a Vietnamese Village in Preparation for a Wolbachia-Based Mosquito Control Strategy to Eliminate Dengue. *PLoS Negl Trop Dis* 3.
30. McMeniman CJ, Lane RV, Cass BN, Fong AW, Sidhu M, et al. (2009) Stable introduction of a life-shortening Wolbachia infection into the mosquito *Aedes aegypti*. *Science* 323: 141–144.
31. Brelsfoard CL, Sechan Y, Dobson SL (2008) Interspecific hybridization yields strategy for South Pacific filariasis vector elimination. *PLoS Negl Trop Dis* 2: e129.
32. Lacroix R, Delatte H, Hue T, Reiter P (2009) Dispersal and survival of male and female *Aedes albopictus* (Diptera: Culicidae) on Reunion Island. *J Med Entomol* 46: 1117–1124.
33. Rogers SO, Bendich AJ (1988) Extraction of DNA from plant tissues. In: Gelvin SB, Schilperoort RA, eds. *Plant Molecular Biology Manual*. Boston: Kluwer Academic Publisher.
34. Wittwer CT, Herrmann MG, Moss AA, Rasmussen RP (1997) Continuous fluorescence monitoring of rapid cycle DNA amplification. *Biotechniques* 22: 130–131, 134–138.
35. Crawley MJ (2007) *The R book*. Chichester: John Wiley & Sons Ltd.