

The hyperdominant tropical tree *Eschweilera coriacea* (Lecythidaceae) shows higher genetic heterogeneity than sympatric *Eschweilera* species in French Guiana

Myriam Heuertz^{1,*}, Henri Caron^{1,2}, Caroline Scotti-Saintagne³, Pascal Pétronelli², Julien Engel^{4,5}, Niklas Tysklind², Sana Miloudi¹, Fernanda A. Gaiotto⁶, Jérôme Chave⁷, Jean-François Molino⁵, Daniel Sabatier⁵, João Loureiro⁸ & Katharina B. Budde^{1,9}

¹Univ. Bordeaux, INRAE, Biogeco, FR-33610 Cestas, France

²INRAE, Cirad, Ecofog, GF-97310 Kourou, French Guiana

³INRAE, URFM, FR-84914 Avignon, France

⁴International Center for Tropical Botany, Department of Biological Sciences, Florida International University, Miami, FL-33199, USA

⁵Université de Montpellier, IRD, Cirad, CNRS, INRAE, AMAP, FR-34398 Montpellier, France

⁶Universidade Estadual de Santa Cruz, Centro de Biotecnologia e Genética, Ilhéus, BR-45662-901, Bahia, Brazil

⁷Université Paul Sabatier Toulouse, CNRS, EBD, FR-31062, Toulouse, France

⁸University of Coimbra, Centre for Functional Ecology, Department of Life Sciences, PT-3000-456 Coimbra, Portugal

⁹Present address: University of Copenhagen, Forest, Nature and Biomass, Rolighedsvej 23, DK-1958 Frederiksberg C, Denmark

*Corresponding author: myriam.heuertz@inrae.fr

Background and aims – The evolutionary history of Amazonia’s hyperabundant tropical tree species, also known as “hyperdominant” species, remains poorly investigated. We assessed whether the hyperdominant *Eschweilera coriacea* (DC.) S.A.Mori (Lecythidaceae) represents a single genetically cohesive species, and how its genetic constitution relates to other species from the same clade with which it occurs sympatrically in French Guiana.

Methods – We sampled 152 individuals in nine forest sites in French Guiana, representing 11 species of the genus *Eschweilera* all belonging to the Parvifolia clade, with emphasis on *E. coriacea*. Samples were genotyped at four simple sequence repeat (SSR) markers. We delimited gene pools, i.e., genetically coherent putative taxa, using STRUCTURE software and principal component analysis. We compared the genetic assignment of individuals with their morphological species determination and estimated genetic diversity and differentiation for gene pools and species. We also estimated genome size using flow cytometry.

Key results – SSR profiles commonly displayed up to four alleles per genotype, suggesting that the investigated *Eschweilera* species bear a paleopolyploid signature. Flow cytometry suggested that the studied species are diploid with haploid genome sizes of 871–1046 Mbp. We detected five gene pools and observed a good correspondence between morphological and genetic delimitation for *Eschweilera sagotiana* Miers and the undescribed morphospecies *E. sp. 3* (which resembles *E. grandiflora* (Aubl.) Sandwith), and to a lesser extent for *E. decolorans* Sandwith and *E. micrantha* (O.Berg) Miers. *Eschweilera coriacea* was the most genetically diverse species and included individuals assigned to each gene pool.

© 2020 Myriam Heuertz, Henri Caron, Caroline Scotti-Saintagne, Pascal Pétronelli, Julien Engel, Niklas Tysklind, Sana Miloudi, Fernanda A. Gaiotto, Jérôme Chave, Jean-François Molino, Daniel Sabatier, João Loureiro, Katharina B. Budde.

This article is published and distributed in Open Access under the terms of the [Creative Commons Attribution License \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/), which permits use, distribution, and reproduction in any medium, provided the original work (author and source) is properly cited.

Plant Ecology and Evolution is published by Meise Botanic Garden and Royal Botanical Society of Belgium

ISSN: 2032-3913 (print) – 2032-3921 (online)

Conclusions – We found no conclusive evidence for cryptic species within *E. coriacea* in French Guiana. SSRs detected fewer gene pools than expected based on morphology in the Parvifolia clade but discriminated evolutionary relationships better than available plastid markers. A positive trend between demographic abundance of species and allelic richness illustrates that hyperdominants may have a high evolutionary potential. This hypothesis can be tested using more powerful genomic data in combination with tree phenotypic trait variation and characterization of niche breadth, to enhance our understanding of the causes of hyperdominance in Amazonian trees.

Keywords – *Eschweilera*; microsatellites; species delimitation; hyperdominant tropical trees; species complex; cryptic species.

INTRODUCTION

Neotropical rainforests are the world's most diverse terrestrial ecosystems, harbouring 90 000–110 000 species of seed plants, which represents ca. 37% of all seed plants worldwide (Gentry 1982; Barthlott et al. 2007; Antonelli & Sanmartín 2011; Eiserhardt et al. 2017). The relative abundances of plant species and their geographic distribution ranges vary strongly in these forests (Peters et al. 1989; Pitman et al. 2001; Macía & Svenning 2005; ter Steege et al. 2013). In Western Amazonia, small sets of common and abundant species, known as oligarchies, are an ubiquitous feature of tree communities at local, landscape and regional scales (Pitman et al. 2001, 2013; Arellano et al. 2014, 2016). Across lowland Amazonia, a similar pattern is observed, with only 227 species with estimated population sizes of $> 3.7 \times 10^8$ trees (ter Steege et al. 2013). These hyperdominant species represent only 1.4% of the estimated total of 16 000 Amazonian tree species, but as much as 50% of the estimated total number of tree stems (ter Steege et al. 2013). Among the 25 families richest in tree species in Amazonia, Arecaceae, Myristicaceae and Lecythydaceae have the highest proportion of hyperdominant species (ter Steege et al. 2013). However, even for the most abundant of these species, it remains unknown whether their taxonomic definition based on morphological characters includes a single, or several evolutionary lineages. We address this question in *Eschweilera coriacea* (DC.) S.A.Mori (Lecythydaceae), the only tree species that qualified as hyperdominant in all six Amazonian regions – Guiana Shield, northwest, southwest, south, east, and central Amazonia (ter Steege et al. 2013).

The causes of hyperdominance in Amazonia are an active field of research. Considerable overlap has been observed in the species composition of regional Western Amazonian oligarchies and Amazonian hyperdominants, suggesting that the basin-wide pattern arises in part from the combined smaller-scale processes (Pitman et al. 2001, 2013; ter Steege et al. 2013). At the regional scale, oligarchic species have been found to display a wider environmental tolerance, on average, than non-oligarchic species in the same communities (Arellano et al. 2014), which may suggest a high adaptive potential. At larger geographic scales, the strength of the oligarchic pattern was found to decrease, due to the pure effect of area and due to reduced landscape connectivity (Arellano et al. 2016). These results suggest geographic and physiological limits to dominance patterns and are congruent with ter

Steege and colleagues' observation that most hyperdominant species are habitat specialists and are only dominant in certain forest types and in certain regions of the basin (ter Steege et al. 2013). Hyperdominants include many species useful to humans, thus humans may have contributed to shaping hyperdominance patterns (Levis et al. 2017; McMichael et al. 2017). Notwithstanding the reasons for the wide-range dominance patterns in Amazonia, their main implication is that a relatively small suite of tree species accounts for a large proportion of Amazonian ecosystem services, such as water, carbon and nutrient cycling, which should have the potential to greatly simplify research and modelling efforts in forest ecology and biogeochemistry (ter Steege et al. 2013).

The inference of the hyperdominant species status is predicated on a correct botanical identification of tree species in inventory plots. However, correct identification is not trivial because many hyperdominants belong to taxonomically difficult, species-rich genera such as *Eschweilera* (Lecythydaceae), *Protium* (Bursaceae), or *Licania* (Chrysobalanaceae) (Funk et al. 2007) in which several species can co-occur sympatrically (Gonzalez et al. 2009; Baraloto et al. 2012). One major limitation to their correct botanical identification is the scarcity of diagnostic characters on sterile herbarium vouchers, because many closely related tree species are difficult to distinguish based on vegetative characters alone, trees are tall and bear little or no flowers or fruits for most of the year (Mori & Prance 1990; Mori et al. 2001; Goodwin et al. 2015). In some clades, tree species are weakly differentiated due to relatively recent speciation events or occasional hybridization (Gonzalez et al. 2009; Pennington & Lavin 2016; Caron et al. 2019). It is thus reasonable to assume that botanical identification errors may occur in hyperdominants; for instance, a rare taxon may be lumped with the local dominant taxon (Hardy et al. 2017). A related possibility is that some hyperdominants may include cryptic species that are morphologically indistinguishable (based on a limited set of characters), but that represent distinct evolutionary lineages (Cavers et al. 2013; Turchetto-Zolet et al. 2013; Torroba-Balmori et al. 2017). Molecular markers may contribute to species delimitation in species complexes where identification based on morphology is challenging (Duminil & Di Michele 2009).

Here we examine if the hyperdominant and morphologically variable Amazonian tree species *Eschweilera coriacea* (DC.) S.A.Mori represents a single genetically cohesive taxon and whether it presents high genetic variation, an indica-

tor of large population size and adaptive potential (Hoffmann et al. 2017). We test these hypotheses in the Guiana shield, more specifically, in French Guiana. *Eschweilera coriacea* is a common canopy tree species, with a maximum height of up to 37m (Lopes 2007). According to Mori et al. (2017), it belongs to the Parvifolia clade, the most diverse clade in the family Lecythidaceae, which encompasses 63 of the 215 Neotropical species in the family. This clade is nested within the Neotropical Bertholletia clade, and represents ca. half of its 125 species (Huang et al. 2015). Species in the previously defined genus *Eschweilera* (Mori & Prance 1990) fall into three unrelated clades (Integrifolia, Tetrpetala and Parvifolia clades) and evolutionary relationships within clades remain poorly resolved, especially in the Parvifolia clade, either based on morphology or on genetic characters (Huang et al. 2015). Numerous species of the Parvifolia clade commonly occur in sympatry, thus forming complexes of sympatric species: for example, 11 and 15 species in forests in French Guiana (La Fumée Mountain) and Central Amazonia (BDFFP 100 ha plot near Manaus), respectively (Mori 1987; Mori & Lepsch-Cunha 1995; Mori et al. 2001; Huang et al. 2015). These sympatric species share plastid DNA haplotypes (Gonzalez et al. 2009; Caron et al. 2019), suggesting that plastid DNA sequences cannot discriminate species, which can either be due to recent common ancestry and incomplete lineage sorting, or to inter-specific hybridization.

We used microsatellites to delimit gene pools and obtain genetic diversity estimates in individuals morphologically determined as *E. coriacea* from several sites across French Guiana and in sympatric *Eschweilera* individuals determined as belonging to closely related species of the Parvifolia clade. We addressed the following specific questions:

- (1) Does the delimitation of gene pools in the Parvifolia clade coincide with the species determination based on morphology in French Guiana? Which species are best supported by genetic data?
- (2) Is the hyperdominant *E. coriacea* a single cohesive species or a complex of cryptic species? Does it harbour indications of stronger genetic structure, indicating cryptic lineages, and/or higher diversity, a proxy for adaptability, in comparison with other species, and how is the variation distributed geographically?

MATERIAL AND METHODS

Study species and sample collection

The Lecythidaceae family in the New World, known as the Lecythidoideae subfamily, comprises ten genera and 215 described species with a centre of geographic distribution in Amazonia (Huang et al. 2015; Mori et al. 2017). Neotropical Lecythidaceae are sub-canopy to canopy-emergent trees with fibrous bark, and distinctive showy and morphologically diverse flowers with either actinomorphic or zygomorphic androecia (Prance & Mori 1979; Mori & Prance 1990). The Parvifolia clade, to which *E. coriacea* belongs, is characterized by a closed androecium and a double coiled androecial hood. Another synapomorphy of this clade is the presence of a lateral aril on the seed (Huang et al. 2015). The flowers are visited and presumably pollinated by bees, as it is the

case for most Lecythidaceae species (Mori & Prance 1990). Lecythidoideae produce woody fruits; their seeds are gravity dispersed and are found in large numbers directly under the parent trees. Additionally, rodents and primates consume the seeds and might play a role in seed dispersal (Mori & Prance 1990). In Paracou, a lowland forest in French Guiana, sympatric *Eschweilera* species have different preferences for soil water availability, e.g. *E. coriacea* prefers wetter habitats than *E. sagotiana* Miers (Allié et al. 2015), although their ecological tolerance is broad and niches are largely overlapping (S. Schmitt, Univ. Bordeaux, INRAE, France, pers. com.). Flowering occurs synchronously in October–November and leaf trait variation is also largely overlapping among species (S. Schmitt and M. Heuertz, pers. obs.).

We sampled sympatric *Eschweilera* species at nine locations mostly within and sometimes close to forest inventory plots of the GUYAFOR and GUYADIV networks (<http://atd-nmorphospecies.myspecies.info/node/781>) in French Guiana, in North-Eastern South America (table 1, supplementary file 1). Leaf material of 152 individual trees was collected, representing 11 species. Botanical determinations were reached through a continual effort over years in repeated inventories of marked trees, using the vegetative and reproductive characters described in Flora Neotropica (Mori & Prance 1990) and “The Lecythidaceae Pages”, a website based on Flora Neotropica (Mori & Prance 1990) and maintained by S.A. Mori, N.P. Smith, X. Cornejo and G.T. Prance (<http://sweetgum.nybg.org/science/projects/lp/>); a subset of individuals had reproductive characters at the time of sampling: *E. apiculata* (Miers) A.C.Sm., *E. chartaceifolia* S.A.Mori, *E. collina* Eyma, *E. coriacea* (DC.) S.A.Mori, *E. decolorans* Sandwith, *E. micrantha* (O.Berg) Miers, *E. parviflora* Mart. ex DC., *E. pedicellata* (Rich.) S.A.Mori, *E. sagotiana* Miers, *E. wachenheimii* (Benoist) Sandwith and *E. sp. 3*, an as yet undescribed putative species which resembles *E. grandiflora* (Aubl.) Sandwith but has distinctly smaller leaves and can co-occur sympatrically with *E. grandiflora* in the absence of morphologically intermediate individuals (J.-F. Molino and D. Sabatier, pers. obs.). As reference specimen for this putative taxon, we used voucher *Sabatier & Molino 4945*: this specimen has flowers, and duplicates are deposited in CAY, K, NY and P. Special emphasis in the sampling was on *E. coriacea*, represented by 56 individuals (table 1, supplementary file 1). The plant material was dried in paper bags with silica gel immediately after collection in the field. A subset of the individuals collected were vouchered and deposited at the Herbarium IRD de Guyane (CAY) or in the GUYADIV working collection at IRD-Cayenne (supplementary file 1).

Microsatellite isolation and screening

For microsatellite (simple sequence repeat, SSR) isolation, four trees identified as *E. coriacea*, *E. parviflora*, *E. simiorum* (Benoist) Eyma or *E. wachenheimii* were sampled in French Guiana. Total DNA was extracted from dried leaf or cambium materials following a CTAB method (Doyle & Doyle 1987). SSR-enriched libraries were constructed following the protocol of Techen et al. (2010). Briefly, DNA was digested with AluI and HaeIII restriction enzymes, and ligated with SSRLIB3 adapter. Three libraries for each species were built by hybridization to biotinylated oligo repeats groups,

Table 1 – Sample sizes per species and per location of 152 *Eschweilera* samples from the Parvifolia clade included in this study.

Species name abbreviations: Eap, *E. apiculata*; Ech, *E. chartaceifolia*; Ecol, *E. collina*; Ecor, *E. coriacea*; Ed, *E. decolorans*; Em, *E. micrantha*; Epa, *E. parviflora*; Epe, *E. pedicellata*; Esa, *E. sagotiana*, Esp3, *E. sp. 3*, Ew, *E. wachenheimii*.

| Sample site | Lat | Lon | Eap | Ech | Ecol | Ecor | Ed | Em | Epa | Epe | Esa | Esp3 | Ew | Total per site |
|---------------------|------|--------|-----|-----|------|------|----|----|-----|-----|-----|------|----|----------------|
| Acarouany | 4.08 | -52.69 | | | | 15 | | | | | | | | 15 |
| Bafog | 5.55 | -53.88 | | | | 13 | | | 1 | | | | | 14 |
| Beiman | 4.46 | -54.13 | 1 | 1 | 1 | 4 | | 4 | 1 | 7 | 13 | 8 | | 40 |
| Mont Emerillon | 3.26 | -53.19 | | | | 1 | | | | | | | | 1 |
| Montagne Tortue | 4.30 | -52.37 | | | | 4 | | | | | | | | 4 |
| Montagne Trésor | 4.60 | -52.22 | | | | 3 | | | | | | | | 3 |
| Nouragues | 4.09 | -52.70 | | | | 3 | | | | | | | | 3 |
| Paracou | 5.33 | -52.92 | | | 1 | 10 | 18 | | | | 17 | | 1 | 47 |
| Piste de Saint Elie | 5.28 | -53.08 | 2 | 1 | | 3 | 3 | 4 | 3 | 3 | 3 | | 3 | 25 |
| Total per species | | | 3 | 2 | 2 | 56 | 21 | 8 | 5 | 10 | 33 | 8 | 4 | 152 |

[(AC)₁₃+(AACG)₅+(AGG)₈], [(AG)₁₂+(ATC)₈+(AAC)₈] or [(ACAG)₆+(ACCT)₆]. Enriched libraries were PCR amplified and PCR products were purified and pooled in equimolar mixtures for selection of 250–450 bp fractions by electrophoresis on a BluePippin (Sage Science, Beverly, MA, USA) instrument. The microsatellite-enriched DNA libraries were sequenced on an Ion Proton™ system (Thermo Fisher Scientific, Waltham, MA, USA) at the Genome Transcriptome Platform of Bordeaux. Adapters were removed using cutadapt, version 1.2.1 (Martin 2011) and reads were trimmed using Sickle (Joshi & Fass 2011) based on a sliding window approach and a minimum Phred score of 20; reads shorter than 80 bases after trimming were removed. The quality of the remaining reads was checked using FastQC version 0.10.0 (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>). The resulting reads were assembled using the default options of CAP3 (Huang & Madan 1999), and microsatellite loci were identified using the QDD pipeline version 3.0 (Meglécz et al. 2010). Primer pairs were designed using Primer3 version 0.4.0 (Rozen & Skaletsky 2000).

In total, 34000 reads were assembled into 7282 sequences which contained SSRs. Twenty primer pairs were tested on 39 *Lecythis* and *Eschweilera* species. Of these, three amplified reliably and were scorable in *Eschweilera* species of the Parvifolia clade: eschw11740, eschw5831 and eschw64683 (supplementary file 2). We also tested nine primer pairs developed for *Eschweilera ovata* (Cambess.) Mart. ex Miers, a species endemic of the Brazilian Atlantic forest (Santos et al. 2019): a single locus, EO25, amplified reliably and was scorable in *Eschweilera* species from the Parvifolia clade (supplementary file 2). Overall, four loci were used for genotyping: the EO25 locus and the three newly developed loci.

DNA extraction and microsatellite genotyping

Genomic DNA was extracted from all 152 samples following the CTAB protocol (Doyle & Doyle 1987). DNA concentrations were measured using a Nanodrop spectrophotometer

(Thermo Fisher Scientific, Waltham, MA, USA) and samples were diluted to 10 ng/μL. The four SSR markers were PCR amplified in two mixes using 5'-labelled forward primers in combination with the Qiagen Multiplex PCR kit (Qiagen, Hilden, Germany). Reactions contained 1 μL DNA (10 ng/μL), 5 μL of 2x Qiagen Multiplex PCR Master Mix, 3 μL ultrapure water and 1 μL of primer mix (10 μM of each forward and reverse primers). The amplification reaction was performed on a Veriti 96-Well Thermal Cycler (Applied Biosystems, Foster City, Canada) using the following protocol: initial denaturation step at 95°C for 15 min, followed by 30 cycles of 30 s denaturation at 94°C, 90 s annealing at 60°C, 60 s extension at 72°C, and a final extension step at 60°C, for 30 min. Amplified fragments were separated on an automated capillary sequencer (ABI 3700, Applied Biosystems, Foster City, CA, USA). Fragment sizes were determined using ABI GeneMapper v4.1 (Applied Biosystems) in comparison with the GeneScan™ 500 LIZ™ dye size standard (Applied Biosystems), and binned into alleles manually using the frequency distribution of observed fragment sizes. Up to four alleles per genotype were found for three of the loci, we thus suspected our taxa to be tetraploid or paleopolyploids that are diploidized to some extent (Parisod et al. 2010), as had previously been suggested for the related Brazil nut, *Bertholletia excelsa* (Buckley et al. 1988).

Genome size and ploidy

The chromosome base number for Lecythidaceae is $x = 17$ (Mori et al. 2007 and references therein) and several species of the Neotropical *Bertholletia* clade belonging to the non-monophyletic genera *Eschweilera* and *Lecythis* are diploid with $x = 17$ (Kowal et al 1977; Kowal 1989 and references therein); for the diploid *Bertholletia excelsa* Bonpl. haploid genome size was estimated by flow cytometry to 930–940 Mbp (de Barros et al. 2019). To obtain information on genome size for comparison with data from the literature and to detect any possible ploidy differences between the studied

Eschweilera species, we collected leaf or flower bud tissue from nine individuals in the Paracou inventory site representing six *Eschweilera* species (*E. coriacea*, *E. grandiflora*, *E. pedicellata*, *E. sagotiana*, *E. squamata* and *E. wachenheimii*) as well as one related outgroup species, *Lecythis persistens* Sagot, and conserved the tissues in RNAlater (Qiagen) until flow cytometry analysis at the University of Coimbra. Nuclear suspensions were obtained following Galbraith et al. (1983) by chopping RNA-later conserved tissue of the studied species and fresh leaf tissue of *Pisum sativum* ‘Ctirad’ (internal reference standard, $2C = 9.09$ pg; Doležel et al. 1998) in 1 ml of WPB buffer (Loureiro et al. 2007). The nuclear suspension was then filtered using a 50 μm nylon mesh and 50 μg / ml of propidium iodide (PI, Fluka, Buchs, Switzerland) and 50 $\mu\text{g}/\text{ml}$ of RNase (Fluka, Buchs, Switzerland) were added. Samples were analysed in a Partec CyFlow Space flow cytometer (Partec GmbH., Görlitz, Germany; 532 nm green solid-state laser, operating at 30 mW) and results were acquired using Partec FloMax software version 2.4d (Partec GmbH, Münster, Germany).

Genetic diversity

Flow cytometry did not detect any ploidy differences between samples (see Results) and data was congruent with the literature (see Discussion), we thus assumed that allele number variation for all species and loci was due to paleopolyploidy, i.e., to a single ancient genome duplication event common to all the species and loci studied (see Discussion). Since duplicated copies of the analysed SSR loci could not be separated in our data, we used an autotetraploid model for downstream data analysis (Hardy 2016) to account for this probable ancient genome duplication. We used SPAGEDI version 1.5a (Hardy & Vekemans 2002) to estimate multi-locus genetic diversity parameters at the level of a) species, b) gene pools (for gene pool delimitation, see below) and c) sampling sites of *E. coriacea*. The genotypes with a single allele or with four alleles were coded as unambiguous tetraploid homozygotes or heterozygotes, respectively, whereas genotypes with two or three alleles were coded as incomplete genotypes to account for allele copy number ambiguity. We calculated the effective number of alleles N_{ae} , the expected heterozygosity H_E , the observed heterozygosity H_O , the inbreeding coefficient F_{IS} , and we estimated the allelic richness for a standardized sample size of eight gene copies, $A_{R(k=8)}$. Standard errors (SE) for H_E and H_O were calculated based on the standard deviation of estimates from data subsets representing the four possible combinations of three SSRs. Significance levels for F_{IS} to deviate from zero, i.e., deviation from Hardy-Weinberg genotypic proportions, were assessed by 10 000 permutations of gene copies among individuals.

We wished to gain insight into the relationship between the abundance of a species and its genetic diversity, to address the hypothesis that hyperdominant species have increased genetic diversity, a proxy for increased adaptive potential (Hoffmann et al. 2017). We attributed a relative rank of demographic abundance to each investigated species based on raw occurrence data (supplementary file 3) from the GUYAFOR and GUYADIV networks of forest inventory plots, representing a total of 316 plots of 0.12 to 0.16 ha and ca. 143 000 stems inventoried, and we performed lin-

ear regression analysis and a Spearman rank correlation test between allelic richness and ranked abundance in R version 3.5.0 (R Development Core Team 2008).

Gene pool delimitation and genetic structure

Gene pool delimitation in the SSR data was conducted using the Bayesian clustering algorithm implemented in STRUCTURE 2.3.4 (Pritchard et al. 2000), using a tetraploid genotypes model that is robust to allele copy number ambiguity (Falush et al. 2007). The data matrix was converted from the GeneMapper output to a STRUCTURE input file that accounts for allele copy number ambiguity using the R package polysat version 1.7-2 (Clark & Jasieniuk 2011) in R version 3.5.0 (R Development Core Team 2008). In STRUCTURE, ambiguity of allele copy number was accounted for by using RECESSIVEALLELES = 1 and setting the recessive allele code to MISSING, “-9”, as described in the software documentation. To infer individual assignment proportions q in K gene pools, or genetic clusters, we used an admixture model with correlated allele frequencies between clusters, running ten repetitions for each K , with $K = 1$ to $K = 18$, using a burn-in length of 100 000 and a run length of 200 000 MCMC steps. The results were summarized using STRUCTURE Harvester web software version 0.6.94 (Earl & vonHoldt 2012) and the Clumpak server (Kopelman et al. 2015). The number of clusters K that best describes the data structure was determined based on the posterior log likelihood of runs plotted against K and using the delta K approach (Evanno et al. 2005).

These analyses were carried out a) on the full data set of 152 *Eschweilera* samples, b) on a reduced data set of 136 samples including only the six species represented by at least eight individuals: *E. coriacea*, *E. decolorans*, *E. micrantha*, *E. pedicellata*, *E. sagotiana* and *E. sp. 3* and c) on 56 individuals determined as *E. coriacea*. We assessed the congruence of genetic and morphological species delimitation by comparing the assignment of individuals to gene pools at a threshold of STRUCTURE ancestry proportion $q > 0.875$ with their botanical determination. The $q > 0.875$ threshold was chosen because this assignment category is expected to contain genetically pure individuals and second- or later-generation backcrosses (Guichoux et al. 2013), and should thus a priori allow us to identify individuals confidently assigned to their respective gene pools, or candidate genetic species.

Additionally, we conducted a principal component analysis (PCA) using the function *dudi.pca()* implemented in the adegenet package (Jombart 2008) in R version 3.5.0 (R Development Core Team 2008). For this, the genotype matrix was converted to a genind object with presence and absence data of alleles using polysat version 1.7-2 (Clark & Jasieniuk 2011).

Genetic differentiation

Population genetic differentiation was assessed as overall and as pairwise F_{ST} (Weir & Cockerham 1984) between species and between gene pools (individuals assigned with $q > 0.875$) in SPAGEDI. Significance was assessed by 10 000 permutations of individuals among species or gene pools. To correct for multiple testing, a false discovery rate approach

was applied using the R package qvalue version 2.8.0 (Storey 2003) in R version 3.5.0 (R Development Core Team 2008).

We also assessed isolation by distance in *E. coriacea* at the level of sampling locations by regressing pairwise $F_{ST}/(1 - F_{ST})$ values on the logarithm of pairwise spatial distance (Rousset 1997) and at the individual level by regressing pairwise kinship coefficients (Loiselle et al. 1995) between individuals on the logarithm of pairwise spatial distances (Vekemans & Hardy 2004), and using permutation tests in SPAGEDI. For the analysis at the individual level, we added random within-location variation to individual coordinates because exact individual coordinates were unknown.

Table 2 – Genome size estimates obtained through flow cytometry in six *Eschweilera* and one *Lecythis* species.

Tissues were sampled in Paracou, French Guiana. ID, individual tree code in the Paracou inventory plot. Holoploid genome size was measured in pg; it was assumed to correspond to 2C.

| ID | Species | 2C (pg) | 1C (pg) | 1C (Mbp) |
|------------|------------------------|---------|---------|----------|
| P13-2-146 | <i>E. coriacea</i> | 1.94 | 0.97 | 950.8 |
| P1-1-1009 | <i>E. coriacea</i> | 1.78 | 0.89 | 871.3 |
| P1-2-221 | <i>E. grandiflora</i> | 1.93 | 0.96 | 942.9 |
| P13-2-881 | <i>E. pedicellata</i> | 2.14 | 1.07 | 1046.1 |
| P13-4-806 | <i>E. sagotiana</i> | 1.98 | 0.99 | 970.4 |
| P13-2-455 | <i>E. squamata</i> | 1.92 | 0.96 | 940.0 |
| P13-1-2911 | <i>E. wachenheimii</i> | 1.72 | 0.86 | 842.3 |
| P13-1-753 | <i>L. persistens</i> | 1.89 | 0.95 | 924.7 |

Table 3 – Genetic diversity in *Eschweilera* species.

Missing genotypes, unsuccessful amplification; incomplete genotypes, genotypes with two or three alleles recorded; N_{ae} , effective number of alleles; $A_{R(k=8)}$, allelic richness for a sample size of 8 allele copies; H_E , expected heterozygosity (standard error); H_O , observed heterozygosity; F_{IS} , inbreeding coefficient (significance levels: n.s., not significant; *, $P < 0.05$; **, $P < 0.01$, ***, $P < 0.001$); n.a., not available.

| Species | n | missing genotypes (%) | incomplete genotypes (%) | N_{ae} | $A_{R(k=8)}$ | H_E (SE) | H_O (SE) | F_{IS} |
|--------------------------|-----|-----------------------|--------------------------|----------|--------------|---------------|---------------|------------|
| <i>E. apiculata</i> | 3 | 25.0 | 41.7 | 4.89 | n.a. | 0.460 (0.089) | 0.417 (0.084) | 0.254 * |
| <i>E. chartaceifolia</i> | 2 | 25.0 | 25.0 | 1.49 | n.a. | 0.505 (0.069) | 0.375 (0.080) | 0.822 * |
| <i>E. collina</i> | 2 | 12.5 | 37.5 | 2.20 | n.a. | 0.497 (0.057) | 0.375 (0.080) | 0.481 n.s. |
| <i>E. coriacea</i> | 56 | 5.8 | 50.9 | 5.34 | 4.20 | 0.751 (0.024) | 0.559 (0.044) | 0.477 *** |
| <i>E. decolorans</i> | 21 | 3.6 | 46.4 | 3.77 | 3.79 | 0.709 (0.015) | 0.487 (0.061) | 0.506 *** |
| <i>E. micrantha</i> | 8 | 12.5 | 25.0 | 3.14 | 3.28 | 0.621 (0.030) | 0.396 (0.059) | 0.540 *** |
| <i>E. parviflora</i> | 5 | 0.0 | 45.0 | 4.05 | 3.22 | 0.663 (0.035) | 0.500 (0.021) | 0.528 *** |
| <i>E. pedicellata</i> | 10 | 10.0 | 45.0 | 3.19 | 3.32 | 0.650 (0.019) | 0.580 (0.069) | 0.233 ** |
| <i>E. sagotiana</i> | 33 | 2.3 | 46.2 | 2.37 | 2.68 | 0.516 (0.034) | 0.498 (0.060) | 0.262 *** |
| <i>E. sp. 3</i> | 8 | 15.6 | 46.9 | 2.33 | 2.65 | 0.524 (0.033) | 0.504 (0.060) | 0.265 ** |
| <i>E. wachenheimii</i> | 4 | 0.0 | 43.8 | 2.67 | 2.67 | 0.583 (0.026) | 0.563 (0.085) | 0.129 n.s. |
| All species confounded | 152 | 6.3 | 46.2 | 4.87 | 4.24 | 0.772 (0.013) | 0.521 (0.051) | 0.524 *** |

RESULTS

Genome size and ploidy

Eight of the nine samples, all representing leaf tissue, were analysed successfully with flow cytometry and yielded holoploid genome size estimates comprised between 1.72 and 2.14 pg for all species (table 2); no differences in ploidy were detected among the samples. Based on the literature (Kowal et al. 1977; Kowal 1989; de Barros et al. 2019) we assumed holoploid genomes to represent diploids, which led to the estimation of haploid genome sizes of $1C = 842$ to 1047 Mbp for the analysed *Eschweilera* species (table 2).

Genetic diversity and differentiation of *Eschweilera* species

All 152 samples belonging to 11 *Eschweilera* species were successfully genotyped at a minimum of three of the four SSR markers. SSR profiles commonly displayed up to four alleles per genotype, suggesting that the investigated *Eschweilera* species represent diploidized paleopolyploids which retain duplicated copies at some loci. Specifically, genotypes with three or four alleles were found in all species, except in the two species with the lowest sample sizes (*E. collina* and *E. chartaceifolia*, $n = 2$ each). Since alleles from duplicated loci could not be told apart, we analysed the data using a tetraploid framework. A total of 56 alleles were detected across the four loci, with 7 to 21 alleles per locus (supplementary file 2). Expected heterozygosity and allelic richness at the species level were highest in *E. coriacea* ($H_E = 0.751$, $A_{R(k=8)} = 4.1$) and lowest in *E. sp. 3* ($H_E = 0.524$, $A_{R(k=8)} = 2.65$) and *E. sagotiana* ($H_E = 0.516$, $A_{R(k=8)} = 2.68$), considering only species with at least eight individuals assessed

Table 4 – Pairwise genetic differentiation (F_{ST}) between *Eschweilera* species.

Only species with at least eight individuals were considered. Numbers in brackets indicate the number of samples. Significance was assessed using 10000 permutations. Significance levels based on corrected P -values: n.s., not significant; ** $P < 0.01$, *** $P < 0.001$.

| | <i>E. decolorans</i> | <i>E. micrantha</i> | <i>E. pedicellata</i> | <i>E. sagotiana</i> | <i>E. sp. 3</i> |
|----------------------------|----------------------|---------------------|-----------------------|---------------------|-----------------|
| <i>E. coriacea</i> (56) | 0.076 *** | 0.141 *** | 0.078 ** | 0.249 *** | 0.107 ** |
| <i>E. decolorans</i> (21) | | 0.163 ** | 0.039 n.s. | 0.253 *** | 0.224 *** |
| <i>E. micrantha</i> (8) | | | 0.184 ** | 0.266 *** | 0.348 *** |
| <i>E. pedicellata</i> (10) | | | | 0.335 *** | 0.289 *** |
| <i>E. sagotiana</i> (33) | | | | | 0.464 *** |
| <i>E. sp. 3</i> (8) | | | | | |

(table 3). The inbreeding coefficients were positive and deviated significantly from zero in most species (table 3).

A positive trend was identified between demographic abundance of species and allelic richness (fig. 1), with the fitted regression equation $A_{R(k=8)} = 2.842 + 0.085 \times$ (abundance rank), $R^2 = 0.138$, however the relationship was not significant (Spearman signed rank correlation $\rho = 0.452$; p -value = 0.268). *Eschweilera sagotiana* was the most abundant species in French Guiana, however, its allelic richness was lower than expected based on the equation.

Genetic differentiation between species represented by at least eight individuals was significant, with global $F_{ST} = 0.193$ ($P < 0.001$). Pairwise F_{ST} was significant for all pairs of species, except for *E. decolorans* and *E. pedicellata*, and was highest between *E. sagotiana* and *E. sp. 3*, $F_{ST} = 0.464$, $P < 0.001$ (table 4).

Inference of gene pools, their composition and genetic diversity

The Bayesian genetic clustering analysis in STRUCTURE revealed a hierarchical structure for the complete data set.




Figure 1 – Regression of allelic richness on ranked demographic abundance of *Eschweilera* species from the Parvifolia clade in French Guiana.

The delta K statistic indicated optima at $K = 2$ and $K = 5$ genetic clusters and log-likelihood values per run increased as a function of K (fig. 2). There was low inter-run variation in log-likelihood (fig. 2) and Clumpak revealed that clustering solutions converged among runs for $K = 5$ (mean similarity score = 0.982; supplementary file 4). For larger K values, additional structuring emerged although the number of individuals assigned to clusters with high ancestry proportion ($q > 0.875$) decreased (supplementary file 4); we thus considered $K = 5$ as a robust clustering solution in view of the resolution of our data. Assignment of individuals to $K = 5$ gene pools




Figure 2 – Evaluation of the number of genetic clusters using STRUCTURE on 152 *Eschweilera* samples. A. The posterior log-likelihood of data as a function of the number of clusters, K , showed increasing values with increasing K . B. The delta K statistic showed an optimum at $K = 2$ and a secondary peak at $K = 5$.

Table 5 – Assignment *Eschweilera* species to STRUCTURE clusters.

Assignment of number of individuals (percentage) per *Eschweilera* species to the five STRUCTURE clusters C11–C15 (individual assignment threshold $q > 0.875$).

| Species | C11 | C12 | C13 | C14 | C15 | Admixed | n |
|--------------------------|-----------|----------|----------|-----------|----------|-----------|-----|
| <i>E. apiculata</i> | | 2 (66.7) | | | | 1 (33.3) | 3 |
| <i>E. chartaceifolia</i> | | 1 (50.0) | 1 (50.0) | | | | 2 |
| <i>E. collina</i> | | | | 1 (50.0) | 1 (50.0) | | 2 |
| <i>E. coriacea</i> | 3 (5.4) | 9 (16.1) | 7 (12.5) | 2 (3.6) | 6 (10.7) | 29 (51.8) | 56 |
| <i>E. decolorans</i> | 12 (57.1) | 3 (14.3) | 1 (4.8) | 2 (9.5) | | 3 (14.3) | 21 |
| <i>E. micrantha</i> | | | | 1 (12.5) | 4 (50.0) | 3 (37.5) | 8 |
| <i>E. parviflora</i> | | 2 (40.0) | 1 (20.0) | | 1 (20.0) | 1 (20.0) | 5 |
| <i>E. pedicellata</i> | | | 2 (20.0) | | 1 (10.0) | 7 (70.0) | 10 |
| <i>E. sagotiana</i> | 3 (9.1) | | | 24 (72.7) | | 6 (18.2) | 33 |
| <i>E. sp. 3</i> | | | 7 (87.5) | | | 1 (12.5) | 8 |
| <i>E. wachenheimii</i> | | 1 (25.0) | | | 3 (75.0) | | 4 |
| Totals | 18 | 18 | 19 | 30 | 16 | 51 | 152 |

($q > 0.875$) showed that only two gene pools coincided well with the species determination based on morphology: most individuals of *Eschweilera sagotiana* were assigned to C14 (73 %) and most *E. sp. 3* individuals were assigned to C13 (87.5%, table 4, fig. 3). *Eschweilera decolorans* and *E. micrantha* also had 50% or more of their individuals assigned to a single gene pool, C11 and C15, respectively, other individuals being admixed or assigned to other gene pools (table 5, fig. 3). For *E. coriacea* the levels of genetic diversity and structure were particularly high: it had individuals assigned

to all five gene pools, and a high proportion (52%) of its individuals were admixed between two or more gene pools (table 5, fig. 3).

Genetic diversity was highest in C12 with $H_E = 0.757$ and $A_{R(k=8)} = 3.95$ (table 5). In C12, nine out of 18 individuals were determined as *E. coriacea* (table 5). Genetic diversity was lowest in C14 ($H_E = 0.379$; $A_{R(k=8)} = 2.18$) which coincided mostly with *E. sagotiana*. Inbreeding coefficients were positive and significant in four of the five gene pools (table 6).




Figure 3 – STRUCTURE bar plot displaying ancestry proportions of *Eschweilera* individuals (vertical bars) in each of five clusters (coloured segments in each bar). The results are very similar for the complete data set (n = 152, above) and the data including only species with at least eight individuals sampled (n = 136, below).

Table 6 – Genetic diversity estimates for STRUCTURE clusters C11–C15 in the *Eschweilera* data set.

Individuals were assigned to clusters based on an assignment threshold $q > 0.875$; n , number of individuals assigned. N_{ae} , effective number of alleles; $A_{R(k=8)}$, allelic richness for a sample size of 8 allele copies; H_E , expected heterozygosity (SE, standard error); H_O , observed heterozygosity; F_{IS} , inbreeding coefficient. Significance levels: n.s., not significant; *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

| Cl | n | # missing genotypes (%) | # incomplete genotypes (%) | N_{ae} | $A_{R(k=8)}$ | H_E (SE) | H_O (SE) | F_{IS} |
|---------|-----|-------------------------|----------------------------|----------|--------------|---------------|---------------|------------|
| C11 | 18 | 4.2 | 50.0 | 2.41 | 2.65 | 0.509 (0.051) | 0.550 (0.070) | 0.051 n.s. |
| C12 | 18 | 12.5 | 41.7 | 4.79 | 3.95 | 0.757 (0.019) | 0.500 (0.040) | 0.565 *** |
| C13 | 19 | 7.9 | 39.5 | 3.04 | 2.85 | 0.581 (0.035) | 0.446 (0.070) | 0.381 *** |
| C14 | 30 | 1.7 | 39.2 | 1.92 | 2.18 | 0.379 (0.027) | 0.428 (0.060) | 0.096 * |
| C15 | 16 | 6.3 | 50.0 | 3.19 | 3.13 | 0.641 (0.015) | 0.607 (0.041) | 0.288 *** |
| All Cls | 101 | 5.9 | 43.3 | 4.86 | 4.24 | 0.772 (0.015) | 0.492 (0.055) | 0.545 *** |

Table 7 – Pairwise genetic differentiation (F_{ST}) between STRUCTURE clusters C1–C15 in the *Eschweilera* data set.

Individuals were assigned to clusters based on an assignment threshold $q > 0.875$; number of individuals assigned in brackets. Significance levels based on corrected P -values: *** $P < 0.001$.

| | C12 | C13 | C14 | C15 |
|----------|-----------|-----------|-----------|-----------|
| C11 (18) | 0.255 *** | 0.327 *** | 0.528 *** | 0.271 *** |
| C12 (18) | | 0.162 *** | 0.467 *** | 0.166 *** |
| C13 (19) | | | 0.530 *** | 0.294 *** |
| C14 (30) | | | | 0.415 *** |
| C15 (16) | | | | |

Genetic differentiation between gene pools (samples assigned with ancestry proportion $q > 0.875$) was significant, with $F_{ST} = 0.304$ for $K = 2$, and $F_{ST} = 0.390$ for $K = 5$, with pairwise values from $F_{ST} = 0.162$ to $F_{ST} = 0.530$ (all values $P < 0.001$, table 7). The highest level of differentiation was found between C13 and C14, largely represented by species *E. sp. 3* and *E. sagotiana*, in agreement with the differentiation pattern found at the species level.

The STRUCTURE analysis on the six species with at least eight individuals confirmed the hierarchical structure in the data (supplementary file 4), with a clustering result for $K = 5$ that was very similar to the STRUCTURE result on the complete data set (fig. 3, supplementary file 4).

The first three axes of the PCA explained jointly 35.09% of the variance in the data. *Eschweilera sagotiana* formed a relatively cohesive cluster at negative values of PCA1 which only overlapped little with other species (fig. 4). *Eschweilera coriacea* was widely scattered in the PCA space (fig. 4). STRUCTURE clusters overlapped with each other but showed a more segregated distribution in the PCA space than morphologically determined *Eschweilera* species (fig. 4).

Genetic structure and diversity in *Eschweilera coriacea*

The STRUCTURE analysis of $n = 56$ individuals determined as *E. coriacea* revealed an optimum at $K = 2$ (supplementary file 4.3) but differentiation between the two clusters was low and not significant ($F_{ST} = 0.010$ n.s.).

No isolation by distance was detected in *E. coriacea*, neither at the sampling site nor at the individual level (supplementary file 5). Genetic diversity estimates in *E. coriacea*

could only be estimated for three sites (Paracou, Acarouany and Bafog) which were represented by at least 10 samples. Heterozygosity and allelic richness were high and very similar in the three sites, $0.710 \leq H_E \leq 0.775$; $3.67 \leq A_{R(k=8)} \leq 3.98$ (table 8).

DISCUSSION

Can SSR markers resolve evolutionary relationships between *Eschweilera* species from the Parvifolia clade?

Our SSR data on eleven species of *Eschweilera* belonging to the Parvifolia clade displayed a signature of paleopolyploidy at both the species and the gene pool levels. Other species of the Bertholletia clade are diploid (e.g., *Bertholletia excelsa*, *Eschweilera pittieri*, *Eschweilera neei*, *Lecythis minor*, *Lecythis turyrana*; Kowal et al. 1977; Kowal 1989 and references therein) and our haploid genome size estimates were of the same order of magnitude, ca. 1 Gbp, as those of *Bertholletia excelsa*. We therefore suggest a single, common paleopolyploid origin for the locus duplications observed in our data set: these loci appear to have remained in duplicated state whereas other parts of the genome appear to have diploidized (Parisod et al. 2010) probably resulting in a diploid karyotype in most species of the Bertholletia clade. The suggested paleopolyploid origin could coincide with an ancient genome duplication event reported at the base of the Ericales order to which the Lecythidaceae belong (Shi et al. 2010).

The power of our genetic data to discriminate species was relatively poor, considering that the most robust clustering solution revealed only five gene pools in our data, as

Table 8 – Genetic diversity in *Eschweilera coriacea*.

Only sampling sites with at least 10 samples were included in this analysis. Missing genotypes, unsuccessful amplification; incomplete genotypes, genotypes with two or three alleles recorded; N_{ae} , effective number of alleles; $A_{R(k=8)}$, allelic richness for a sample size of 8 allele copies; H_E , expected heterozygosity (SE, standard error); H_O , observed heterozygosity; F_{IS} , inbreeding coefficient (significance levels: ***, $P < 0.001$).

| Site | n | missing genotypes (%) | incomplete genotypes (%) | N_{ae} | $A_{R(k=8)}$ | H_E (SE) | H_O (SE) | F_{IS} |
|-----------|----|-----------------------|--------------------------|----------|--------------|---------------|---------------|-----------|
| Paracou | 10 | 5.0 | 40.0 | 4.65 | 3.98 | 0.775 (0.009) | 0.436 (0.048) | 0.656 *** |
| Acarouany | 15 | 3.3 | 68.3 | 4.39 | 3.67 | 0.710 (0.024) | 0.709 (0.020) | 0.262 *** |
| Bafog | 13 | 5.8 | 44.2 | 5.40 | 3.98 | 0.721 (0.030) | 0.495 (0.060) | 0.501 *** |
| All sites | 56 | 5.8 | 50.9 | 5.34 | 4.20 | 0.751 (0.024) | 0.559 (0.044) | 0.477 *** |

opposed to eleven species determined based on morphology. Marker number and information content of each marker strongly affect the clustering solution (Rosenberg et al. 2005). Despite the coarse resolution, we are confident that the five clusters recovered meaningful evolutionary relationships for the multilocus marker set in our study. Our genetic taxon delimitation represents an improvement over the use of plastid DNA sequences: a reexamination of the data of Gonzalez et al. (2009) and Caron et al. (2019) showed extensive plastid haplotype sharing among species and did not allow to resolve evolutionary relationships in *Eschweilera* in the Parvifolia clade.

Groups of individuals assigned to genetic clusters were genetically more distinct from each other ($F_{ST} = 0.390$) than groups of individuals assigned to species based on morphology ($F_{ST} = 0.193$). This result should be interpreted with caution, considering the discrimination power of the data in combination with the markers and methods employed. The clustering solution of the STRUCTURE software can be biased by the sampling scheme, notably by unbalanced

sampling among populations (Kalinowski 2011), and by the stochastic lineage sorting specific to each marker (Orozco-Wengel et al. 2011). We believe that the first possible bias was limited, given that STRUCTURE results were consistent when using a subset of the data with even representation among species. The paucity of loci did not allow us to address the second possible bias. Ideally, genetic species delimitation should rely on a set of complementary approaches, and conclusions should only be based on a conservative interpretation of congruent results among methods (Carstens et al. 2013). According to Carstens et al. (2013), “in most contexts it is better to fail to delimit species than it is to falsely delimit entities that do not represent actual evolutionary lineages”. In our case, STRUCTURE, PCA and differentiation-based methods yielded congruent results for the delimitation of genetic clusters with our marker set.

Our data delimited gene pools that largely coincided with groups of samples morphologically determined as *E. sp. 3* (Cl3) or *E. sagotiana* (Cl4) and, to a lesser extent, determined as *E. decolorans* (Cl1) or *E. micrantha* (Cl5). The




Figure 4 – Principal component analysis (PCA) of 152 *Eschweilera* individuals genotyped at four SSR loci. Colours and symbols indicate the species determination of samples in plots A and B and the assignment to STRUCTURE clusters (threshold > 0.875 ancestry proportion) in plots C and D.

clusters identified with greatest confidence corresponded to the species pair that displayed the highest level of differentiation, *E. sp. 3* and *E. sagotiana*, suggesting that these two species may represent the most divergent taxa in this sympatric species complex. In the fifth cluster, C12, 50% of individuals corresponded to *E. coriacea*, and there was a markedly lower agreement in the cluster definition vs. its species composition (see also below). The genetic heterogeneity of *E. coriacea* may largely account for such mismatch, although hybridization between closely related taxa in the Parvifolia clade (Caron et al. 2019) may also contribute to hindering taxon delimitation with a limited number of markers.

Reasons for poor congruence between morphological and genetic species delimitation

Morphological and genetic species delimitations represent different abstractions to deal with the complex reality that biodiversity represents. Since both abstractions rely on different species concepts (de Queiroz 2007), they can be congruent, but are not necessarily expected to be. In our study, where marker resolution was low, only two out of eleven species showed a good congruence between genetic and morphological species delimitation. Hybridization and introgression, as suggested in the Parvifolia clade (Gonzalez et al. 2009; Huang et al. 2015; Caron et al. 2019) inevitably leads to low genetic differentiation between species, which causes challenges for genetic delimitation. In rainforest tree species complexes that contain lower numbers of species than considered in our study, and where more powerful markers were used, genetic species delimitation has proven successful, e.g., in *Carapa* (Meliaceae), *Erythrophleum* (Fabaceae) and *Milicia* (Moraceae) (Duminil et al. 2006, 2010; Daïnou et al. 2016). Another reason for poor congruence between morphological and genetic delimitation could be mistaken species identification based on morphology in our data, e.g., three individuals morphologically identified as *Eschweilera sagotiana* were assigned genetically to the cluster mainly identified as *E. decolorans* (C11 in red, in fig. 3).

Choice of molecular approaches for taxon delimitation

We chose to use SSR data and methods based on allele frequency differences for gene pool delimitation in the Parvifolia clade. Given the expected large population sizes in *E. coriacea* (ter Steege et al. 2006, 2013) and the lack of phylogenetic signal using plastid DNA markers in the Parvifolia clade (Gonzalez et al. 2009; Caron et al. 2019), we assumed that evolutionary relationships may be shallow in the clade, which is why we opted for population genetic, rather than phylogenetic approaches for species delimitation (Medrano et al. 2015; Luo et al. 2018). Indeed, coalescent theory shows that the expected time to the most recent common ancestor (TMRCA) for any two homologous sequences is equal to the effective population size, N_e , in numbers of generations, N_e being the size of a (diploid) population evolving according to a Wright-Fisher model with random mating and discrete generations (Nordborg 2001). In the absence of inter-specific gene flow, the performance of phylogenetic methods for species delimitation depends on the ratio of population size to divergence time (Luo et al. 2018): phylogenetic methods

tend to succeed if species divergence time is (substantially) older than the mean TMRCA of gene copies within species (Maddison 1997; Rosenberg & Nordborg 2002). For hyperabundant and widespread tree species that maintain huge population sizes over large areas due to efficient seed and pollen dispersal, this condition is unlikely to be fulfilled. Even if N_e is often much smaller than the census population size N , for example because of variation in reproductive success (Hartl 2000), the TMRCA of gene copies in hyperabundant species is likely to be many million years in the past, and should thus regularly fall within the ancestral species, before the speciation event(s) of interest. The large effective population sizes of common rainforest trees are thus the main reason why phylogenetic trees are often poorly resolved (Pennington & Lavin 2016). An analogous situation is observed in the conifer genus *Pinus*, in which evolutionary relationships were long debated (Willyard et al. 2009), and where it took a set of 21 low-copy nuclear genes with 665 SNPs to obtain a phylogeny with concordant placement of > 75% of the species in the subgenus *Pinaster*, the Mediterranean pines (Grivet et al. 2013). Conversely, SSR markers and population genomic approaches led to successful genetic species delimitation in tropical tree species complexes (Duminil et al. 2012; Daïnou et al. 2016). A prospect for a better phylogenetic resolution and a correct inference of evolutionary relationships is nevertheless offered by the use of multi-locus approaches in a multi-species coalescent framework (Knowles & Carstens 2007; Degnan & Rosenberg 2009; Mirarab et al. 2014).

Genetic constitution and hyperdominance

Our results based on four SSRs suggested that *Eschweilera coriacea* was genetically more diverse and more heterogeneous than related *Eschweilera* species, i.e., *E. sagotiana*, *E. sp. 3*, *E. decolorans* and *E. pedicellata* occurring sympatrically with *E. coriacea* in French Guiana. Although *E. coriacea* individuals were assigned to several genetic clusters when other species were included in the analysis, significant evidence of several genetic clusters was not found when only the morphologically determined *E. coriacea* individuals were analysed. Thus, given the limited power of our SSR markers, our data did not contain robust evidence for *E. coriacea* to be a complex of cryptic species in French Guiana. However, absence of evidence is not evidence of absence! Given the wide distribution of the species, with presence in all six Amazonian regions, and the weak but nevertheless significant genetic structure in the species ($F_{ST} = 0.059$ in French Guiana), spatial and temporal population genetic processes are expected to occur, which make it indeed likely that *E. coriacea* may contain several biological species across Amazonia.

We observed a weak linear trend between allelic richness and ranked abundance of *Eschweilera* species. This relationship is not a robust biological result as it would most likely vary by excluding or adding taxa, sampling sites, loci. This relationship simply serves to illustrate our expectation that the level of genetic diversity of a population, the effectiveness of selection and the strength of genetic drift all depend on the effective population size N_e (Charlesworth 2009; Hoban et al. 2014; Hoffmann et al. 2017). The high diversity and heterogeneity of *E. coriacea* thus suggest that it harbours a larger N_e and a higher adaptive potential than other sympa-

trically occurring species, as expected for a hyperdominant tree species with a census population size as large as 5×10^9 individuals across Amazonia (ter Steege et al. 2013). On the other hand, a high census population size is not necessarily a surrogate for high N_e . A notable outlier of our identified trend is *E. sagotiana* in French Guiana, in which diversity was more reduced, despite it being the most common *Eschweilera* species in our inventories, and despite its large distribution across the Guianas and the Brazilian states of Amapá and Pará (The Lecythidaceae Pages, <http://sweetgum.nybg.org/science/projects/lp/>). The two most common *Eschweilera* species in French Guiana appear thus to have contrasting evolutionary histories. This observation also illustrates that it is difficult to derive any causal relationship when observing a biological pattern, such as that of hyperdominance. For instance, Arellano et al. (2014) observed a wider environmental tolerance in oligarchic than non-oligarchic species, which the authors interpreted as niche breadth causing dominance. But the opposite could also be true: dominant species are more widespread and thus they appear in more habitats, which results in greater observed realized niches, whereas rare species are observed less frequently, thus their niche breadth may be poorly estimated or even biased.

A substantially larger set of genetic markers and a larger and more balanced sampling design should help to shed additional light on the genetic constitution of *E. coriacea*, the characterization of hybridization and introgression in the Parvifolia clade, and the evolutionary history of abundant vs. rare *Eschweilera* species, to understand the genetic underpinnings of hyperdominance in Amazonian tree species.

Conclusions

Our data revealed high genetic diversity and heterogeneity, indicative of high adaptive potential, in the hyperdominant *Eschweilera coriacea* in comparison with other *Eschweilera* species of the Parvifolia clade with which it occurs sympatrically in French Guiana. However, we found no conclusive evidence for cryptic species within *E. coriacea*. Our data set had relatively poor power to delimit species in *Eschweilera* individuals from the Parvifolia clade, although delimitation power was improved in comparison with available plastid DNA markers. Promising avenues for future research on species delimitation and adaptive evolution in species complexes such as *Eschweilera*, Parvifolia clade, will be the combined use of morphological trait data, data on ecological niche characterization and genomic resequencing data using high throughput approaches.

SUPPLEMENTARY FILES

Five supplementary files are associated to this paper:

- (1) Sampling information on 152 *Eschweilera* individuals, STRUCTURE ancestry proportions and SSR data: <https://doi.org/10.5091/plecevo.2020.1565.2053>
- (2) Characteristics of SSR markers used in *Eschweilera*: <https://doi.org/10.5091/plecevo.2020.1565.2055>
- (3) Ranked census data of *Eschweilera* species belonging to the Parvifolia clade in French Guiana: <https://doi.org/10.5091/plecevo.2020.1565.2057>

(4) STRUCTURE analysis in *Eschweilera*: <https://doi.org/10.5091/plecevo.2020.1565.2059>

(5) Isolation by distance analysis in *Eschweilera coriacea*: <https://doi.org/10.5091/plecevo.2020.1565.2061>

ACKNOWLEDGEMENTS

We thank Saintomer Cazal, Valérie Troispoux, Jocelyn Cazal and Patrick Heuret (UMR Ecofog) for help with field and/or laboratory work, and Alain Franc, Stéphanie Mariette and Quentin Jehanne (UMR Biogeco) for stimulating discussions and preliminary work on the study system. We are grateful to our reviewers Gabriel Arellano and one anonymous colleague, as well as the editor André Simões for helpful suggestions on the manuscript. The accuracy of botanical identifications in the GUYADIV and GUYAFOR plot networks is largely due to the availability, quality and continuous improvement of the collections held in the Herbarium IRD de Guyane (CAY). The GUYADIV network is managed by IRD and the GUYAFOR network is managed by CIRAD and CNRS, with the contribution of Office National des Forêts Guyanaises (ONF). This work has benefited from an “Investissement d’Avenir” grant managed by Agence Nationale de la Recherche (CEBA, ref. ANR-10-LABX-25-01). We also acknowledge support from INRA through the Innovation project OUTIREC, and from IRD through the GUYAMAZON programme, LECYTOMICS project. KBB acknowledges an IdEx individual postdoc fellowship from the University of Bordeaux and an AgreenSkills+ mobility grant.

REFERENCES

- Allié E., Pélissier R., Engel J., Petronelli P., Freycon V., Deblauwe V., Soucémariadin L., Weigel J., Baraloto C. (2015) Pervasive local-scale tree-soil habitat association in a tropical forest community. *PLoS One* 10(11): e0141488. <https://doi.org/10.1371/journal.pone.0141488>
- Antonelli A., Sanmartín I. (2011) Why are there so many plant species in the Neotropics? *Taxon* 60(2): 403–414. <https://doi.org/10.1002/tax.602010>
- Arellano G., Cala V., Macía M.J. (2014) Niche breadth of oligarchic species in Amazonian and Andean rain forests. *Journal of Vegetation Science* 25(6): 1355–1366. <https://doi.org/10.1111/jvs.12180>
- Arellano G., Jørgensen P.M., Fuentes A.F., Loza M.I., Torrez V., Macía M.J. (2016) Oligarchic patterns in tropical forests: role of the spatial extent, environmental heterogeneity and diversity. *Journal of Biogeography* 43(3): 616–626. <https://doi.org/10.1111/jbi.12653>
- Baraloto C., Hardy O.J., Paine C.E.T., Dexter K.G., Cruaud C., Dunning L.T., Gonzalez M.-A., Molino J.-F., Sabatier D., Savolainen V., Chave J. (2012) Using functional traits and phylogenetic trees to examine the assembly of tropical tree communities. *Journal of Ecology* 100(3): 690–701. <https://doi.org/10.1111/j.1365-2745.2012.01966.x>
- Barthlott W., Hostert A., Kier G., Küper W., Kreft H., Mutke J., Rafiqpoor M.D., Sommer J.H. (2007) Geographic patterns of vascular plant diversity at continental to global scales (Geographische Muster der Gefäßpflanzenvielfalt im kontinentalen und globalen Maßstab). *Erdkunde* 61(4): 305–315. <https://www.jstor.org/stable/25648042>

- Buckley D.P., O'Malley D.M., Apsit V., Prance G.T., Bawa K.S. (1988) Genetics of Brazil "Nut" (*Berholletia excelsa* Humb. & Bonpl.: Lecythidaceae): I. Genetic variation in natural populations. *Theoretical and Applied Genetics* 76(6): 923–928. <https://doi.org/10.1007/BF00273682>
- Caron H., Molino J.-F., Sabatier D., Léger P., Chaumeil P., Scotti-Saintagne C., Frigério J.-M., Scotti I., Franc A., Petit R.J. (2019) Chloroplast DNA variation in a hyperdiverse tropical tree community. *Ecology and Evolution* 9(8): 4897–4905. <https://doi.org/10.1002/ece3.5096>
- Carstens B.C., Pelletier T.A., Reid N.M., Satler J.D. (2013) How to fail at species delimitation. *Molecular Ecology* 22(17): 4369–4383. <https://doi.org/10.1111/mec.12413>
- Cavers S., Telford A., Arenal Cruz F., Pérez Castañeda A.J., Valencia R., Navarro C., Buonamici A., Lowe A., Vendramin G.G. (2013) Cryptic species and phylogeographical structure in the tree *Cedrela odorata* L. throughout the Neotropics. *Journal of Biogeography* 40(4): 732–746. <https://doi.org/10.1111/jbi.12086>
- Charlesworth B. (2009) Effective population size and patterns of molecular evolution and variation. *Nature Reviews Genetics* 10: 195–210. <https://doi.org/10.1038/nrg2526>
- Clark L.V., Jasieniuk M. (2011) POLYSAT: an R package for polyploid microsatellite analysis. *Molecular Ecology Resources* 11(3): 562–566. <https://doi.org/10.1111/j.1755-0998.2011.02985.x>
- Daïnou K., Blanc-Jolivet C., Degen B., Kimani P., Ndiade-Bourboubou D., Donkpegan A.S.L., Tosso F., Kaymak E., Bourland N., Doucet J.-L., Hardy O.J. (2016) Revealing hidden species diversity in closely related species using nuclear SNPs, SSRs and DNA sequences – a case study in the tree genus *Milicia*. *BMC Evolutionary Biology* 16: 259. <https://doi.org/10.1186/s12862-016-0831-9>
- de Barros L.R.F., de Oliveira Wadt L.H., Mondin M., Pappas Junior G., Rocha R.T., de Castro Rodrigues Pappas M., Kimura R.K., Martins K. (2019) Draft genome assembly of the tropical tree *Bertholletia excelsa* using long-read sequence data. In: XXV IUFRO World Congress, 29 sept - 5 October 2019, Curitiba, PR, Brazil, Abstracts: 318. Pesquisa Florestal Brasileira, Colombo, vol. 39, e201902043, Special issue.
- Degnan J.H., Rosenberg N.A. (2009) Gene tree discordance, phylogenetic inference and the multispecies coalescent. *Trends in Ecology & Evolution* 24(6): 332–340. <https://doi.org/10.1016/j.tree.2009.01.009>
- Doležal J., Greilhuber J., Lucretti S., Meister A., Lysák M.A., Nardi L., Obermayer R. (1998) Plant genome size estimation by flow cytometry: interlaboratory comparison. *Annals of Botany* 82(Suppl. A): 17–26. <https://doi.org/10.1093/oxfordjournals.aob.a010312>
- De Queiroz K. (2007). Species concepts and species delimitation. *Systematic Biology* 56(6): 879–886. <https://doi.org/10.1080/10635150701701083>
- Doyle J., Doyle J.L. (1987) Genomic plant DNA preparation from fresh tissue-CTAB method. *Phytochemical Bulletin* 19(1): 11–15.
- Duminil J., Di Michele M. (2009) Plant species delimitation: a comparison of morphological and molecular markers. *Plant Biosystems* 143(3): 528–542. <https://doi.org/10.1080/11263500902722964>
- Duminil J., Caron H., Scotti I., Casal S.-O., Petit R. J. (2006) Blind population genetics survey of tropical rainforest trees. *Molecular Ecology* 15(12): 3505–3513. <https://doi.org/10.1111/j.1365-294X.2006.03040.x>
- Duminil J., Heuertz M., Doucet J.-L., Bourland N., Cruaud C., Gavyro F., Doumenge C., Navascués M., Hardy O.J. (2010) CpDNA-based species identification and phylogeography: application to African tropical tree species. *Molecular Ecology* 19(24): 5469–5483. <https://doi.org/10.1111/j.1365-294X.2010.04917.x>
- Duminil J., Kenfack D., Viscosi V., Grumiau L., Hardy O.J. (2012) Testing species delimitation in sympatric species complexes: the case of an African tropical tree, *Carapa* spp. (Meliaceae) *Molecular Phylogenetics and Evolution* 62(1): 275–285. <https://doi.org/10.1016/j.ympev.2011.09.020>
- Earl D.A., vonHoldt B.M. (2012) STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conservation Genetics Resources* 4: 359–361. <https://doi.org/10.1007/s12686-011-9548-7>
- Eiserhardt W.L., Couvreur T.L.P., Baker W.J. (2017) Plant phylogeny as a window on the evolution of hyperdiversity in the tropical rainforest biome. *New Phytologist* 214(4): 1408–1422. <https://doi.org/10.1111/nph.14516>
- Evanno G., Regnaut S., Goudet J. (2005) Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular Ecology* 14(8): 2611–2620. <https://doi.org/10.1111/j.1365-294X.2005.02553.x>
- Falush D., Stephens M., Pritchard J.K. (2007) Inference of population structure using multilocus genotype data: dominant markers and null alleles. *Molecular Ecology Notes* 7(4): 574–578. <https://doi.org/10.1111/j.1471-8286.2007.01758.x>
- Funk V., Hollowell T., Berry P., Kelloff C., Alexander S.N. (2007) Checklist of the Plants of the Guiana Shield (Venezuela: Amazonas, Bolívar, Delta Amacuro; Guyana, Surinam, French Guiana) Contributions from the United States National Herbarium, vol. 55. Washington, DC, National Museum of Natural History.
- Galbraith D.W., Harkins K.R., Maddox J.M., Ayres N.M., Sharma D.P., Firoozabady E. (1983) Rapid flow cytometric analysis of the cell cycle in intact plant tissues. *Science* 220(4601): 1049–1051. <https://doi.org/10.1126/science.220.4601.1049>
- Gentry A.H. (1982) Neotropical floristic diversity: phytogeographical connections between Central and South America, Pleistocene climatic fluctuations, or an accident of the Andean orogeny? *Annals of the Missouri Botanical Garden* 69(3): 557–593. <https://doi.org/10.2307/2399084>
- Gonzalez M.A., Baraloto C., Engel J., Mori S.A., Pétronelli P., Riéra B., Chave, J. (2009) Identification of Amazonian trees with DNA barcodes. *PloS One* 4: e7483. <https://doi.org/10.1371/journal.pone.0007483>
- Goodwin Z.A., Harris D.J., Filer D., Wood J.R.I., Scotland R.W. (2015) Widespread mistaken identity in tropical plant collections. *Current Biology* 25(22): R1066–R1067. <https://doi.org/10.1016/j.cub.2015.10.002>
- Grivet D., Climent J., Zabal-Aguirre M., Neale D.B., Vendramin G.G., González-Martínez S.C. (2013) Adaptive evolution of Mediterranean pines. *Molecular Phylogenetics and Evolution* 68(3): 555–566. <https://doi.org/10.1016/j.ympev.2013.03.032>
- Guichoux E., Garnier-Géré P., Lagache L., Lang T., Boury C., Petit R.J. (2013) Outlier loci highlight the direction of introgression in oaks. *Molecular Ecology* 22(2): 450–462. <https://doi.org/10.1111/mec.12125>
- Hardy O.J. (2016) Population genetics of autopolyploids under a mixed mating model and the estimation of selfing rate. *Molecular Ecology Resources* 16(1): 103–117. <https://doi.org/10.1111/1755-0998.12431>
- Hardy O.J., Vekemans X. (2002) Spagedi: a versatile computer program to analyse spatial genetic structure at the individual

- or population levels. *Molecular Ecology Notes* 2(4): 618–620. <https://doi.org/10.1046/j.1471-8286.2002.00305.x>
- Hardy O.J., Dainou K., Donkpegan A., Duminil J., Ewedje E.-E., Ikabanga D.U. (2017) Are we underestimating the number of plant species in the tropics? New insights from population genetics approaches applied on African forest trees. In: Scientific abstracts from the 7th International Barcode of Life Conference. *Genome* 60: 942.
- Hartl D. (2000) A primer of population genetics. 3rd Ed. Sunderland, Sinauer Associates, Inc.
- Hoban S., Arntzen J.A., Bruford M.W., Godoy J.A., Rus Hoelzel A., Segelbacher G., Vilà C., Bertorelle G. (2014) Comparative evaluation of potential indicators and temporal sampling protocols for monitoring genetic erosion. *Evolutionary Applications* 7(9): 984–998. <https://doi.org/10.1111/eva.12197>
- Hoffmann A.A., Sgrò C.M., Kristensen T.N. (2017) Revisiting adaptive potential, population size, and conservation. *Trends in Ecology & Evolution* 32(7): 506–517. <https://doi.org/10.1016/j.tree.2017.03.012>
- Huang X., Madan A. (1999) CAP3: A DNA sequence assembly program. *Genome Research* 9: 868–877. <https://doi.org/10.1101/gr.9.9.868>
- Huang Y.-Y., Mori S.A., Kelly L.M. (2015) Toward a phylogenetic-based generic classification of Neotropical Lecythidaceae - I. Status of *Bertholletia*, *Corythophora*, *Eschweilera* and *Lecythis*. *Phytotaxa* 203: 85–121. <https://doi.org/10.11646/phytotaxa.203.2.1>
- Jombart T. (2008) adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics* 24(11): 1403–1405. <https://doi.org/10.1093/bioinformatics/btn129>
- Joshi N.A., Fass J.N. (2011) Sickle: a sliding-window, adaptive, quality-based trimming tool for FastQ files (Version 1.33) [Software]. Available at <https://github.com/najoshi/sickle> [accessed 27 Jan. 2020].
- Kalinowski S.T. (2011) The computer program STRUCTURE does not reliably identify the main genetic clusters within species: simulations and implications for human population structure. *Heredity* 106: 625–632. <https://doi.org/10.1038/hdy.2010.95>
- Knowles L.L., Carstens B.C. (2007) Delimiting species without monophyletic gene trees. *Systematic Biology* 56(6): 887–895. <https://doi.org/10.1080/10635150701701091>
- Kopelman N.M., Mayzel J., Jakobsson M., Rosenberg N.A., Mayrose I. (2015) Clumpak: a program for identifying clustering modes and packaging population structure inferences across *K*. *Molecular Ecology Resources* 15(5): 1179–1191. <https://doi.org/10.1111/1755-0998.12387>
- Kowal R.R. (1989) Chromosome numbers of *Asteranthos* and the putatively related Lecythidaceae. *Brittonia* 41: 131–135. <https://doi.org/10.2307/2807517>
- Kowal R.R., Mori S.A., Kallunki J.A. (1977) Chromosome numbers of Panamanian Lecythidaceae and their use in subfamilial classification. *Brittonia* 29: 399. <https://doi.org/10.2307/2806482>
- Levis C., Costa F.R.C., Bongers F., Peña-Claros M., Clement C.R., Junqueira A.B., Neves E.G., et al. (2017) Persistent effects of pre-Columbian plant domestication on Amazonian forest composition. *Science* 355(6328): 925–931. <https://doi.org/10.1126/science.aal0157>
- Loiselle B.A., Sork V.L., Nason J., Graham C. (1995) Spatial genetic structure of a tropical understory shrub, *Psychotria officinalis* (Rubiaceae). *American Journal of Botany* 82(11): 1420–1425. <https://doi.org/10.2307/2445869>
- Lopes M.A. (2007) Population structure of *Eschweilera coriacea* (DC.) S. A. Mori in forest fragments in eastern Brazilian Amazonia. *Revista Brasileira de Botânica* 30(3): 509–519. <https://doi.org/10.1590/S0100-84042007000300015>
- Loureiro J., Rodriguez E., Doležel J., Santos C. (2007) Two new nuclear isolation buffers for plant DNA flow cytometry: a test with 37 species. *Annals of Botany* 100(4): 875–888. <https://doi.org/10.1093/aob/mcm152>
- Luo A., Ling C., Ho S.Y.W., Zhu C.-D. (2018) Comparison of methods for molecular species delimitation across a range of speciation scenarios. *Systematic Biology* 67(5): 830–846. <https://doi.org/10.1093/sysbio/syy011>
- Macía M.J., Svenning J.-C. (2005) Oligarchic dominance in western Amazonian plant communities. *Journal of Tropical Ecology* 21(6): 613–626. <https://doi.org/10.1017/S0266467405002579>
- Maddison W.P. (1997) Gene trees in species trees. *Systematic Biology* 46(3): 523–536. <https://doi.org/10.1093/sysbio/46.3.523>
- Mallet J., Besansky N., Hahn M.W. (2016) How reticulated are species? *BioEssays* 38(2): 140–149. <https://doi.org/10.1002/bies.201500149>
- Martin M. (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet Journal* 17(1): 10–12. <https://doi.org/10.14806/ej.17.1.200>
- McMichael C.H., Feeley K.J., Dick C.W., Piperno D.R., Bush M.B. (2017) Comment on “Persistent effects of pre-Columbian plant domestication on Amazonian forest composition.” *Science* 358(6361): eaan8347. <https://doi.org/10.1126/science.aan8347>
- Medrano M., López-Perea E., Herrera C.M. (2015) Population genetics methods applied to a species delimitation problem: endemic trumpet daffodils (*Narcissus* section *Pseudonarcissi*) from the Southern Iberian Peninsula. *International Journal of Plant Sciences* 175(5): 501–517. <https://doi.org/10.1086/675977>
- Megléc E., Costedoat C., Dubut V., Gilles A., Malausa T., Pech N., Martin J.-F. (2010) QDD: a user-friendly program to select microsatellite markers and design primers from large sequencing projects. *Bioinformatics* 26 (3): 403–404. <https://doi.org/10.1093/bioinformatics/btp670>
- Mirarab S., Reaz R., Bayzid M.S., Zimmermann T., Swenson M.S., Warnow T. (2014) ASTRAL: genome-scale coalescent-based species tree estimation. *Bioinformatics* 30(17): i541–i548. <https://doi.org/10.1093/bioinformatics/btu462>
- Mori S.A. (1987) The Lecythidaceae of a lowland Neotropical forest: La Fumée Mountain, French Guiana. *Memoirs of the New York Botanical Garden*, vol. 44. New York, New York Botanical Garden Press.
- Mori S.A., Lepsch-Cunha N. (1995) The Lecythidaceae of a Central Amazonian moist forest. *Memoirs of the New York Botanical Garden*, vol. 75. New York, New York Botanical Garden Press.
- Mori S.A., Prance G.T. (1990) Lecythidaceae, Part 2. The zygomorphic-flowered New World genera (*Couroupita*, *Corythophora*, *Bertholletia*, *Couratari*, *Eschweilera*, & *Lecythis*), with a study of secondary xylem of Neotropical Lecythidaceae by Carl H. de Zeeuw. *Flora Neotropica*, Monograph 21 (II). New York, New York Botanical Garden Press. <https://www.jstor.org/stable/4393724>
- Mori S.A., Becker P., Kincaid D. (2001) Lecythidaceae of a Central Amazonian lowland forest. In: Bierregaard R.O. Jr, Gascon C., Lovejoy T., Mesquita R. (eds) *Lessons from Amazonia: The ecology and conservation of a fragmented forest*: 54–67. New Haven and London, Yale University Press.
- Mori S.A., Tsou C.-H., Wu C.-C., Cronholm B., Anderberg A.A. (2007) Evolution of Lecythidaceae with an emphasis on the

- circumscription of Neotropical genera: information from combined *ndhF* and *trnL-F* sequence data. *Annals of Botany* 94(3): 289–301. <https://doi.org/10.3732/ajb.94.3.289>
- Mori S.A., Kiernan E.A., Smith N.P., Kelley L.M., Huang Y.-Y., Prance G.T., Thiers B. (2017) Observations on the phylogeography of the Lecythidaceae (Brazil nut family). *Phytoneuron* 30: 1–85.
- Nordborg M. (2001) Coalescent theory. In: Balding D.J., Bishop M.J., Cannings C. (eds) *Handbook of Statistical Genetics*: 179–212. Chichester, John Wiley & Sons.
- Orozco-terWengel P., Corander J., Schlötterer C. (2011) Genealogical lineage sorting leads to significant, but incorrect Bayesian multilocus inference of population structure. *Molecular Ecology* 20(6): 1108–1121. <https://doi.org/10.1111/j.1365-294X.2010.04990.x>
- Parisod C., Holderegger R., Brochmann C. (2010) Evolutionary consequences of autopolyploidy. *New Phytologist* 186(1): 5–17. <https://doi.org/10.1111/j.1469-8137.2009.03142.x>
- Pennington R.T., Lavin M. (2016) The contrasting nature of woody plant species in different neotropical forest biomes reflects differences in ecological stability. *New Phytologist* 210(1): 25–37. <https://doi.org/10.1111/nph.13724>
- Peters C.M., Balick M.J., Kahn F., Anderson A.B. (1989) Oligarchic forests of economic plants in Amazonia: Utilization and conservation of an important tropical resource. *Conservation Biology* 3(4): 341–349. <https://www.jstor.org/stable/2386215>
- Pitman N.C.A., Terborgh J.W., Silman M.R., Percy Núñez V., Neill D.A., Cerón C.E., Palacios W.A., Aulestia M. (2001) Dominance and distribution of tree species in Upper Amazonian terra firme forests. *Ecology* 82(8): 2101–2117. [https://doi.org/10.1890/0012-9658\(2001\)082\[2101:DADOTS\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2001)082[2101:DADOTS]2.0.CO;2)
- Pitman N.C.A., Silman M.R., Terborgh J.W. (2013) Oligarchies in Amazonian tree communities: a ten-year review. *Ecography* 36(2): 114–123. <https://doi.org/10.1111/j.1600-0587.2012.00083.x>
- Prance G.T., Mori S.A. (1979) Lecythidaceae—Part 1. The actinomorphic-flowered New World Lecythidaceae (*Asteranthos*, *Gustavia*, *Grias*, *Allantoma*, & *Cariniana*). *Flora Neotropica*, Monograph 21(I). New York, New York Botanical Garden Press. <http://www.jstor.org/stable/4393721>
- Pritchard J.K., Stephens M., Donnelly P. (2000) Inference of population structure using multilocus genotype data. *Genetics* 155(2): 945–959.
- R Development Core Team (2008) R: a language and environment for statistical computing. Vienna, Austria, R Foundation for Statistical Computing. Available at <https://www.r-project.org/> [accessed 27 Jan. 2020].
- Rosenberg N.A., Nordborg M. (2002) Genealogical trees, coalescent theory and the analysis of genetic polymorphisms. *Nature Reviews Genetics* 3: 380–390. <https://doi.org/10.1038/nrg795>
- Rosenberg N.A., Mahajan S., Ramachandran S., Zhao C., Pritchard J.K., Feldman M.W. (2005) Clines, clusters, and the effect of study design on the inference of human population structure. *PLoS Genetics* 1: e70. <https://doi.org/10.1371/journal.pgen.0010070>
- Rousset F. (1997) Genetic differentiation and estimation of gene flow from F-statistics under isolation by distance. *Genetics* 145(4): 1219–1228.
- Rozen S., Skaletsky H. (2000) Primer3 on the WWW for general users and for biologist programmers. *Methods in Molecular Biology* 132: 365–386. <https://doi.org/10.1385/1-59259-192-2:365>
- Santos A.S., Borges D.B., Vivas C.V., Berg C.V.D., Rodrigues P.S., Tarazi R., Gaiotto F.A. (2019) Gene pool sharing and genetic bottleneck effects in subpopulations of *Eschweilera ovata* (Cambess.) Mart. ex Miers (Lecythidaceae) in the Atlantic Forest of southern Bahia, Brazil. *Genetics and Molecular Biology* 42(3): 655–665. <https://doi.org/10.1590/1678-4685-gmb-2018-0140>
- Shi T., Huang H., Barker M.S. (2010) Ancient genome duplications during the evolution of kiwifruit (*Actinidia*) and related Ericales. *Annals of Botany* 106(3): 497–504. <https://doi.org/10.1093/aob/mcq129>
- Storey J.D. (2003) The positive false discovery rate: a Bayesian interpretation and the *q*-value. *The Annals of Statistics* 31(6): 2013–2035. <https://doi.org/10.1214/aos/1074290335>
- Teuchen N., Arias R.S., Glynn N.C., Pan Z., Khan I.A., Scheffler B.E. (2010) Optimized construction of microsatellite-enriched libraries. *Molecular Ecology Resources* 10(3): 508–515. <https://doi.org/10.1111/j.1755-0998.2009.02802.x>
- ter Steege H., Pitman N.C.A., Phillips O.L., Chave J., Sabatier D., Duque A., et al. (2006) Continental-scale patterns of canopy tree composition and function across Amazonia. *Nature* 443: 444–447. <https://doi.org/10.1038/nature05134>
- ter Steege H., Pitman N.C.A., Sabatier D., Baraloto C., Salomão R.P., Guevara J.E., et al. (2013) Hyperdominance in the Amazonian tree flora. *Science* 342(6156): 1243092. <https://doi.org/10.1126/science.1243092>
- Torroba-Balmori P., Budde K.B., Heer K., González-Martínez S.C., Olsson S., Scotti-Saintagne C., Casalis M., Sonké B., Dick C.W., Heuertz M. (2017) Altitudinal gradients, biogeographic history and microhabitat adaptation affect fine-scale spatial genetic structure in African and Neotropical populations of an ancient tropical tree species. *PLoS One* 12: e0182515. <https://doi.org/10.1371/journal.pone.0182515>
- Turchetto-Zolet A.C., Pinheiro F., Salgueiro F., Palma-Silva C. (2013) Phylogeographical patterns shed light on evolutionary process in South America. *Molecular Ecology* 22(5): 1193–1213. <https://doi.org/10.1111/mec.12164>
- Vekemans X., Hardy O.J. (2004) New insights from fine-scale spatial genetic structure analyses in plant populations. *Molecular Ecology* 13(4): 921–935. <https://doi.org/10.1046/j.1365-294X.2004.02076.x>
- Weir B.S., Cockerham C.C. (1984) Estimating *F*-statistics for the analysis of population structure. *Evolution* 38(6): 1358–1370. <https://doi.org/10.1111/j.1558-5646.1984.tb05657.x>
- Willyard A., Cronn R., Liston A. (2009) Reticulate evolution and incomplete lineage sorting among the ponderosa pines. *Molecular Phylogenetics and Evolution* 52(2): 498–511. <https://doi.org/10.1016/j.ympev.2009.02.011>

Communicating Editor: André Simões.

Submission date: 4 Dec. 2018

Acceptance date: 28 Jan. 2020

Publication date: 26 Mar. 2020