

Main Earthquakes in the region

Event	Year	Province and Locality	Intensity MM
1	1692	Salta (Talavera de Esteco)	VIII
2	1826	Tucumán (Trancas)	VIII
3	1844	Salta (Palomitas)	VIII
4	1863	Jujuy (San Salvador de Jujuy)	VII
5	1871	Salta (Orán)	VII
6	1874	Salta (Orán)	VIII
7	1892	Catamarca (Pomán)	VIII
8	1899	Yacuiba (Limite Argentino - Boliviano)	VII
9	1906	Tucumán (Tafí del Valle)	VII
10	1930	Salta (La Poma)	VIII
11	1931	Tucumán (El Naranjo)	VII
12	1948	Salta (Palomitas)	IX
13	1959	Salta (San Andrés)	VIII
14	1966	Catamarca (Belén)	VII
15	1973	Salta (Santa Clara)	VII
16	1974	Salta (Santa Victoria)	VII
17	1974	Salta (Orán)	VII

THE SEISMOTECTONIC SETTING

The region under study is located in a tectonic setting resulting from the ongoing subduction of the Nazca Plate eastward beneath the South American Plate. This process has created major structural features as complex faulting, the Puna Plateau and to the east, the Subandean Ranges.

Considering the existing megastructures in the region, it has been observed that these structures constitute barriers limiting shallows from intermediate earthquakes.

SEISMOGENIC SOURCES

The seismotectonic process is quite complex towards east, especially after the 3,000 m contour. Taking the megastructures into account besides the 3,000 m contour, it is observed that the main earthquakes direction will be SSW to NNE, starting from Catamarca to Yacuiba (at the Argentina-Bolivia border). In this zone the most dangerous earthquakes have occurred.


LOCAL TECTONIC STRESSES

The W-E direction of the maximum horizontal tectonic stresses, which are caused by the collision between both plates, are altered by the local horizontal stresses. Therefore, the final orientation results almost perpendicular to the 3,000 m of the Puna contour.

Some of the focal mechanisms for this zone have been very well determined in previous studies and their representations confirms the above statements. Besides, it is possible to assert that the high Andes are under a tensional process that is transmitted to the puna step by step (e.g. the normal faults near to the 3,000 contour). East of the 3,000 m contour the maximum horizontal tectonic stress caused by the collision between both plates is predominant and therefore the principal stress direction suffers no changes.

CONCLUSIONS

Main shallow earthquakes in this region have occurred close to the 3,000 m contour, which indicates that



MAXIMUM REPORTED INTENSITIES

