

## Intraseasonal Variability - A Critical Component of ENSO ?

Stephen E. ZEBIAK

*Lamont-Doherty Geological Observatory  
Palisades, NY 10964 - U.S.A.*


### ABSTRACT

The impact of intraseasonal variability on ENSO is studied in the context of the Zebiak and Cane (1987) coupled atmosphere-ocean model, and an idealized representation intraseasonal forcing. The effects of the parameterized forcing are examined in both simulation and forecast experiments, with similar results: the intraseasonal variability generally plays a minor role in altering the model behavior. Despite the uncertainties inherent in both the coupled model and the specified forcing, the results clearly suggest that intraseasonal variability is not an essential component of ENSO. At the same time, they present evidence of occasional sensitive periods or states of the coupled system in which intraseasonal forcing (and possibly other forcings) can indeed disrupt the future course of events.

### 1. Introduction

Since the pioneering work of Madden and Julian (1971,1972), the so-called 40-50 day oscillation (also known as the 30-60 day oscillation, or Madden-Julian oscillation) has been the subject of intense study. It is a prominent mode of variability on intraseasonal timescales throughout most of the tropics, with clear signatures in upper and lower level tropospheric zonal wind, and convective activity (Madden and Julian 1971,1972; Lau and Chan 1985,1986a; Knutson and Weickmann 1987; Chen and Murakami 1988). Its significance is further heightened by suggested relationships with a number of other phenomena; among them, fluctuations of the Indian monsoon (Krishnamurti and Subrahmanyam 1982; Murakami et al. 1984,1986; Cadet and Greco 1987), fluctuations of the Southeast Asian monsoon (Chen et al. 1988), westerly wind bursts in the western Pacific (Lau et al. 1988), and the onset of El Niño (Lau and Chan 1986b,1988). The latter two are related inasmuch as westerly wind episodes typically characterize the early stages of El Niño events (Luther et al. 1983). The proposed relationship between 30-60 day activity (hereafter referred to as intraseasonal variability or ISV) and El Niño is the subject of this study. Specifically, we will assess the impact of ISV on the Zebiak and Cane (1987) coupled model in both simulation mode and forecast mode.

One can easily imagine, following the authors cited above, that the energetic ISV forcing, if not an essential element of ENSO, is often influential in the timing of events. On the other hand, the theory emerging from simple coupled models points to an essentially deterministic interannual cycle. If one accepts this theory, then the role of ISV would seem to be secondary; or more correctly, the role of that component of ISV not controlled by the slower, ENSO-mode variability would seem secondary. Whereas one perspective treats El Niño occurrences as individual events, dependent on some triggering mechanism(s), the other views warm (and cold) events as the inevitable consequence of a systematic preconditioning process, and part of an ongoing, self-sustaining cycle. Of course, there is the possibility that both viewpoints are partially correct. A






Fig. 1. Three 48-month realizations of the stochastic ISV model defined by Eq. (1).




Fig. 2. Spectrum of model ISV. Abscissa has been scaled so that zero represents a 40-day period (logarithms are in base 10).

deterministic, but chaotic, ENSO cycle likely would be characterized by periods of lesser and greater susceptibility to various external forcing factors, including ISV. Even if this forcing were not usually sufficient to disrupt the cycle, at particular times or in extreme instances it might well do so. It is the purpose of this paper to determine which of these scenarios obtains in the context of the ZC model.

## 2. A Model of Intraseasonal Forcing

Intraseasonal variability appears naturally in atmospheric models with time-dependent dynamics, and reasonable representations of convective heating (e.g., Lau and Lau, 1986; Chang and Lim, 1988; Hendon, 1988; Sui and Lau, 1989). It does not occur in the simpler, steady-state models such as that of ZC. Thus, for the present purposes it is necessary to specify the intraseasonal variability in some fashion. To do this, several observational results are utilized: (i) the low-level wind signal is dominantly zonal in the equatorial region, (ii) the 30-60 day period band contains most of the power, and (iii) the disturbances are energetic in the west Pacific, but weaker (at the surface) in the eastern Pacific. In the model to be constructed here the spatial structure of the imposed anomalies is fixed in time, and positioned in the western Pacific at the equator. The eastward propagation aspect is not included as it is considered relatively unimportant in terms of the ocean response (especially the remote response to the east of the forcing region). Specifically, the following form is chosen for fluctuations in zonal wind stress:

$$\tau^{(x)}(t) = A \left[ R(t) + 2R(t-\Delta t) + R(t-2\Delta t) \right] \cos(\omega_0 t + t_0) \exp\left[\frac{y}{10^\circ}\right]^2 \exp\left[\frac{x-x_0}{30^\circ}\right]^2, \quad (1)$$

where  $R$  is a normal random variable with zero mean and unit variance, and  $t_0$  represents a uniform random variable on  $(0, 2\pi)$ . This forcing is evaluated at time intervals of  $\Delta t$ , which for the ZC model is 10 days. The parameters  $\omega_0$  and  $x_0$  were taken to be  $2\pi/40\text{days}$  and  $146^\circ\text{E}$ , respectively, and the amplitude  $A$  was set at  $.015 \text{ N/m}^2$ .

Fig. 1 shows three 48-month realizations of this forcing function. Anomalies as large as the climatological mean stress (about  $.05 \text{ N/m}^2$ ) occur frequently; the model forcing is at least as strong, and probably stronger, than what is observed (Madden, personal communication). A spectrum of the forcing (Fig. 2) shows that, like the real ISV, most of the total power resides in the 20-60 day band. Subsequent coupled model runs incorporate this forcing with one exception: the amplitude  $A$  is reduced during simulated warm episodes. We reduce  $A$  in proportion to  $(\text{NINO3})^{1/2}$  for  $\text{NINO3} > 1$ , where  $\text{NINO3}$  is the area averaged SST anomalies for the region ( $90^\circ\text{W}-150^\circ\text{W}, 5^\circ\text{S}-5^\circ\text{N}$ ). The rationale behind this is that convective activity is generally observed to decrease in the west Pacific during El Niño events. The reduction in model forcing is modest (always less than a factor of two), so that the simulated ISV is still energetic by observed standards.

## 3. Model Simulations/Forecasts with Imposed ISV

Two 1000-year simulations were done, one with and one without ISV forcing, for the standard physics case as described in ZC. Such a lengthy simulation reveals a wide range of behavior in the unperturbed coupled model (Fig. 3), including periods of large, regular warm events, periods of more irregular cycles, and even periods of almost no variability. How much can a powerful ISV forcing alter this pattern? Given this range of behavior in the basic model, it is perhaps not surprising to find that there is little effect (Fig. 4). The favored 4 year period, and transitions between regimes of larger and smaller amplitude variability all occur analogously. A more quantitative comparison is shown in Fig. 5. The two time series were divided into segments of 24 years duration, Fourier analyzed (with a Welch window applied to minimize aliasing effects), and then averaged to form a mean spectrum. The two spectra are nearly identical; only




Fig. 3. NINO3 index from a 1000-year run of the coupled model with no ISV forcing.




Fig. 4. As Fig. 3, except with added ISV forcing.




Fig. 5. Spectra of the time series in Figs. 3 and 4, for the period range 1-24 years.




Fig. 6. Observed NINO3 (heavy solid line), and forecasts of NINO3 made with the coupled model alone (light solid line) and with added ISV forcing (dashed lines), starting from July 1970 through October 1970.






Fig. 7. As Fig. 6, except for September 1975 through December 1975.




Fig. 8. As Fig. 6, except for February 1978 through May 1978.




Fig. 9. As Fig. 6, except for September 1986 through December 1986.




Fig. 10. Root mean square separation among all pairs of forecasts starting from the same initial conditions, but with different realizations of ISV (dashed line), compared with the same quantity computed from pairs of forecasts without ISV beginning one month apart (solid line).



will be much larger, and the predictability limit will be reached sooner.

For the present purposes we wish to compare the spreading rates associated with ISV forcing and one-month differences in start time. The former is determined by computing the mean square difference among all pairs of forecasts starting from the same initial conditions, but with different realizations of intraseasonal forcing. Similarly, the latter is determined from the mean square difference between forecasts starting one month apart, and with no ISV (only the overlapping time period enters for this calculation). Both are shown in Fig. 10. The results confirm what was suggested in the sets of individual forecasts: the ISV forcing has less overall effect on the forecasts than does moving the start time by one month.

#### 4. Discussion

The question of intraseasonal variability and its impact on ENSO has been addressed in the context of the Zebiak and Cane (1987) coupled model and an idealized representation of ISV. From several different analyses, the results are unambiguous: ISV has only a marginal effect on ENSO. Lengthy integrations of the model with and without ISV yield nearly identical statistics. The majority of individual forecast experiments show little spreading over a two year period as a result of this forcing. Predictability calculations show that, even relative to the differences among forecasts starting one month apart, the ISV influence is small. (The comparison with forecasts starting two months or more apart is more dramatic.)

It is worth noting that the initial conditions in our forecast experiments are generated with a filtered (and detrended) version of the FSU pseudo-stress fields. This involves a temporal 1-2-1 smoothing of monthly averages, and therefore has nearly all intraseasonal variability removed. Thus the changes in initial conditions from month to month in the forecasts do not simply reflect ISV, but rather reflect somewhat lower frequency wind changes at varying preceding times depending on position. The results show that the sensitivity to these lower frequency components of the wind field (whether real or spurious) supersedes the effects of ISV.

Despite the consistency of the results, conclusions should be considered tentative for two reasons. Our ISV model, though motivated by observations, is highly idealized in many respects and may not represent the real phenomenon faithfully. Also, the ZC model has systematic biases, and quite possibly is overly insensitive to the imposed west Pacific forcing. Both of these potential problems in principle can be reduced with better or more complete models. Awaiting such further studies, the present results can be regarded as suggestive of a secondary role for ISV in relation to ENSO.

Finally, it is interesting to consider the relatively infrequent periods where the imposed intraseasonal forcing *did* have an important effect on forecasts. This leads to question: what characteristics of the coupled system determine whether or not the future is well determined? Is there some feature symptomatic of an uncertain future evolution, beyond our imperfect knowledge of the state of the system at any given time? In general, can we know when predictability is more and less limited? It seems certain that a careful study of these issues will give new insight into the model dynamics, but may also contribute to a better understanding of the real climate system.

#### ACKNOWLEDGMENTS

This work was supported by the National Science Foundation under grant ATM-8612570, and by NOAA under grant NA-84-AA-D-00031. Computations were carried out at the John Von Neumann National Supercomputer Center under award NAC 401.



**WESTERN PACIFIC INTERNATIONAL MEETING  
AND WORKSHOP ON TOGA COARE**

**Nouméa, New Caledonia**

**May 24-30, 1989**

**PROCEEDINGS**

*edited by*


**Joël Picaut \***

**Roger Lukas \*\***

**Thierry Delcroix \***

\* ORSTOM, Nouméa, New Caledonia

\*\* JIMAR, University of Hawaii, U.S.A.



## TABLE OF CONTENTS

<b>ABSTRACT</b> .....	i
<b>RESUME</b> .....	iii
<b>ACKNOWLEDGMENTS</b> .....	vi
<b>INTRODUCTION</b>	
<b>1. Motivation</b> .....	1
<b>2. Structure</b> .....	2
<b>LIST OF PARTICIPANTS</b> .....	5
<b>AGENDA</b> .....	7
<b>WORKSHOP REPORT</b>	
<b>1. Introduction</b> .....	19
<b>2. Working group discussions, recommendations, and plans</b> .....	20
a. Air-Sea Fluxes and Boundary Layer Processes .....	20
b. Regional Scale Atmospheric Circulation and Waves .....	24
c. Regional Scale Oceanic Circulation and Waves .....	30
<b>3. Related programs</b> .....	35
a. NASA Ocean Processes and Satellite Missions .....	35
b. Tropical Rainfall Measuring Mission .....	37
c. Typhoon Motion Program .....	39
d. World Ocean Circulation Experiment .....	39
<b>4. Presentations on related technology</b> .....	40
<b>5. National reports</b> .....	40
<b>6. Meeting of the International Ad Hoc Committee on TOGA COARE</b> .....	40
<b>APPENDIX: WORKSHOP RELATED PAPERS</b>	
<b>Robert A. Weller and David S. Hosom: Improved Meteorological     Measurements from Buoys and Ships for the World Ocean     Circulation Experiment</b> .....	45
<b>Peter H. Hildebrand: Flux Measurement using Aircraft     and Radars</b> .....	57
<b>Walter F. Dabberdt, Hale Cole, K. Gage, W. Ecklund and W.L. Smith:     Determination of Boundary-Layer Fluxes with an Integrated     Sounding System</b> .....	81

## MEETING COLLECTED PAPERS

## WATER MASSES, SEA SURFACE TOPOGRAPHY, AND CIRCULATION

<b>Klaus Wyrtki: Some Thoughts about the West Pacific Warm Pool</b> .....	99
<b>Jean René Donguy, Gary Meyers, and Eric Lindstrom: Comparison of the Results of two West Pacific Oceanographic Expeditions FOC (1971) and WEPOCS (1985-86)</b> .....	111
<b>Dunxin Hu, and Maochang Cui: The Western Boundary Current in the Far Western Pacific Ocean</b> .....	123
<b>Peter Hacker, Eric Firing, Roger Lukas, Philipp L. Richardson, and Curtis A. Collins: Observations of the Low-latitude Western Boundary Circulation in the Pacific during WEPOCS III</b> .....	135
<b>Stephen P. Murray, John Kindle, Dharma Arief, and Harley Hurlburt: Comparison of Observations and Numerical Model Results in the Indonesian Throughflow Region</b> .....	145
<b>Christian Henin: Thermohaline Structure Variability along 165°E in the Western Tropical Pacific Ocean (January 1984 - January 1989)</b> .....	155
<b>David J. Webb, and Brian A. King: Preliminary Results from Charles Darwin Cruise 34A in the Western Equatorial Pacific</b> .....	165
<b>Warren B. White, Nicholas Graham, and Chang-Kou Tai: Reflection of Annual Rossby Waves at The Maritime Western Boundary of the Tropical Pacific</b> .....	173
<b>William S. Kessler: Observations of Long Rossby Waves in the Northern Tropical Pacific</b> .....	185


<b>John S. Godfrey, K. Ridgway, Gary Meyers, and Rick Bailey:</b> Sea Level and Thermal Response to the 1986-87 ENSO Event in the Far Western Pacific .....	291
<b>Joël Picaut, Bruno Camusat, Thierry Delcroix, Michael J. McPhaden, and Antonio J. Busalacchi:</b> Surface Equatorial Flow Anomalies in the Pacific Ocean during the 1986-87 ENSO using GEOSAT Altimeter Data .....	301

#### THEORETICAL AND MODELING STUDIES OF ENSO AND RELATED PROCESSES

<b>Julian P. McCreary, Jr.:</b> An Overview of Coupled Ocean-Atmosphere Models of El Nino and the Southern Oscillation .....	313
<b>Kensuke Takeuchi:</b> On Warm Rossby Waves and their Relations to ENSO Events .....	329
<b>Yves du Penhoat, and Mark A. Cane:</b> Effect of Low Latitude Western Boundary Gaps on the Reflection of Equatorial Motions .....	335
<b>Harley Hurlburt, John Kindle, E. Joseph Metzger, and Alan Wallcraft:</b> Results from a Global Ocean Model in the Western Tropical Pacific .....	343
<b>John C. Kindle, Harley E. Hurlburt, and E. Joseph Metzger:</b> On the Seasonal and Interannual Variability of the Pacific to Indian Ocean Throughflow .....	355
<b>Antonio J. Busalacchi, Michael J. McPhaden, Joël Picaut, and Scott Springer:</b> Uncertainties in Tropical Pacific Ocean Simulations: The Seasonal and Interannual Sea Level Response to Three Analyses of the Surface Wind Field .....	367
<b>Stephen E. Zebiak:</b> Intraseasonal Variability - A Critical Component of ENSO ? .....	379
<b>Akimasa Sumi:</b> Behavior of Convective Activity over the "Jovian-type" Aqua-Planet Experiments .....	389
<b>Ka-Ming Lau:</b> Dynamics of Multi-Scale Interactions Relevant to ENSO .....	397
<b>Pecheng C. Chu and Roland W. Garwood, Jr.:</b> Hydrological Effects on the Air-Ocean Coupled System .....	407
<b>Sam F. Iacobellis, and Richard C.J. Somerville:</b> A one Dimensional Coupled Air-Sea Model for Diagnostic Studies during TOGA-COARE .....	419
<b>Allan J. Clarke:</b> On the Reflection and Transmission of Low Frequency Energy at the Irregular Western Pacific Ocean Boundary - a Preliminary Report .....	423
<b>Roland W. Garwood, Jr., Pecheng C. Chu, Peter Muller, and Niklas Schneider:</b> Equatorial Entrainment Zone : the Diurnal Cycle .....	435
<b>Peter R. Gent:</b> A New Ocean GCM for Tropical Ocean and ENSO Studies .....	445
<b>Wasito Hadi, and Nuraini:</b> The Steady State Response of Indonesian Sea to a Steady Wind Field .....	451
<b>Pedro Ripa:</b> Instability Conditions and Energetics in the Equatorial Pacific .....	457
<b>Lewis M. Rothstein:</b> Mixed Layer Modelling in the Western Equatorial Pacific Ocean .....	465
<b>Neville R. Smith:</b> An Oceanic Subsurface Thermal Analysis Scheme with Objective Quality Control .....	475
<b>Duane E. Stevens, Qi Hu, Graeme Stephens, and David Randall:</b> The hydrological Cycle of the Intraseasonal Oscillation .....	485
<b>Peter J. Webster, Hai-Ru Chang, and Chidong Zhang:</b> Transmission Characteristics of the Dynamic Response to Episodic Forcing in the Warm Pool Regions of the Tropical Oceans .....	493

**MOMENTUM, HEAT, AND MOISTURE FLUXES BETWEEN  
ATMOSPHERE AND OCEAN**

**W. Timothy Liu: An Overview of Bulk Parametrization and Remote Sensing of Latent Heat Flux in the Tropical Ocean ..... 513**  
**E. Frank Bradley, Peter A. Coppin, and John S. Godfrey: Measurements of Heat and Moisture Fluxes from the Western Tropical Pacific Ocean ..... 523**  
**Richard W. Reynolds, and Ants Leetmaa: Evaluation of NMC's Operational Surface Fluxes in the Tropical Pacific ..... 535**  
**Stanley P. Hayes, Michael J. McPhaden, John M. Wallace, and Joël Picaut: The Influence of Sea-Surface Temperature on Surface Wind in the Equatorial Pacific Ocean ..... 543**  
**T.D. Keenan, and Richard E. Carbone: A Preliminary Morphology of Precipitation Systems In Tropical Northern Australia ..... 549**  
**Phillip A. Arkin: Estimation of Large-Scale Oceanic Rainfall for TOGA ..... 561**  
**Catherine Gautier, and Robert Frouin: Surface Radiation Processes in the Tropical Pacific ..... 571**  
**Thierry Delcroix, and Christian Henin: Mechanisms of Subsurface Thermal Structure and Sea Surface Thermo-Haline Variabilities in the South Western Tropical Pacific during 1979-85 - A Preliminary Report ..... 581**  
**Greg. J. Holland, T.D. Keenan, and M.J. Manton: Observations from the Maritime Continent : Darwin, Australia ..... 591**  
**Roger Lukas: Observations of Air-Sea Interactions in the Western Pacific Warm Pool during WEPOCS ..... 599**



<b>Chung-Hsiung Sui, and Ka-Ming Lau: Multi-Scale Processes in the Equatorial Western Pacific</b> .....	<b>747</b>
<b>Stephen E. Zebiak: Diagnostic Studies of Pacific Surface Winds</b> .....	<b>757</b>

#### MISCELLANEOUS

<b>Rick J. Bailey, Helene E. Phillips, and Gary Meyers: Relevance to TOGA of Systematic XBT Errors</b> .....	<b>775</b>
<b>Jean Blanchot, Robert Le Borgne, Aubert Le Bouteiller, and Martine Rodier: ENSO Events and Consequences on Nutrient, Planktonic Biomass, and Production in the Western Tropical Pacific Ocean</b> .....	<b>785</b>
<b>Yves Dandonneau: Abnormal Bloom of Phytoplankton around 10°N in the Western Pacific during the 1982-83 ENSO</b> .....	<b>791</b>
<b>Cécile Dupouy: Sea Surface Chlorophyll Concentration in the South Western Tropical Pacific, as seen from NIMBUS Coastal Zone Color Scanner from 1979 to 1984 (New Caledonia and Vanuatu)</b> .....	<b>803</b>
<b>Michael Szabados, and Darren Wright: Field Evaluation of Real-Time XBT Systems</b> .....	<b>811</b>
<b>Pierre Rual: For a Better XBT Bathy-Message: Onboard Quality Control, plus a New Data Reduction Method</b> .....	<b>823</b>