

Decreased ability of *Pasteuria penetrans* spores to attack to successive generations of *Meloidogyne javanica* ⁽¹⁾

Emmanuel. A. TZORTZAKAKIS*, Simon. R. GOWEN** and Dimitrios. E. GOUMAS*

* National Agricultural Research Foundation, Plant Protection Institute, 71110, Heraklion, Crete, Greece and

** Natural Resources Institute, Chatham, Kent, ME4 4TB, UK.

Accepted for publication 28 February 1995.

Summary – The decreasing ability to attachment of *Pasteuria penetrans* spores on a field population of *Meloidogyne javanica* from Malawi and single egg mass lines of the same species from Crete was recorded in the laboratory. When the bacterial parasite does not provide sufficient control of the nematode, a selection of *Pasteuria* spores with high attachment potential against the target nematode might be an alternative for delaying the decrease of spore attachment ability.

Résumé – Décroissance de l'aptitude des spores de *Pasteuria penetrans* à l'attaque de générations successives de *Meloidogyne javanica* – La décroissance de l'aptitude à l'attaque des spores du parasite bactérien *Pasteuria penetrans* est mise en évidence au laboratoire vis-à-vis d'une population sauvage de *Meloidogyne javanica* originaire du Malawi et de lignées provenant d'une seule masse d'œufs de cette même espèce originaire de Crète. Dans le cas où le parasite bactérien ne permet pas un contrôle suffisant du nématode, une sélection de spores de *Pasteuria* à haut potentiel d'adhésion vis-à-vis du nématode cible pourrait constituer une alternative à la décroissance de l'aptitude à l'adhésion des spores.

Key-words : attachment, biological control, *Pasteuria penetrans*, *Meloidogyne javanica*, spore adhesion.

Pasteuria penetrans (*Pp*) is an endospore forming bacterium pathogenic to root-knot nematodes (Sayre & Starr, 1985) with potential for biocontrol when employed in container grown crops, small plots and polytunnels (Stirling, 1984; Brown *et al.*, 1985; Dube & Smart, 1987; Gowen & Ahmed, 1990; Gowen & Tzortzakakis, 1994; Tzortzakakis & Gowen, 1994 *b*). Previous work demonstrated that a mixed population of *Meloidogyne javanica* and *M. incognita* was subjected to selection pressure for decreased susceptibility to spore attachment and infectivity after exposure to a population of the parasite for four generations. That could be successfully prevented when a mixture of parasite isolates (*Pp* blend) was used instead of a single isolate (Tzortzakakis & Gowen, 1994 *a*). Several authors addressed the problem of specificity of spore attachment on several *Meloidogyne* populations and concluded that for durable control, *Pp* isolates with a broad spectrum of host specificity should be used (Stirling, 1985; Davies *et al.*, 1988 *a, b*; Channer & Gowen, 1992). This investigation aims to study the effectiveness of a mixture of parasite isolates (*Pp* blend) in preventing selection for decreased susceptibility to spore attachment in *Meloidogyne* species with narrow genetic diversity, such as a population or single egg mass lines.

The studies were carried out in pot experiments at the University of Reading, UK and the Plant Protection Institute of Heraklion, Greece.

Materials and methods

Pp ISOLATES AND NEMATODE POPULATIONS ORIGINS

Spore suspensions were prepared according to the method of Stirling and Wachtel (1980). Powdered tomato roots containing *Pp* infected females were wet ground in a mortar and pestle and the resulting slurry passed through a 38 µm sieve to separate the spores from coarse plant material. The suspensions prepared at the beginning of the experiments were stored in a domestic refrigerator. The origins of spore isolates were: *Pp* 1 (Australia), *Pp* 2 (USA), *Pp* 3 (S. Africa), *Pp* PNG (Papua New Guinea), *Pp* M (Malawi) and *Pp* IC (Ivory Coast). The tested *Meloidogyne* populations were identified morphologically (perineal pattern of mature female, tail length and hyaline portion of J2) and by the North Carolina differential host test (Hartman & Sasser, 1985; Jepson, 1987).

The nematode population used in the first experiment was a population of *M. javanica* originally isolated from

⁽¹⁾ The work carried out at Reading University represents a portion of a PhD Thesis submitted by the senior author in the Department of Agriculture.

a field site in Malawi and maintained on tomato cv. Tiny Tim for many years in a glasshouse at Reading University.

The populations used in the second experiment were single egg mass lines of *M. javanica* originating from two plastic mini-greenhouse frames in Crete and maintained for one generation on tomato cv. Rutgers in a growth chamber at the Plant Protection Institute of Heraklion, Greece. These populations were chosen from several populations of the same species after testing for susceptibility to spore attachment.

EXPERIMENT 1

Juveniles (J2s) were collected from egg masses incubated on extraction filters (Southey 1986) for 4 days and were immersed in a spore suspension (3×10^4 spores/ml) at 28 °C. Spore attachment was recorded on 50 individuals under an inverted microscope at $\times 400$. The minimum level of spore attachment before plant inoculation was that at which at least 80 % of the nematodes were encumbered with more than five spores. Nematodes from the same source left in tap water under the same conditions served as controls (original population which was not exposed to spores). Juveniles contaminated with spores of *Pp* blend (the six isolates blended) and *Pp*-free controls were inoculated on tomato plants cv. Tiny Tim. Cross contamination was avoided by placing saucers beneath individual plant pots and watering by hand with care. Each treatment was replicated four times. After a 5-week-growing period in a glasshouse with air temperature 25–30 °C, the egg masses were collected from the roots and incubated in extraction dishes. The same procedure of exposure to spores was repeated for four generations using highly concentrated spore suspensions (up to 10^5 spores/ml) to hasten the selection and achieve the desired attachment level. Plant inoculation was done with 1000–4000 nematodes per pot depending on the size of the seedlings.

At the end of each crop cycle 100 J2s collected from the egg masses of each plant (selected and original popu-

lations) were incubated in 3 ml spore suspensions of *Pp* blend and *Pp2* containing a total of 20 000 spores in 2.5 cm Petri dishes at 28 °C. Spore attachment was recorded on 20 J2s from each of the four replicate dishes after 24 h. The J2s finally collected after the completion of the fourth generation were also exposed to *Pp3*, *PpPNG*, *PpM* and *PpIC* under the same conditions.

EXPERIMENT 2

The procedure was the same as that described in the first experiment. The plants grew for six weeks in a growth room with air temperature 26–30 °C. Attachment tests were conducted as previously described, using the *Pp* blend (used for selection) and some of the individual isolates (*Pp3*, *PpPNG*, *PpM*, and *PpIC*) in 5 cm diameter Petri dishes containing 30 000 spores of the parasite in a total volume of 5 ml. The experiment was terminated after recording spore attachment on the first generation J2s.

Data were transformed to square roots before analysis when the value for the standard error of the mean was higher than a mean value (Mead & Curnow, 1990). Comparisons of main effects were made using the F value calculated by two factor analysis of variance between nematode populations and *Pasteuria* isolates.

Results

EXPERIMENT 1

Significantly fewer spores of *Pp* blend and *Pp2* attached to the J2s of the selected nematode population after the completion of the third ($P < 0.01$) and fourth generations ($P < 0.001$) (Table 1). The selected population indicated also decreased susceptibility when exposed to four of the *Pp* isolates ($P < 0.001$) composing the blend (Table 2).

EXPERIMENT 2

Several single egg mass populations of *M. javanica* originating from plastic tunnels in Crete were preliminarily tested indicating mostly either high (> 15 spores/J2) or low (0–5 spores/J2) susceptibility to spore attach-

Table 1. Number of attached spores of *Pasteuria penetrans* (*Pp* blend and *Pp2*) on the original population of *Meloidogyne javanica* (Malawi) and the selected population previously exposed to *Pp* blend for four generations.

	Generations							
	1		2		3		4	
	<i>Pp</i> blend	<i>Pp2</i>	<i>Pp</i> blend	<i>Pp2</i>	<i>Pp</i> blend	<i>Pp2</i>	<i>Pp</i> blend	<i>Pp2</i>
Original	10.88	10.88	10.30	7.95	10.77	10.82	13.52	7.58
Selected	9.47	8.67	9.13	8.56	8.77	7.02	1.51	2.35
	NS		NS			**	*** a	

Number of spores attached to 20 J2s per dish (average of four replicate dishes); a : analysis on square root transformed data; **, *** significant differences at $P < 0.01$ and 0.001 , respectively, between original and selected nematode population; NS : non significant difference.

- DAVIES, K. G., KERRY, B. R. & FLYNN, C. A. (1988 b). Observations on the pathogenicity of *Pasteuria penetrans*, a parasite of root-knot nematodes. *Ann. appl. Biol.*, 112 : 491-501.
- DUBE, B., SMART, G. C. Jr. (1987). Biological control of *Meloidogyne incognita* by *Paecilomyces lilacinus* and *Pasteuria penetrans*. *J. Nematol.*, 19 : 222-227.
- GOWEN, S. R., & AHMED, R. (1990). *Pasteuria penetrans* for control of pathogenic nematodes. *Aspects appl. Biol.*, 22 : 437-438.
- GOWEN, S. & TZORTZAKAKIS, E. (1994). Biological control of *Meloidogyne* spp with *Pasteuria penetrans*. *EPPO Bull.*, 24 : 495-500.
- HARTMAN, K. M. & SASSER, J. N. (1985). Identification of *Meloidogyne* species on the basis of differential host test and perineal-pattern morphology. In : Barker, K. R., Carter, C. C. & Sasser, J. N. (Eds). *An advanced treatise on Meloidogyne. Vol II. Methodology*. North Carolina State University, USA : 69-77.
- JEPSON, S. B. (1987). *Identification of root knot nematodes (Meloidogyne species)*. Wallingford, UK, CAB International, 252 p.
- MEAD, R. & CURNOW, R. N. (1990). *Statistical methods in agriculture and experimental biology*. Chapman & Hall : 104-124.
- SAYRE, R. M. & STARR, M. P. (1985). *Pasteuria penetrans* (ex Thorne 1940) nom. rev., comb. n., sp. n., a mycelial and endospore forming bacterium parasitic in plant parasitic nematodes. *Proc. helminth. Soc. Wash.*, 52 : 149-165.
- SOUTHEY, J. F. (1986). *Laboratory methods for work with plant and soil nematodes*. London, HMSO, MAFF Reference Book 402, 202 p.
- STIRLING, G. R. (1984). Biological control of *Meloidogyne javanica* with *Bacillus penetrans*. *Phytopathology*, 74 : 55-60.
- STIRLING, G. R. (1985). Host specificity of *Pasteuria penetrans* within the genus *Meloidogyne*. *Nematologica*, 31 : 203-209.
- STIRLING, G. R. & WACHTEL, M. F. (1980). Mass production of *Bacillus penetrans* for the biological control of root knot nematodes. *Nematologica*, 26 : 308-312.
- TZORTZAKAKIS, E. A. & GOWEN, S. R. (1994 a). Resistance of a population of *Meloidogyne* spp. to parasitism by the obligate parasite *Pasteuria penetrans*. *Nematologica*, 40 : 258-266.
- TZORTZAKAKIS, E. A. & GOWEN, S. R. (1994 b). The evaluation of *Pasteuria penetrans* alone and in combination with oxamyl, plant resistance and solarization for control of *Meloidogyne* spp. on vegetables grown in greenhouses in Crete. *Crop Protect.*, 13 : 455-462.