

NEMATODES AND AGRICULTURE IN CONTINENTAL ARGENTINA.  
AN OVERVIEW

Marcelo E. DOUCET and María M.A. DE DOUCET

Laboratorio de Nematología, Centro de Zoología Aplicada, Facultad de Ciencias Exactas,  
Físicas y Naturales, Universidad Nacional de Córdoba, Casilla de Correo 122, 5000 Córdoba, Argentina.

Accepted for publication 5 November 1996.

**Summary** – In Argentina, soil nematodes constitute a diverse group of invertebrates. This widely distributed group includes more than two hundred currently valid species, among which the plant-parasitic and entomopathogenic nematodes are the most remarkable. The former includes species that cause damages to certain crops (mainly *Meloidogyne* spp, *Nacobbus aberrans*, *Ditylenchus dipsaci*, *Tylenchulus semipenetrans*, and *Xiphinema index*), the latter includes various species of the Mermithidae family, and also the genera *Steinernema* and *Heterorhabditis*. There are few full-time nematologists in the country, and they work on taxonomy, distribution, host-parasite relationships, control, and different aspects of the biology of the major species. Due to the importance of these organisms and the scarcity of information existing in Argentina about them, nematology can be considered a promising field for basic and applied research.

**Résumé** – *Les nématodes et l'agriculture en Argentine. Un aperçu général* – Les nématodes du sol représentent en Argentine un groupe très diversifié. Ayant une vaste répartition géographique, il comprend actuellement plus de deux cents espèces, celles parasitant les plantes et les insectes étant considérées comme les plus importantes. Les espèces du genre *Meloidogyne*, ainsi que *Nacobbus aberrans*, *Ditylenchus dipsaci*, *Tylenchulus semipenetrans* et *Xiphinema index* représentent un réel danger pour certaines cultures. Les parasites d'insectes comprennent plusieurs espèces de Mermithidae, ainsi que des représentants des genres *Steinernema* et *Heterorhabditis*. Les nématologistes travaillant en Argentine sont peu nombreux. Ils poursuivent des recherches très diversifiées: taxinomie, répartition, relation hôte-parasite, résistance, contrôle, biologie des principales espèces. Etant donné l'importance de ces organismes et leur connaissance encore incomplète, la nématologie peut être considérée comme ayant en Argentine un futur plein de promesses dans les domaines des recherches fondamentale et appliquée.

**Key words:** Agriculture, Argentina, nematology.

This article is an overview of the main features of the history of nematology in Argentina and its current status, with emphasis on its association with agriculture.

The information presented here comes from a critical analysis of the most relevant bibliographic information, from the personal experience of the authors and from their interaction with other specialists in the country.

Mainland Argentina has an area of 2 791 810 km<sup>2</sup> and is characterised by a great diversity of climates (Chiozza & van Domselaar, 1958) (Fig. 1) and soils (Papadakis, 1964; Moscatelli, 1990). This diversity has favoured the development of diversified agriculture, with both extensive and intensive farming systems in different regions of the country (Fig. 2). Extensive farming is used for crops such as wheat, soybean, corn, sorghum, sunflower, rice, sugar cane, barley, rye, alfalfa, peanut, rape, tobacco, grapevine, sweet potato, and potato; intensive farming produces all kinds of vegetables, and mostly tomato, garlic, pepper, beans, asparagus, and lentils. In addition, many

types of fruit trees are grown in wide areas throughout the country (Gomez Riera, 1992). Aromatic and ornamental plants also are important items in the vegetal production of Argentina.

Throughout its development, Argentine agriculture has adapted to the demand of international markets. Because of this, different crops have received priority in different regions. Currently, soybean, wheat, corn, sunflower and potato are the most important species from the viewpoint of exports and domestic consumption (Cap *et al.*, 1993a).

Within this context of marked agricultural diversity, there exist numerous species of soil nematodes belonging to the major trophic categories (plant-parasitic, free-living, mycophagous, predators, and entomopathogenic species).

Plant-parasitic nematodes include some species that can cause serious damages to several crops, while various entomopathogenic species could be used as biological control agents against harmful insects. Recent results (Doucet & Doucet, 1996) and older data as




Fig. 1. Types of climate in Argentina (adapted from Chiozza & van Domselaar, 1958).




Fig. 2. Distribution of the main crops in Argentina.

presented in this article emphasise the great diversity of soil nematodes in Argentina.

### Historical background

The first reference to soil nematodes in Argentina is related to entomopathogenic species. A German naturalist (Weyembergh, 1876) working in the National Academy of Sciences (Córdoba City) described under the name of *Gordius acridiorum* a parasite associated to one of the most important agricultural pest at that time: the locust *Schistocerca paranensis* (Burm). Later, the same investigator, with others also from Europe, reported the afore mentioned species and *G. acrydii* in locusts from Buenos Aires and Córdoba (Weyembergh, 1878, 1879; Conil, 1878, 1879; Bruner, 1898). The descriptions made at that time were not detailed, but the data published by these authors on the life-cycle and interactions of these parasites with their hosts make it obvious that these nematodes belonged to the family Mermithidae (Berg, 1898).

Only fifteen years separate the first reference to entomopathogenic species and the discovery of a plant-parasitic nematode. At the beginning of the 1890's, the Head of the Chemistry Office of the Municipality of Córdoba city – a chemist and naturalist from Germany – informed the Head of the National Department of Agriculture at Buenos Aires of the finding of "*Heterodera radicolica* de Greef" associated with grapevine, and reported some problems caused by phylloxera. He also mentioned that this pathogen had also been observed on roots of peach trees (Doering, 1891) and he added: "we have found the little animal together with the eggs in the cavities of the root-knot" (clearly a reference to the genus *Meloidogyne*).

The subsequent development of nematology in the country followed different paths for plant-parasitic, entomopathogenic, and free-living soil nematodes.

Plant-parasitic nematodes, because of their close association with economic losses to the agriculture, have received the most attention. We will now describe the activities that took place in Argentina during the development of nematological research concerning each of the three types of nematodes.

### Plant-parasitic nematodes

At the beginning to this century, several articles were published on *Heterodera radicolica* (now *Meloidogyne* spp.) associated with tomato plants. Those papers are remarkable by the good information they provide on yield loss in the attacked plants, some aspects of the life cycle of the nematode and its relationships with some environmental factors, and the most convenient methods for reduction of the nematode damages (Lynch Arribáizaga, 1901; Huergo 1902, 1903).

From that time on, several species were sporadically detected in different regions, and linked to problems observed in several crops.

As time went by, some people from national organizations – i) Ministry of Agriculture; Instituto Nacional de Tecnología Agropecuaria (INTA): Estación Regional Experimental Balcarce (Buenos Aires province), Roque Saenz Peña (Chaco province), San Rafael (Mendoza province); ii) Universities (Universidad de Buenos Aires, Universidad Nacional de Córdoba, Universidad Nacional de Cuyo, Universidad Nacional de La Plata) ; provincial organizations – Dirección de Sanidad Vegetal, Ministerio de Asuntos Agrarios, Buenos Aires province; Estación Experimental Agro Industrial Obispo Colombes, Tucumán province ; and private organizations – Centro de Investigaciones Nematológicas de Corrientes – became interested in the studies of these organisms. Agriculture Engineers Burkart, Fresa and Marchionatto and Dr. R. Gutierrez were some of the people who out stood in some of those institutions for their work. Thus, "centres" of nematological activities emerged in the Buenos Aires, Mendoza and Tucumán provinces; in these last two provinces, Engineers Eduardo Vega and Miguel Costilla developed important applied works related to management of plant-parasitic nematodes associated to numerous crops. The first articles defining the status of plant-parasitic nematodes in relation to agriculture in Argentina were prepared in Buenos Aires and Mendoza (Moreno, 1964; López Cristóbal, 1965; Vega, 1971). Dr. Amalia Moreno worked in the Patología Vegetal Institute (Centro de Investigaciones Agropecuarias, INTA Castelar, Buenos Aires) where she analysed samples from different sites in the country and obtained the initial information about the distribution of different species and their relationships with their hosts.

In 1980, a new "centre" of nematological activities was created at the National University of the city of Córdoba, under the direction of the authors of this article. Both had studied nematology at "Station de Recherches sur les Nématodes" (INRA, Antibes, France). In Córdoba, they engaged in the study of soil nematodes (plant-parasitic, entomopathogenic and free-living species) and organised courses on Agricultural nematology. The first course – "Introduction to nematology" – in Argentina was presented in the Centro by the present authors in July 1980.

Almost at the same time, another nematologist, Dr. E. Chaves, was being formed at the University of Ghent, Belgium, who later joined the Unidad Integrada EEA Balcarce, Universidad Nacional de Mar del Plata, Buenos Aires.

Subsequently nematological activities started at the Universidad Nacional de Cuyo, Mendoza province (control of plant-parasitic nematodes) and at the Uni-

versidades Nacionales of Buenos Aires and La Plata (Buenos Aires province) (survey of plant-parasitic nematodes and taxonomy-biology of entomopathogenic nematodes, respectively).

More recently, other young researchers from Argentina have done postgraduate studies in Belgium and the USA, on taxonomy and the search for resistant characters.

### Agricultural losses caused by plant-parasitic nematodes

In general, agricultural losses caused by nematodes are not yet known with any degree of accuracy.

According to specialised agronomists, it is possible to evaluate approximate yield reductions in some zones or average production losses for specific cultures. For example, in the western part of the country, tomato and peach trees were affected until 1970 by *Meloidogyne* spp. that reduced the yields by 20 % and 15 %, respectively. *Meloidogyne* spp. and *Xiphinema index* attacks on grapevine reduced yields by about 5 %. The use of resistant varieties and nematicides products have now reduced the losses to 2, 1, and 3 %, respectively. On garlic, *Ditylenchus dipsaci* caused damages that affected 30 % of the production between 1965-1970; the selection of non contaminated seeds and different methods of control have reduced the damages to less than 1 % of the production (Vega, pers. comm.).

Potato can be attacked by *Meloidogyne* spp. and/or *Nacobbus aberrans*, depending on the regions. The main problems occur for seed potato production. Species of *Meloidogyne* have been observed in the West (Vega *et al.*, 1986; Doucet & Ponce de León, 1996a) affecting 40 % of the production, the North-West (Costilla, comm. pers.), and the South-East (Chaves & Torres, 1993; Doucet & Ponce de León, 1996a) affecting 30 % of the production (Vega, comm. pers.).

*Nacobbus aberrans* has been found so far only in Tucumán and Buenos Aires areas (Costilla *et al.*, 1978; Chaves & Torres, 1993). The current legislation prohibits the use of infected tubercles for seeds; they are sold for food at lower prices.

These and other nematodes can cause a decrease of the production of several other crops. This situation has been evident for some years in horticultural crops under hot-house conditions; the economical loss has not yet been evaluated with precision.

### Current list of nematode species in the country

The number of nematologists specialised in population characterisation, and identification and description of species has been and still is very low.

Unfortunately, some doubts have been expressed on past identification of some species because the vari-

ability of the characters used was not taken into account and detailed analyses of the populations were not conducted. Some identifications were probably inaccurate because those species have never been found again, although many samples were analysed. Furthermore, voucher specimens were almost never kept in local institutions or deposited in known collections.

As a consequence, it is difficult to recognise which species are of quarantine importance for the country. In some cases, the commercialisation of some agricultural products is difficult on international markets.

The list of identified species is given in the Appendix.

### Active fields of research

The studies on plant-parasitic nematodes in Argentina can be divided into two main categories.

The first one is related to the acquisition of basic information: distribution, host range, some host-parasite relationships, identification and characterisation of populations of various species, and some aspects of the biology of new species. This kind of information has been obtained on the most common plant-parasitic nematodes associated with the major crops of the country (López Cristóbal, 1946, 1965; Moreno, 1958; La Red & Vega, 1968; Vega, 1971; Costilla *et al.*, 1976; Chaves, 1984; Doucet, 1986, 1989; Chaves & Torres, 1993). These activities are still being developed, and expanded to different genera and species of nematodes.

The second category is related to the management of problems caused by harmful nematodes.

From the very first projects on plant-parasitic nematodes in Argentina, the necessity of controlling nematode populations has been emphasised. The use of various chemical products has been proposed and it has been recommended to use preventive measures to avoid spreading these pathogens (Doering, 1891; Lynch Arribálzaga, 1901; Huergo, 1902, 1903). Then, several methods of crop protection were implemented.

In different cases, resistant varieties were selected (Burkart, 1934, 1937; Galmarini, 1978; Pucci & Avila, 1978; Sisler & Casaurang, 1983; March *et al.*, 1985; Costilla, 1985, 1986; Castellanos & Del Toro, 1994), crop rotation was established (Vega, 1981; Costilla, 1991), and different chemical methods were used, as explained below under the major nematodes species. Several nematicides have been tested, including DD, DBCP, methyl bromide (Gargiulo & Moyano, 1948; Vega & Gatica, 1968, 1972; Del Toro & Vega, 1982), carbofuran (Costilla, 1973), and aldicarb (Del Toro, 1988a). Because of their high cost, these are used generally in crops with high profitability.

Experiments have started on the use of solarization techniques as an alternative control method.

Finally, surveys have recently been initiated to delimitate areas free of specific harmful nematodes. This should make it possible to commercialise agricultural products in accordance with international regulations (Del Toro *et al.*, 1994a).

### Entomopathogenic nematodes

After the discovery that marks the beginning of Argentine nematology (Weyembergh, 1876), more than half a century passed before a second entomopathogenic nematode was detected: the mermithid *Agamermis decaudata* found parasitizing specimens of the locust *Dichroplus arrogans* Stal in Buenos Aires (Gutierrez, 1945).

At the beginnings of the 1970's, nematodes belonging to the Steinernematidae and Heterorhabditidae families, natural enemies of Coleoptera of agronomical importance (genus *Graphognatus*), were reported for the first time in the humid pampa soils (Ahmad, 1972).

The last 15 years have seen the beginning of systematic studies of entomopathogenic nematodes. Research concentrates almost exclusively on specimens of the Mermithidae, Steinernematidae and Heterorhabditidae families. Only a few studies concern nematodes of the Thelastomatidae, Rhabditidae and Diplogasteridae families (Miralles, 1982; Doucet & Doucet, 1989, 1992; Stock & Camino, 1991; Stock, 1993c).

In general, the nematode families which are being studied are those that include species that are being used, or may be used, as an alternative for chemical control of insect pests.

### Active fields of research

Research has been developed on the following subjects:

#### TAXONOMIC STUDIES

These include the characterisation of local populations of known species and also descriptions of new genera and species in the Mermithidae (Doucet, 1982, 1986a; Doucet & Poinar, 1984; Camino, 1985a, b, c, 1987, 1988a, 1990, 1991, 1993; Camino & Poinar, 1989; Camino & Stock, 1989, 1994; Stock & Camino, 1992a,b), Steinernematidae (Doucet, 1986b, 1995; Poinar *et al.*, 1988; Doucet & Doucet, 1990; Stock, 1993b), and Heterorhabditidae families (Stock, 1993a; Doucet *et al.*, 1996).

Several articles on species of the last two families have emphasised the need to consider the intraspecific variability of the characters used (Doucet & Doucet,

1986; Doucet *et al.*, 1990, 1991, 1992a, Doucet *et al.*, 1996).

#### BIOLOGICAL STUDIES

The published studies described the embryonic development and life cycle of the mermithids (Camino, 1986; Camino & Rebores, 1994) and the life cycle of heterorhabditids (Doucet & Poinar, 1985; Doucet *et al.*, 1996).

Studies on host range and aggressivity of *Heterorhabditis bacteriophora* Poinar, 1975 have recently started (Doucet *et al.*, 1992b; Doucet & Giayetto, 1994; Doucet *et al.*, 1996).

#### ECOLOGICAL STUDIES

Studies concern the geographic distribution of mermithids, steinernematids, and heterorhabditids (Doucet & Cagnolo, 1995a; Doucet & Bertolotti, 1995; Doucet & Doucet, 1996) and the spatial distribution patterns of Heterorhabditidae and Steinernematidae infective larvae in the soil (Doucet, 1992).

These studies continue in the Centro de Estudios Parasitológicos y Vectores (La Plata, Buenos Aires province), the Laboratorio de Nematología of the Centro de Zoología Aplicada, and the Cátedra de Parasitología, Facultad de Ciencias Exactas, Físicas y Naturales (Universidad Nacional de Córdoba).

### Free-living soil nematodes

Most of the articles on free-living nematodes concern descriptions of species from various parts of the country. In particular, *Cruzinema tripartitum* has been studied in relation to its possible association with a disease of garlic (Gutierrez, 1949; Avila & Pucci, 1980; Doucet & Ponce de León, 1994) and its eventual behaviour as a facultative entomopathogenic species (Doucet & Doucet, 1989; Doucet, 1994). Another interesting case is that of the free-living species *Distolabrellus veechi* Anderson, 1983 which, in specific circumstances, becomes a facultative entomopathogenic species (Doucet & Doucet, 1992).

### The Argentine Society of Nematology

In 1980, at the request of INTA, the Argentine Society of Nematology was created with the objective of coordinating nematological activities in the field and promoting contacts between the specialists in the country. Its first President was the senior author of the present article and one of the most important actions of the Society was to create an awareness of the agricultural significance of soil nematodes, the damages they cause, and the possible ways to manage these parasites. To this aim, courses were organised in several institutions all over the country for teachers, technicians, researchers and other professionals

involved in plant pathology. These courses proved to be one of the best ways of promoting nematology. Among those courses, one of the most remarkable was the one sponsored by ORSTOM (France) in 1986 at the Centro de Zoología Aplicada. The growing awareness of nematode problems initiated by these courses among agriculture technicians and farmers resulted in the passing by the national and provincial Ministries of Agriculture of laws intended to limit the spread of nematodes harmful to agriculture. In this manner, not only is agricultural production preserved but also the products obtained are clean and can be commercialised abroad.

### Present situation

At the present, there are about ten researchers working full time on nematology. The studies that are carried out concern several subjects (survey, taxonomy, biology, management, search for resistance sources) and are principally related to plant-parasitic and entomopathogenic nematodes. Occasionally, the few specialised laboratories offer identification services for the species of importance for various crops and act as consultants for organisations related to agriculture at the national, provincial and private levels.

### Major genera and species of plant-parasitic nematodes reported in Argentina and the problems they cause

From the beginning, attention has focused on harmful species in the genera *Ditylenchus*, *Meloidogyne*, *Nacobbus*, *Tylenchulus*, and *Xiphinema*.

In this context, it is necessary to mention the occurrence of the species *Ditylenchus dipsaci*, *Meloidogyne* spp., *Nacobbus aberrans*, *Tylenchulus semipenetrans* and *Xiphinema index* associated to cultivated alfalfa and garlic, potatoes and several vegetables, citrus and grapevine, respectively. Because their possible economic importance, other genera (*Pratylenchus* and *Helicotylenchus*) are also taken into account.

#### GENUS *DITYLENCHUS*

Two species are found in Argentina: *D. intermedius* and *D. dipsaci*.

#### *Ditylenchus intermedius*

Found in association with the tree *Platanus* sp. (Moreno & Turica, 1960), *P. acerifolia* (Brugnoni, 1980), *Citrus* sp., and grass (Moreno, 1961). The impact of this nematode on agriculture has not yet been evaluated.

#### *Ditylenchus dipsaci*

This species has created problems mostly on alfalfa and garlic. It was first recorded in the country associated with damages on alfalfa in the North-West (Anon., 1929). Several studies on *D. dipsaci* were carried out before the 1950's. The species was believed to be widely distributed in the central and northern parts of the country and to be responsible for damages of variable importance (Baez, 1925; Marchionatto *et al.*, 1926; Anon., 1929; Blanchard, 1930; Burkart, 1943; López Cristóbal, 1946; Moreno, 1964). Observations made at that time revealed the presence of plants immune to this nematode in some fields in Buenos Aires. Resistant varieties of alfalfa were then selected for the management of *D. dipsaci* (Burkart, 1934, 1937; Ragonese & Marcó, 1943; Miccio Peralta, 1956; Godeck & Favret, 1965; Godeck & Stilinovic, 1966). Since then, no more damages caused by this plant-parasitic nematode have been detected in alfalfa fields.

The crop that was the most affected by *D. dipsaci* is garlic. The climatic and edaphic conditions of the western, central-southern and south-eastern regions contributed to the fast increase of populations wherever infested seeds were used (Cucchi *et al.*, 1967; Urbietta Salvarredi *et al.*, 1971). *D. dipsaci* was first found on garlic in 1965 when it provoked a total loss of the garlic production in some areas of the province of Mendoza, the major producer of garlic in the country (Cucchi *et al.*, 1967). Simultaneously, the close association of this nematode with fungi to form a major pathologic complex was detected (Avila & Pucci, 1980). This situation was at the origin of detailed studies on the behaviour of the nematode (Urbietta Salvarredi, 1972), which resulted in the implementation of several forms of control, including thermotherapy (Del Toro & Mavrigh, 1977), chemical control with synthetic nematicides (Rivera & Del Toro, 1982; Del Toro *et al.*, 1988), and use of natural products (Del Toro, 1988b).

Today, garlic is grown on about 15 000 ha distributed in various regions, mainly the Central-West (Burba, 1992). Given the specific requirements of the Argentine regulatory legislation, special care is taken for the production of nematode-free seeds, which is enough to ensure an excellent production of garlic without nematode damages (Del Toro *et al.*, 1994b).

It seems that the host range of *D. dipsaci* is wide in Argentina. This nematode has been found in pastures (Burkart, 1935; Moreno & Fresa, 1969), wheat (Brugnoni, 1966), ornamental plants and weeds (Burkart, 1937; Moreno, 1956a; Doucet, 1992; Margegiani & Russo, 1992), cotton (Mallo, 1961), and forest trees (Moreno & Turica, 1960; Brugnoni, 1980).

*Ditylenchus* spp.

Unidentified species of *Ditylenchus* have been reported in association with sugar cane in the north-western part of Argentina (Costilla *et al.*, 1976) and with cotton (Gutierrez, 1959).

GENUS *PRATYLENCHUS*

This genus is widely distributed throughout the country. So far, fourteen species have been reported in Argentina. The population densities of these species are generally low, except in the North-West where *Pratylenchus* spp. cause damages to banana plantations (Costilla, 1973) and can affect the development and yield of corn (Costilla, 1992).

GENUS *NACOBBUS*

This genus was first found in the mountainous region of the province of Tucumán (2000 m above sea level), in soils of potato fields where it parasitises roots of cultivated and wild potatoes, *Cucurbita maxima*, *Beta* spp., and some weeds (Costilla *et al.*, 1977).

Subsequent studies showed that this nematode is widely distributed in the country (Chaves & Sisler, 1980; Chaves, 1984; Costilla, 1985; Doucet, 1989) and that it can be associated with a large number of plants, such as potatoes, vegetables and weeds (Ojeda *et al.*, 1978; Agüero *et al.*, 1984; Costilla, 1985; Doucet & Ponce de León, 1986; Ponce de León & Doucet, 1989; Doucet *et al.*, 1992; Doucet, 1992).

The analysis of morphological characters of various populations of *Nacobbus* from Argentina showed that they all belong to the species *N. aberrans* (Chaves, 1984; Costilla, 1985; Doucet, 1989; Doucet & Di Rienzo, 1991). However, preliminary observations of the behaviour of different populations in relation to different hosts revealed significant differences (Costilla, 1986). Subsequent studies based on morphometric characters (Doucet & Di Rienzo, 1991) and isoenzyme phenotypes (Doucet & Gardenal, 1992; Doucet *et al.*, 1996) revealed important differences between populations. This supports the hypothesis of a complex of forms within the species *N. aberrans* (Jatala & Golden, 1977).

The major problems caused by this species are related to the cultivation of potato (Ojeda *et al.*, 1978; Costilla, 1985).

Besides forming knots in the roots, third and fourth stage juveniles settle below the lenticels of the tuber where they reach a state of anhydrobiosis. In this state, the nematodes remain viable during storage and may be dispersed by contaminated tuber seed (Costilla, 1985). *N. aberrans* causes significant damages to crops, seriously affecting production. Because of this, the legislation requires the use of non-contaminated potato seed.

Various strategies are used against this nematode, such as the use of nematicides (Costilla & Gomez, 1981), the assessment of resistance of different cultivars (Sisler & Casaurang, 1983), and crop rotation (Costilla, 1991).

A search for sources of resistance in *Lycopersicon* has recently been launched (Cap *et al.*, 1993b).

GENUS *HELICOTYLENCHUS*

This genus is widely distributed in Argentina. It includes several populations, most of which have not been identified at the species level.

The most important species identified so far – *H. multincinctus* – was found in the North-West on roots of banana trees, where it produces characteristic lesions (Costilla *et al.*, 1979).

*GLOBODERA ROSTOCHIENSIS*

This nematode was first reported in Argentina in the soil of a garlic field in the western part of the country (Moreno, 1956b). Later, it was mentioned in the Andes, Province of Jujuy, parasitizing roots of wild potatoes (Brücher, 1960, 1961). Years later, it was reported in the soil of a potato field in another mountainous region in the West (Virsoo, 1967).

Since then, the numerous analyses carried out by different laboratories on soil samples and tubers from many different regions all had negative results (Chaves, 1993). Work on cyst nematodes from Argentina suggests that this species is not present in the country (Chaves, 1987, 1993; Chaves & Torres, 1993).

Because of the first four bibliographic references, some countries hinder exports of certain agricultural products and demand regulatory certifications. However, there are no objective proofs of the occurrence of *G. rostochiensis* in the country.

For the last two years, surveys have been in progress in order to demonstrate that specific areas of garlic, potato and onion production are free of *G. rostochiensis* (Del Toro *et al.*, 1994a).

GENUS *MELOIDOGYNE*

As mentioned above, the discovery of a species of this genus marks the beginning of the study of plant-parasitic nematodes in Argentina. Since then, species of the genus *Meloidogyne* have been found on roots of various plants: vegetables, potatoes, tobacco, soybean, cotton, ornamental plants, fruit trees, forest trees, and weeds.

Reviews of the existing information on the genus have recently been published (Doucet & Pinochet, 1992; Doucet, 1993; Doucet & Ponce de León, 1996b).

Nine species have been detected: *M. acrita*, *M. arenaria* (race 2), *M. chitwoodi*, *M. cruciani*, *M. decalineata*, *M. hapla*, *M. incognita* (races 1, 2,



and 3), *M. javanica*, *M. ottersoni*, in addition to some populations reported as *Meloidogyne* spp. whose specific identity is unknown.

One hundred plant species (belonging to 33 families) can be parasitised by the eight species of *Meloidogyne*. *M. incognita* and *M. arenaria* are the most frequently found species, and they parasitise 56 % and 27 % of plants, respectively (Doucet, 1993). Cultivated as well as wild plants (native and introduced) can be attacked by these nematodes, but mostly weeds and vegetables.

Although the genus is widely distributed and has a very wide range of hosts, populations, and damages, appear to be limited to some relatively small areas.

Major damages are observed in vegetables (particularly tomato and pepper), potato, grapevine, and soybeans. They are managed by conventional nematicides controlling soil populations, crop rotations, and resistant varieties.

Potatoes present a particular challenge because the nematodes parasitise the tubers and can thus be transported to new sites when contaminated tubers are used for seed. Because of the high reproductive potential of these nematodes and their polyphagous behaviour, planting pieces of infested tubers will create a serious problem to soils and future crops (Chaves & Torres, 1993; Doucet & Ponce de León, 1996a). The legislation prohibits the use of infested potato seed, which helps preventing the dispersal of these nematodes.

#### TYLENCHULUS SEMIPENETRANS

*Tylenchulus semipenetrans* was found for the first time in the North-East of the country on roots of citrus trees (Fresa, 1943). Later, it was found associated with different varieties of citrus in several regions, mainly the North-West and the North-East (Marchionatto, 1945, 1946, 1947; Schultz, 1945; Gutierrez, 1947; López Cristóbal, 1965; Moreno, 1969). In all cases, the nematode was linked to a decrease in citrus yields. The distribution of this nematode is very limited. The problems it causes are managed by resistant graft (Doucet & Ponce de León, 1992; Costilla, pers. comm.).

#### FAMILY LONGIDORIDAE

So far, three genera have been detected: *Xiphinema*, *Xiphidorus*, and *Paraxiphidorus*. The first genus has a low diversity; only seven species are present in Argentina. On the contrary, all the known species of the second genus are represented in the country to the exception of *X. uruguayensis*. The third genus includes only the type species (Coomans & Chaves, 1995).

Regarding the relative importance of these genera, it is necessary to mention the following species:

#### *Xiphinema index*

*Xiphinema index* was first found in the West on grapevines infested with grape-vine fan leaf virus and grape yellow mosaic virus (Feldman & Pontis, 1963, 1964). This species is widely distributed in the wine-growing region of the Province of Mendoza (La Red & Vega, 1968; Cucchi *et al.*, 1971; Vega, 1971, 1978). It was also found in grapevine plantations in the central part of the country – without any virus-related diseases – and on citrus roots in the North-East (Luc & Doucet, 1990).

The damages caused by this species on grapevines were controlled by nematicides (La Red & Vega, 1970).

#### *Xiphinema americanum sensu lato*

*X. americanum sensu lato* is the species of the genus *Xiphinema* the most frequently found in the country. It is associated with a great variety of crops (tobacco, citrus, sugar cane, grapevine, barley, tomato, several fruit trees) as well as wild plants. It is widely distributed (Luc & Doucet, 1990). It is not known whether the populations of this species can transmit pathogenic viruses.

### Main genera and species of entomopathogenic nematodes recorded in Argentina

#### FAMILY MERMITHIDAE

From the results of the studies made so far, it can be said that the Mermithidae family is characterised by a great generic and specific diversity in the country. Fifteen genera have been observed: three terrestrial and twelve aquatic genera. Two of them were found for the first time in Argentina: *Ditremamermis* Camino & Poinar, 1989 and *Divisispiculimermis* Doucet, 1986. The best represented genera are *Mesomermis* and *Gastromermis*, both including mostly species parasites of simuliids.

So far, *Strelkovimermis spiculatus* – a parasite of mosquitoes – is the only species studied in the laboratory to obtain information on its physiology and modalities of infection (Camino, 1988b,c; Camino & García, 1991, 1992).

Among terrestrial mermithids, the most remarkable are the genus *Hexamermis* including the largest number of species and the species *Agamermis decaudata* because of its large geographic distribution (Camino *et al.*, 1986; Doucet & Cagnolo, 1995b).

#### FAMILY HETERORHABDITIDAE

The genus *Heterorhabditis* is represented by the species *H. argentiniensis* and *H. bacteriophora*. Only one

population of the first species has been observed (Stock, 1993a) whereas several populations of the second species have been found in various locations in the country (Doucet & Bertolotti, 1995; 1996).

In the laboratory, two *H. bacteriophora* populations from Córdoba have shown promises as efficient biological control agents against Lepidoptera and Coleoptera (Scherma & Rodriguez Mosquera, 1989; Doucet & Giayetto, 1994).

#### FAMILY STEINERNEMATIDAE

So far, six species of the genus *Steinernema* are known in Argentina. *S. feltiae* was found in the Santa Fe province (Stock, 1993b), *S. scapterici* in Buenos Aires (Stock, 1992), and *S. carpocapsae*, *S. glaseri*, *S. rara*, and *S. ritteri* were found in Córdoba.

Except for *S. glaseri*, which was observed in mountain forest soil, the *Steinernema* species were found in cultivated soils.

Two species, *S. carpocapsae* and *S. rara* could become good control agents of harmful Homoptera, Lepidoptera and Coleoptera (Doucet, 1981).

#### Conclusion

Our knowledge of the soil nematodes that are present in the country is still insufficient. Because of the great diversity of conditions, it is reasonable to suppose that there is a marked diversity of species. It may be necessary to confirm the previous identification of some of the species by means of detailed studies.

Concerning the species of plant-parasitic nematodes, it remains necessary to assess more accurately the impact they have on the crops they attack. Because of a lack of symptoms, the attack of these pathogens sometimes goes unnoticed or is ascribed to other agents.

In those cases in which the harmful association nematode-plant is confirmed, it is essential to obtain more information on the biology of these species. The knowledge of the type of behaviour and the characterisation of the different populations of particular species (particularly *Meloidogyne* spp. and *Nacobbus aberrans*) will allow the development of strategies aimed at ensuring plant production and preserving good soil conditions.

In Argentina, research remains aimed at the study of the major species of plant-parasitic nematodes associated with potato and soybean and with vegetables in general, and toward the development of different types of control measures.

In relation to entomopathogenic nematodes, the objective of the investigations is to increase our knowledge on the species present, on the aggressivity of different populations, and to evaluate their potentials as biological control agents.

#### Acknowledgements

The authors thank the Consejo de Investigaciones Científicas y Técnicas (CONICET), the Consejo de Investigaciones Científicas y Tecnológicas de la Provincia de Córdoba (CONICOR), the Secretaria de Ciencia y Técnica of the Universidad Nacional de Córdoba and the Comisión Administradora del Fondo de Promoción de la Tecnología Agropecuaria (CAFPTA) for their financial support that made it possible to obtain part of the information reviewed here. They are also grateful to the members of the: Sociedad Argentina de Nematología, Biblioteca del Ministerio de Agricultura de la Nación and Archivo Histórico de la Municipalidad de Córdoba, who collaborated in the search and compilation of bibliographic references. They also thank Dr. M. Luc for his valuable suggestions and comments.

#### References

- AGÜERO, E. J. A., ROJAS, M. A. & VATTUONE, E. M. (1984). El falso nematode del nudo, *Nacobbus aberrans* (Thorne, 1935) Thorne & Allen, 1944 en cultivos de pimiento del Valle de Santa María. *Subsecret. Asuntos rur. & Univ. nac. Catamarca*, 10 p.
- AHMAD, R. (1972). Studies on *Graphognathus leucoloma* (Boh) (Coleoptera: Curculionidae) and its natural enemies in the central provinces of Argentina. *Techn. Bull. Commonw. Inst. biol. Control*, 17: 19-28.
- ANON. (1929). La anguilulosis de la alfalfa. Una amenaza para alfalfares. *Agríc. moderna*, 4: 667-679.
- AVILA, E. R. & PUCCI, A. (1980). Complejo etiológico en relación con la pudrición mohosa del ajo. *Fitopatología*, 15: 32-34.
- BAEZ, J. R. (1925). Enfermedades de nuestros alfalfares. Los manchones. *Revta mensual, Buenos Aires*, 8: 29-30.
- BERG, C. (1898). Sobre los enemigos pequeños de la langosta peregrina *Schistocerca paranensis* (Burm.). *Comun. Museo nac. Buenos Aires*, 1: 25-30.
- BLANCHARD, E. E. (1930). Principales parásitos que dañan el cultivo de la alfalfa en la República Argentina. Anguilulosis del tallo. *Dir. gen. Agríc., Def. Agríc., Buenos Aires*, 645: 14.
- BRÜCHER, H. (1960). Über das Wildvorkommen des Nematoden *Heterodera rostochiensis* in Nord-Argentinien. *Naturwissenschaften*, 47: 21.
- BRÜCHER, H. (1961). Primer hallazgo de *Heterodera rostochiensis* Woll. sobre papas silvestres (*Solanum* sect. *tuberosarium*) en la República Argentina y su significación como plaga de las especies cultivadas. *Revta Fac. Cienc. agr. Univ. nac. Cuyo*, 8: 7-18.
- BRUGNONI, H. C. (1966). Sobre la presencia de *Ditylenchus radicum* parasitando a trigo *Triticum aestivum*. *Revta Fac. Agron., La Plata*, 26: 139-142.
- BRUGNONI, H. C. (1980). *Plagas forestales*. Buenos Aires, Ed. Hemisferio Sur, 216 p.
- BRUNER, L. (1898). *Primer informe de la comisión del comercio de Buenos Aires para la investigación de la langosta*. Buenos Aires. 41 p.

- BURBA, J. L. (1992). Estado de la producción de ajo en la Argentina. In: Gómez Riera, P. (Ed.) *Argentina frutihortícola 92.*, EEA Mendoza: 37-45.
- BURKART, A. (1934). Alfalfa inmune al nematodo del tallo. *Revta Arg. Agric.*, 1: 304-306.
- BURKART, A. (1935). Un nematode parásito de *Poa annua* en Buenos Aires. *Revta Arg. Agric.*, 2: 136-138.
- BURKART, A. (1937). La selección de alfalfa inmune al nematode del tallo (*Anguillulina dipsaci*). *Revta Arg. Agric.*, 4: 171-196.
- BURKART, A. (1943). El nematodo del tallo (*Anguillulina dipsaci*) en alfalfares de Salta. *Revta Arg. Agric.*, 10: 190.
- CAMINO, N. B. (1985a). Estudio de cuatro especies del género *Mesomermis* Daday, 1911 (Nematoda: Mermithidae) parásitas de larvas de simúlidos (Diptera: Simuliidae). *Revta Museo La Plata*, 14: 1-19.
- CAMINO, N. B. (1985b). Contribución al estudio de la Familia Mermithidae (Nematoda) parásita de larvas de dípteros acuáticos. II. *Octomyomermis albicans* sp. n. *Neotrópica*, 31: 83-87.
- CAMINO, N. B. (1985c). Contribución al estudio de la Familia Mermithidae (Nematoda) parásita de larvas de dípteros acuáticos. I. *Gastromermis vaginiferous* sp. n. *Neotrópica*, 31: 143-147.
- CAMINO, N. B. (1986). Observaciones sobre el desarrollo embrionario y el estado infectante de *Mesomermis subandina* Camino, 1985 (Nematoda: Mermithidae), parásito de larvas de simúlidos. *Spheniscus*, 4: 1-7.
- CAMINO, N. B. (1987). Two new species of the genus *Isomermis* Coman, 1953 (Nematoda: Mermithidae) parasitic on aquatic larvae of Diptera in Argentina. *Revta ibér. Parasit.*, 47: 153-158.
- CAMINO, N. B. (1988a). Description of *Octomermis bonaerensis* n. sp. (Mermithidae, nematoda) parasitizing larvae of *Simulium bonaerense* Coscaron & Wygodzinski (Simuliidae, Diptera) in Argentina. *Revta ibér. Parasit.*, 48: 183-186.
- CAMINO, N. B. (1988b). Efecto del parasitismo múltiple en la determinación del sexo de *Strelkovimermis spiculatus* Poinar & Camino, 1986 (Nematoda: Mermithidae) en larvas de *Culex pipiens fatigans* Wiedemann, 1828. *Iheringia*, 2: 93-97.
- CAMINO, N. B. (1988c). Crecimiento larval de *Strelkovimermis spiculatus* Poinar y Camino, 1986 (Nematoda: Mermithidae) en *Culex pipiens fatigans* Wiedemann, 1828, como hospedador alternativo. *Iheringia*, 2: 31-37.
- CAMINO, N. B. (1990). *Mesomermis delponteiana* sp. n. (Nematoda: Mermithidae) parásito de larvas de *Simulium delponteianum* Wigodzinsky (Diptera: Simuliidae) en Argentina. *Revta ibér. Parasit.*, 50: 273-276.
- CAMINO, N. B. (1991). *Octomyomermis arecoensis* n. sp. (Nematoda: Mermithidae) parasitizing midges (Diptera: Chironomidae) in Argentina, with some observations on its bionomics. *Revue Nématol.*, 14: 375-379.
- CAMINO, N. B. (1993). A new species of the genus *Bathymermis* Daday, 1911 (Nematoda: Mermithidae) parasitizing blackfly larvae (Diptera: Simuliidae) in Argentina. *Res. Rev. Parasit.*, 53: 125-127.
- CAMINO, N. B. & GARCÍA, J. J. (1991). Influence of salinity and pH on the parasitism of *Strelkovimermis spiculatus* Poinar and Camino, 1986 (Nematoda: Mermithidae) in larvae of *Culex pipiens* Wied (Diptera: Culicidae). *Neotrópica*, 37: 107-112.
- CAMINO, N. B. & GARCÍA, J. J. (1992). Algunos factores que afectan al parasitismo de *Strelkovimermis spiculatus* Poinar & Camino, 1986 (Nematoda: Mermithidae) in mosquitos (Diptera: Culicidae). *Neotrópica*, 38: 75-80.
- CAMINO, N. B., MIRALLES, D. A. B. & MARCHISSIO, S. L. (1986). Aportes al conocimiento de la fauna argentina de mermitidos. I. Sobre *Agamermis decaudata* Cobb, Steiner y Christie, 1926 (Nematoda). *Neotrópica*, 32: 67-70.
- CAMINO, N. B. & POINAR, G. O., JR. (1989). *Ditremamermis simuliae* gen. nov., sp. nov. (Nematoda: Mermithidae) a parasite of *Simulium bonaerense* Coscaron & Wygodzinski in Argentina. *Neotrópica*, 34: 93-97.
- CAMINO, N. B. & REBORDA, G. R. (1994). Biología de *Strelkovimermis spiculatus* Poinar y Camino 1986 (Nematoda: Mermithidae, parásito de mosquitos (Diptera: Simuliidae) en condiciones de laboratorio. *Neotrópica*, 40: 45-48.
- CAMINO, N. B. & STOCK, S. P. (1989). *Hexamermis hortensis* sp. n. (Nematoda: Mermithidae) parásita de larvas de *Spodoptera frugiperda* (Smith) Lepidoptera: Noctuidae) en Argentina. *Revta Ibér. Parasit.*, 49: 329-333.
- CAMINO, N. B. & STOCK, S. P. (1994). *Hexamermis macrostoma* n. sp. (Nematoda: Mermithidae) parasitizing the cricket *Grylodes laplatae* (Orthoptera: Gryllidae) in Argentina. *Fundam. appl. Nematol.*, 17: 397-399.
- CAP, E. J., DE OBSCHATKO, E. S., CASTRONOVO, A. J. P., MIRANDA, O. & SERIGNESE, A. (1993a). *Perfil tecnológico de la producción agropecuaria argentina. Vol. I: Análisis descriptivo y prospectivo*. Buenos Aires, Inst. nac. Tecnol. agropec., 182 p.
- CAP, G. B., ROBERTS, P. A. & THOMASON, I. J. (1993b). Sources of resistance in *Lycopersicon* to the false root-knot nematode *Nacobbus aberrans*. 25th ann. Meet. ONTA, Cochabamba, Bolivia: 12. [Abstr.]
- CASTELLANOS, S. J. & DEL TORO, M. S. (1994). Evaluación de la resistencia de patrones de vid a *Meloidogyne incognita* (Kofoid & White) Chitwood en Mendoza, Argentina. *Nematropica*, 24: 75. [Abstr.]
- CHAVES, E. (1984). *Observations on plant parasitic nematodes from Argentina*. Thesis Doctorat, Univ. Gent, Belgium, 106 p.
- CHAVES, E. (1987). Cyst-nematodes (Heteroderidae) from Argentina. *Nematologica*, 33: 22-33.
- CHAVES, E. (1993). Revisión de los nematodos formadores de quistes en la República Argentina. *INTA, EEA Balcarce, Buenos Aires, Boln técn.*, 113: 1-11.
- CHAVES, E. & DE SISLER, G. M. (1980). Presencia de *Nacobbus aberrans* (Thorne, 1935) Thorne & Allen, 1944 (Nematodea: Nacobbiidae) en cultivos hortícolas de las provincias de Buenos Aires y Santa Fe, asociados con otros nematodos endoparásitos. *Inf. Invest. agropec.*, 385-386: 13-15.

- CHAVES, E. & TORRES, M. (1993). Nematodos parásitos de la papa del sudeste bonaerense. *INTA, EEA Balcarce, Buenos Aires, Boln técn.*, 115: 1-21.
- CHIOZZA, E. M. & VAN DOMSELAAR, Z. G. (1958). Clima. In: *La Argentina. Suma de geografía. Vol. II.* Buenos Aires, Argentina, Peuser Ed.: 92-183
- CONIL, P. A. (1878). Études sur l'*Acridium paranense* Burm., ses variétés et plusieurs insectes qui le détruisent. *Period. zool. Córdoba*, 3: 177-254.
- CONIL, P. A. (1879). Études sur l'*Acridium paranense* Burm., ses variétés et plusieurs insectes qui le détruisent. *Boln Acad. nac. Cienc. Córdoba*, 3: 385-469.
- COOMANS, A. & CHAVES, E. (1995). *Paraxiphidurus michel-luci* n. gen., n. sp. from Argentina (Nematoda: Longidoridae). *Fundam. appl. Nematol.*, 18: 303-306.
- COSTILLA, M. A. (1973). El nematodo *Pratylenchus zaeae* Graham en caña de azúcar. *Revta ind. agric. Tucumán*, 50: 39-43.
- COSTILLA, M. A. (1985). El falso nematode del nudo *Nacobbus aberrans* (Thorne, 1935) Thorne & Allen, 1944 y su relación con el cultivo de papa en el noroeste argentino. *Revta ind. agric. Tucumán*, 62: 79-97.
- COSTILLA, M. A. (1986). Comportamiento de diferentes poblaciones del falso nematode del nudo *Nacobbus aberrans* (Thorne, 1935) Thorne & Allen, 1944 en cultivos de papa. *6th Jorn. Fitosan. Arg., Neuquén, 8-11 April 1986*: 255-261.
- COSTILLA, M. A. (1991). Importancia de la frutilla (*Fragaria vesca*) en el manejo de *Nacobbus aberrans*, nematode del cultivo de tomate en Argentina. *23th Reun. anual ONTA, San Antonio, Texas, USA, 21-26 July 1991*: 123. [Abstr.]
- COSTILLA, M. A. (1992). Primer registro del nematodo de las lesiones *Pratylenchus zaeae* Graham, 1951 y su incidencia como plaga del maíz en la Argentina. *24th Reun. anual ONTA, Teguise, Lanzarote, España, 27 April-1 Mai 1992*: 120-121. [Abstr.]
- COSTILLA, M. A. & DE GOMEZ, T. A. H. (1981). Ensayo de control químico del nematode *Nacobbus aberrans* en tubérculos de papa. *Nematropica*, 11: 78-79.
- COSTILLA, M. A., DE GOMEZ, T. A. H. & DE OJEDA, S. G. (1976). Nematodos identificados en cultivos de caña de azúcar en las provincias del noroeste Argentino. *Revta ind. agric. Tucumán*, 53: 55-59.
- COSTILLA, M. A., DE OJEDA, S. G. & DE GÓMEZ, T. A. H. (1977). Contribución al estudio de "falso nematodo del nudo" *Nacobbus aberrans*. *Nematropica*, 7: 7-8.
- COSTILLA, M. A., DE OJEDA, S. G. & DE GÓMEZ, T. A. H. (1978). El falso nematode del nudo *Nacobbus aberrans* (Thorne, 1935) Thorne & Allen, 1944 (Nematoda, Nacobbiidae) en cultivos de papa de Tucumán. *3rd Jorn. fitosan. Arg., Tucumán, 6-8 Sept. 1978*: 323-340.
- COSTILLA, M. A., DE OJEDA, S. G. & DE GÓMEZ, T. A. H. (1979). *Helicotylenchus multicinctus* en raíces de banano en el noroeste de Argentina. *Nematropica*, 9: 138-139.
- CUCCHI, J. N. A., ORIOLANI, E. J., NADAL, J., URBIETA, A. & PUIATTI, A. (1971). Difusión del nematodo *Xiphinema index* Thorne & Allen en la zona centro, este y norte de la provincia de Mendoza. *Infivivo Investig. agropec.*, 283: 18-22.
- CUCCHI, N. J. A., PUIATTI, A. E. & SALVARREDI, A. U. (1967). Experiencias sobre el control de nematodos en ajo. *Infivivo Investig. agropec.*, 230: 54-58.
- DEL TORO, M. S. (1988a). Empleo de aldicarb en el control de nematodos fitoparásitos en viñedos de Mendoza y San Juan, Argentina. *Nematropica*, 18: 3.
- DEL TORO, M. S. (1988b). Utilización de orujo de uva para el control de nematodos en ajo. *Revta Fac. Cienc. agric. Mendoza*, 24: 25-34.
- DEL TORO, M. S., CASTELLANOS, S. J. & MOYANO, E. (1994a). Declaración de Mendoza, Argentina, como zona libre de "Nematodo dorado" en papa. *Nematropica*, 24: 93. [Abstr.]
- DEL TORO, M. S., CASTELLANOS, S. J. & MOYANO, E. (1994b). Estudio y certificación de "ajo semilla libre de nematodos" en Mendoza, Argentina. *Nematropica*, 24: 93 [Abstr.]
- DEL TORO, M. S., GARAY, J. A. & COLOMBINO, M. A. A. (1988). Experiencias para el control de *Ditylenchus dipsaci* (Kühn) Filipjev, en ajo, cv. Colorado, en San Luis, Argentina. *Revta Fac. Cienc. agric. Mendoza*, 24: 35-47.
- DEL TORO, M. S. & MAVRICH, E. (1977). Control del nematode *Ditylenchus dipsaci* (Kühn) Filipjev en semilla de ajo con termoterapia. *Revta Fac. Cienc. agric. Mendoza*, 21: 75-95.
- DEL TORO, M. S. & VEGA, E. (1982). Comparación entre el agregado de materia orgánica al suelo y el uso de nematocidas en el control de nematodos en tomate. *2nd Congr. Latinoam. Fitopatol., Buenos Aires, 22-26 November 1982*. [Abstr.]
- DOERING, A. (1891). Viticultura en Córdoba. *Boln Dept. nac. Agric., Buenos Aires*, 15: 381.
- DOUCET, M. E. (1986). Los nematodos fitófagos en Argentina: realidad y perspectivas. *6th Jorn. Fitosan. Arg., April 8-11 1986, Neuquén*: 243-246.
- DOUCET, M. E. (1989). The genus *Nacobbus* Thorne & Allen, 1944 in Argentina. 1. Study of a population of *N. aberrans* (Thorne, 1935) Thorne & Allen, 1944 on *Chenopodium album* L. from Rio Cuarto, province of Córdoba. *Revue Nématol.*, 12: 17-26.
- DOUCET, M. E. (1992). Asociación entre nematodos fitófagos y malezas en la República Argentina. *Agriscientia*, 9: 103-112.
- DOUCET, M. E. (1993). Consideraciones acerca del género *Meloidogyne* Goeldi, 1887 (Nemata: Tylenchida) y su situación en Argentina. *Agriscientia*, 10: 63-80.
- DOUCET, M. E. (1994). Variability on *Cruz nema tripartitum* (Linstow, 1906) Zullini, 1982 (Nemata: Rhabditida). *Stud. neotrop. Fauna Envir.*, 29: 33-41.
- DOUCET, M. E. & DI RIENZO, J. A. (1991). El género *Nacobbus* Thorne & Allen, 1944 en Argentina. 3. Caracterización morfológica y morfométrica de poblaciones de *N. aberrans* (Thorne, 1935) Thorne & Allen, 1944. *Nematropica*, 21: 19-35.
- DOUCET, M. E. & DE DOUCET, M. M. A. (1989). *Cruz nema tripartitum* (Linstow, 1906) Zullini, 1982: un

- nuevo nematodo entomófago facultativo. *7th Jorn. fitosan. Arg.*, 5-8 June 1989, Salta, Argentina. [Abstr.]
- DOUCET, M. E. & DE DOUCET, M. M. A. (1992). Contribution to the knowledge of *Distolabrellus veechi* Anderson, 1983 (Rhabditida: Nematoda). *Fundam. appl. Nematol.*, 15: 413-420.
- DOUCET, M. E. & DE DOUCET, M. M. A. (1996). Nematodos de suelo y agua dulce de la Provincia de Córdoba. In: Bucher, E. & di Tada, I. (Eds). *Biodiversidad de la Provincia de Córdoba. Vol. 1. Fauna*. Río Cuarto, Argentina, Univ. nac. Río Cuarto: 41-61.
- DOUCET, M. E. & GARDENAL, C. N. (1992). The genus *Nacobbus* in Argentina. 4. Preliminary comparison of populations of *N. aberrans* (Thorne, 1935) Thorne & Allen, 1944 by means of isoenzyme phenotypes. *Nematropica*, 22: 243-246.
- DOUCET, M. E., MONTAMAT, E. & GIAYETTO, A. (1996). Intrapopulation variability of enzyme phenotypes in *Nacobbus aberrans* from Argentina. *3rd int. Nematology Congr., Gosier, Guadalupe, 7-12 July 1996*: 144. [Abstr.]
- DOUCET, M. E. & PINOCHET, J. (1992). Occurrence of *Meloidogyne* spp. in Argentina. *J. Nematol.*, 24: 765-770.
- DOUCET, M. E. & DE PONCE DE LEÓN, E. L. (1986). *Chenopodium album* L.: eficiente hospedador de *Nacobbus aberrans* (Thorne, 1935) Thorne & Allen, 1944 y *Meloidogyne javanica* (Treub, 1885) Chitwood, 1949 en la Provincia de Córdoba. *Infivivo Investig. agropec.*, 437-440: 36-43.
- DOUCET, M. E. & DE PONCE DE LEÓN, E. L. (1992). *Tylenchulus semipenetrans* Cobb, 1913 (Nematoda: Tylenchida) y su asociación con *Citrus aurantium* L. y *Citrus reshni* Hort. ex Tan. *Agriscientia*, 9: 113-116.
- DOUCET, M. E. & DE PONCE DE LEÓN, E. L. (1994). Asociación de *Cruz nema tripartitum* con plántulas cloróticas de ajo. *Nematol. medit.*, 22: 149-151.
- DOUCET, M. E. & DE PONCE DE LEÓN, E. L. (1996a). *Meloidogyne* spp. (Nematoda): una seria amenaza para la papa *Solanum tuberosum* L. *Revta Investig. agropec.*, 26: 45-51.
- DOUCET, M. E. & DE PONCE DE LEÓN, E. L. (1996b). Alteraciones histológicas ocasionadas por *Meloidogyne incognita* (Kofoid & White, 1919) Chitwood, 1949 en raíces de plantas ornamentales florales. *Revta Investig. agropec.*, 26: 83-91.
- DOUCET, M. E. & DE PONCE DE LEÓN, E. L. & COSTILLA, M. A. (1992). Histopatología en *Capsicum annum* L. inducida por *Nacobbus aberrans* (Thorne, 1935) Thorne & Allen, 1944. *Revta Investig. agropec.*, 23: 69-76.
- DOUCET, M. M. A. DE (1981). *Neoalectana carpocapsae* Weiser (Steinernematidae) su posible utilización como agente en el control de *Zulia entrerriana* (Berg) (Cercopidae). *4th Jorn. fitosan. Arg. Córdoba, 19-21 August 1981*: 149-150. [Abstr.]
- DOUCET, M. M. A. DE (1982). Una nueva especie de *Gastromermis* Micoletzky, 1923 (Nematoda: Mermithidae) parásito de *Simulium wolffhengeli* (Enderlein) (Diptera: Simuliidae). *Revta Museo Arg. Cienc. nat. "Bernardino Rivadavia"*, 2: 11-17.
- DOUCET, M. M. A. DE (1986a). *Divisipiculimermis mirus* n. gen., n. sp., (Nematoda: Mermithidae) parasitizing midges in Córdoba (Argentina). *J. Nematol.*, 18: 247-251.
- DOUCET, M. M. A. DE (1986b). A new species of *Neoalectana* Steiner, 1929 (Nematoda: Steinernematidae) from Córdoba, Argentina. *Revue Nematol.*, 9: 317-323.
- DOUCET, M. M. A. DE (1992). Distribución de nematodos entomófagos en suelos de Córdoba, Argentina. *6th Congr. latinoam. Fitopat. & 6th Congr. nac. Soc. Esp. Fitopat. Torremolinos, España, 8-12 April 1992*: 112. [Abstr.]
- DOUCET, M. M. A. DE (1995). Caracterización de una población de *Steinernema carpocapsae* (Nematoda: Steinernematidae) aislada en Córdoba, República Argentina. *Nematol. medit.*, 23: 181-189.
- DOUCET, M. M. A. DE & BERLOTTI, M. A. (1995). New records of entomogenous nematodes (Steinernematidae; Heterorhabditidae) in Córdoba, Argentina. *Congr. int. Nematol. trop., Rio Quente, Brasil, 4-9 June 1995*: 74. [Abstr.]
- DOUCET, M. M. A. DE & BERLOTTI, M. A. (1996). Una nueva población de *Heterorhabditis bacteriophora* Poinar, 1975 de Argentina. Caracterización y acción sobre el huésped. *Nematol. medit.*, 24: 169-174.
- DOUCET, M. M. A. DE & CAGNOLO, S. (1995a). Diversity of mermithids in Córdoba, Argentina. *Congr. int. Nematol. trop., Rio Quente, Brasil, 4-9 June 1995*: 75. [Abstr.]
- DOUCET, M. M. A. DE & CAGNOLO, S. (1995b). *Agamer-mis decaudata* Cobb, Steiner y Christie, 1923 (Nematoda) parásito de isópodos y acrididos de Córdoba, Argentina. *9th Jorn. fitosan. Arg., Mendoza, 13-15 November 1995*: 110. [Abstr.]
- DOUCET, M. M. A. DE, CAGNOLO, S. R. & BERLOTTI, M. A. (1996). New *Heterorhabditis bacteriophora* Poinar, 1975 from Argentina. Variability and description of hermaphrodites from second and third generation. *Fundam. appl. Nematol.*, 19: 415-420.
- DOUCET, M. M. A. DE, CARO, K., MIRANDA, M. & BERLOTTI, M. A. (1996). Efficacy of *Heterorhabditis bacteriophora* (strain OLI) in relation to temperature, concentration and origin of the infective juvenile. *Nematropica*, 26: 129-133.
- DOUCET, M. M. A. DE & DOUCET, M. E. (1986). Nuevos datos para el conocimiento de *Heterorhabditis bacteriophora* Poinar, 1975. *Revta Investig. agropec.*, 21: 1-10.
- DOUCET, M. M. A. DE & DOUCET, M. E. (1989). *Cruz nema tripartitum* (Linstow, 1906) Zullini, 1982: un nuevo nematodo entomófago facultativo. *7th Jorn. fitosan. Arg., Salta, 5-8 June 1989*: 1 p. [Abstr.]
- DOUCET, M. M. A. DE & DOUCET, M. E. (1990). Description of *Steinernema ritteri* n. sp. (Nematoda: Steinernematidae) with a key to the species of the genus. *Nematologica*, 36: 257-265.
- DOUCET, M. M. A. DE & DOUCET, M. E. & BERLOTTI, M. A. (1991). Efecto de la temperatura de cría e intensidad de infestación sobre los caracteres morfológicos de *Heterorhabditis bacteriophora* Poinar, 1975. Análisis de variabilidad. *Nematropica*, 21: 37-49.
- DOUCET, M. M. A. DE, DOUCET, M. E. & DI RIENZO, J. (1990). Análisis de variabilidad de los caracteres morfo-

- métricos de *Heterorhabditis bacteriophora* Poinar, 1975. *Nematropica*, 20: 4-5.
- DOUCET, M. M. A. DE, DOUCET, M. E. & DI RIENZO, J. (1992a). Diferencias entre las larvas infectantes de *Heterorhabditis bacteriophora* Poinar, 1975 (Nemata: Heterorhabditidae) según su generación de origen. *Nematropica*, 22: 121-122.
- DOUCET, M. M. A. DE, DOUCET, M. E. & NIENSTEDT, K. M. (1992b). Diferencias inter e intraespecificas en la capacidad infectiva de poblaciones de *Heterorhabditis* y *Steinernema* aislados en Argentina. *Nematropica*, 22: 237-242.
- DOUCET, M. M. A. DE & GIAYETTO, A. (1994). Gama de huéspedes y especificidad de *Heterorhabditis bacteriophora* Poinar, 1975 (Nematoda: Heterorhabditidae). *Nematol. medit.*, 22: 171-178.
- DOUCET, M. M. A. DE & POINAR, G. O., JR. (1984). *Gastromermis kolleoni* n. sp. (Nematoda: Mermithidae) a parasite of midge (*Chironomus* sp., Chironomidae) from Argentina. *J. Nematol.*, 16: 252-255.
- DOUCET, M. M. A. DE & POINAR, G. O., JR. (1985). Estudio del ciclo de vida de una población de *Heterorhabditis* sp. proveniente de Río Cuarto, Provincia de Córdoba. *Revta Univ. nac. Rio Cuarto*, 5: 253-258.
- FELDMAN, J. M., & PONTIS, R. E. (1963). *Xiphinema index* y *Xiphinema americanum* en suelos de viñedos mendocinos. *Revta Fac. Cienc. agric. Mendoza*, 1-2: 3-12.
- FELDMAN, J. M., & PONTIS, R. E. (1964). *Xiphinema index* in vineyard soils in Mendoza, Argentina. *Pl. Dis. Repr.*, 48: 373-374.
- FRESA, R. (1943). El nematode de las raicillas de los cítricos. *Suelo Argent.*, 2: 695-696.
- GALMARINI, H. (1978). Nuevos cultivares de tomate. *2nd Reun. nac. Soc. argent. Olericult.*, 13-16 September 1978, Córdoba: 43. [Abstr.]
- GARGIULO, A. G. & MOYANO, A. J. (1948). Ensayo de DD en tomate. *Revta Fac. Cienc. agric. Mendoza*, 7: 39-47.
- GODECK, W. & FAVRET, E. A. (1965). Observaciones sobre la selección de alfalfa resistente al nematode del tallo. *Revta Investig. agropec.*, 3: 41-54.
- GODECK, W. & STILINOVIC, A. (1966). Comparación y posibilidades de distintos métodos para mejorar el rendimiento agronómico de la alfalfa. *Infvivo Investig. agropec.*, 226: 59-64.
- GOMEZ RIERA, P. (1992). Indicadores macroeconómicos. In: Gómez Riera, P. (Ed.). *Argent. Frutihortic.* 92, EEA Mendoza: 11-36.
- GUTIERREZ, R. O. (1945). Observaciones sobre Mermithidae de Acrididos. *Informe Inst. Sanid. veg., Min. Agric. nación.*: 352-364.
- GUTIERREZ, R. O. (1947). El nematode de las raicillas de los citrus *Tylenchulus semipenetrans* en la República Argentina. *Revta Investig. agric., Buenos Aires*, 1: 119-146.
- GUTIERREZ, R. O. (1949). Nuevo género y especie de nematodo saprobionte. *Min. Agric. Revta Investig. agric.*, 3: 403-412.
- GUTIERREZ, M. (1959). Inventario preliminar de la difusión de la "marchitez" del algodón en la Argentina. *Boln. INTA, EEA Presidencia Roque Saenz Peña*, 2: 1-96.
- HUERGO, J. M. (1902). Enfermedad radicular del tomate. *Boln Min. Agric., Buenos Aires*, 42: 1040-1059.
- HUERGO, J. M. (1903). Enfermedad radicular de la vid causada por *Heterodera radicola* o *Anguilulina radicola* de Greef (Anguillulosis). *Boln Agric. Ganad., Buenos Aires*, 61: 679-710.
- JATALA, P. & GOLDEN, A. M. (1977). Taxonomic status of *Nacobbus* species attacking potatoes in South-America. *Nematropica*, 7: 9-10.
- LA RED, F. C. & VEGA, E. (1968). Identificación y distribución geográfica del nematodo *Xiphinema index* Thorne & Allen en el sur de la provincia de Mendoza. *Infvivo Investig. agropec.*, 246: 49-54.
- LA RED, F. C. & VEGA, E. (1970). Control de nematodos en viñedos implantados. *Revta Investig. agropec.*, 7: 31-45.
- LÓPEZ CRISTÓBAL, U. (1946). La anguillulosis de la alfalfa en la República Argentina. *Publnes Min. Agric., Buenos Aires*, 20: 1-36.
- LÓPEZ CRISTÓBAL, U. (1965). Nematodos fitófagos. Anguillulosis de las plantas cultivadas en Argentina. *Agro*, 12: 31 p.
- LUC, M. & DOUCET, M. E. (1990). La Familia Longidoridae Thorne, 1935 (Nemata) en Argentina. 1. Distribución. *Revta Cienc. agropec.*, 7: 19-25.
- LYNCH ARRIBALZAGA, E. (1901). Trabajos de la extinguida sección entomológica. Informe de su ex-Director. V. Los Anguillulidos. *Boln Agric. Ganad.*, 1: 29-34.
- MALLO, R. G. (1961). Insectos, ácaros y nematodos enemigos del algodón en la República Argentina. *Infvivo Investig. agropec.*, 165: 10-22.
- MARCH, G. J., ORNAGHI, J. A., BEVIACQUA, J. E., ASTORGA, E. M. & MARCELLINO, J. (1985). Comportamiento de cultivares de soja frente al nematodo causante de agallas *Meloidogyne javanica* (Treub.) Chitwood. *Infvivo Investig. agropec.*, 441-444: 70-77.
- MARCHIONATTO, J. B. (1945). *La podredumbre de la raicilla del naranjo*. Buenos Aires, Argentina, Inst. San. Veg., Min. Agric., 15 p.
- MARCHIONATTO, J. B. (1946). Nota relacionada con la etiología de la podredumbre de la raicilla del naranjo. *Revta argent. Agron.*, 13: 96-100.
- MARCHIONATTO, J. B. (1947). La podredumbre de las raicillas de los citrus provocada por *Tylenchulus semipenetrans*. *Min. Agric., Inst. Sanid. veg., Buenos Aires, Serie A*, 3: 1-6.
- MARCHIONATTO, J. B., BAZZI, R. & CASTAÑEDA VEGA, R. (1926). Encuesta sobre alfalfares. *Minist. Agric., Secc. Propag. Inform.*, 634: 14.
- MAREGGIANI, G. & RUSSO, S. (1992). Nematodes asociados con ornamentales en Buenos Aires y sus alrededores. *Revta Fac. Agron., Buenos Aires*, 13: 145-150.
- MICCIO PERALTA, L. B. (1956). Alfalfa resistente al nematodo del tallo. *Infvivo Investig. agropec.*, 106-108: 58-59.
- MIRALLES, D. A. B. DE (1982). Presence of the genus *Pseudononymus* (Nematoda: Thelastomatidae) in the Republic of Argentina. *Neotropica*, 28: 170.
- MORENO, A. F. (1956a). Nematodo en primula. *Infvivo Investig. agropec.*, 106-108: 57-58.

- MORENO, A. F. (1956b). Un nematodo hallado en cultivos de ajos interfiere su exportación. *Infivivo Investig. agropec.*, 106-108: 58.
- MORENO, A. F. (1958). Algunos nematodos parásitos que perjudican cultivos de interés económico. *An. Soc. cient. Arg.*, 166: 115-121.
- MORENO, A. F. (1961). Nematodos que perjudican cultivos de interés económico. *Infivivo Investig. agropec.*, 163: 30-32.
- MORENO, A. F. (1964). The present state of nematology in Argentine. *Nematologica*, 10: 68.
- MORENO, A. F. (1969). Nematodos hallados en raíz y tierra de plantas cítricas. *CNIA, Inst. Pat. veg., INTA Castelar, Buenos Aires, Hoja inform.*, N° 38, 2 p.
- MORENO, A. F. & FRESA, A. R. (1969). Causa de muerte del pasto llorón. *CNIA, Inst. Pat. veg., INTA, Buenos Aires, Hoja inform.*, N° 36, 2 p.
- MORENO, A. F. & TURICA, A. (1960). Contribución al conocimiento de los nematodos perjudiciales para forestales. *Infivivo Investig. agropec.*, 147: 27-31.
- MOSCATELLI, G. (1990). *Atlas de suelos de la República Argentina. Vol. I.* Buenos Aires, Argentina, Secret. Agric., Ganad. Pesca., Inst. nac. Tecnol. agropec., Centro Investig. Recursos naturales, 731 p.
- OJEDA, S. DE, COSTILLA, M. A. & DE GOMEZ, T. H. (1978). Nematodes identificados en cultivos de papa de la provincia de Tucumán. *Revta Ind. Agric. Tucumán*, 55: 65-69.
- PAPADAKIS, J. (1964). Reseña ecológica de la Argentina. In: Parodi, R. (Ed.). *Enciclopedia argentina de agricultura y jardinería. Vol. II.* Buenos Aires, Argentina, ACME: 31-45.
- POINAR, G. O., JR., MRÁČEK, Z. & DE DOUCET, M. M. A. (1988). A re-examination of *Neoaplectana rara* Doucet, 1986 (Steinernematidae: Rhabditoidea). *Revue Nématol.*, 11: 447-449.
- DE PONCE DE LEÓN, E. L. & DOUCET, M. E. (1989). The genus *Nacobbus* Thorne & Allen, 1944 in Argentina. 2. Association between *N. aberrans* (Thorne, 1935) Thorne & Allen, 1944 and the weed *Sisymbrium irio* L. *Revue Nématol.*, 12: 269-271.
- PUCCI, A. & AVILA, E. R. (1978). Tomates resistentes a *Meloidogyne incognita* (Kofoid & White, 1919) Chitwood, 1949. *3rd Jorn. fitosan. Arg., Tucumán, 6-8 September 1978*, 1: 341-361.
- RAGONESE, A. E. & MARCÓ, P. R. (1943). Resistencia al nematodo del tallo de diversas líneas y procedencias de alfalfa. *Revta argent. Agric.*, 10: 378-384.
- RIVERA, J. C. & DEL TORO, M. S. (1982). Control químico de nematodos en ajo *Allium sativum* L. cv. Colorado en Mendoza, Argentina. *Inf. Investig. agropec.*, 401-404: 62-82.
- SCHERMA, J. C. & RODRIGUEZ MOSQUERA, M. (1989). *Heterorhabditis bacteriophora* (Poinar), potencial agente para el control de los gorgojos de la alfalfa. Aspectos preliminares. *7th Jorn. fitosan. Arg., Salta, 5-8 June 1978*: 1 p. [Abstr.]
- SCHULTZ, E. F. (1945). Algunas observaciones sobre la podredumbre de las raicillas del naranjo agrio injertado. *Boln Estac. exp. agric. Tucumán*, 54: 1-22.
- SHAHINA, F. & DE LEY, P. (1997). Two new species of Cephalobidae from Valle de la Luna, Argentina, and observations on the genera *Acrobeles* and *Nothacrobeles* (Nematoda: Rhabditida). *Fundam. appl. Nematol.*, 20: 329-348.
- DE SISLER, G. M. & DE CASAURANG, A. P. (1983). Reacción de cultivares de tomate y pimiento a *Nacobbus aberrans* (Nematoda, Nacobbidae). *Revta Fac. Agron. Buenos Aires*, 4: 79-82.
- STOCK, S. P. (1992). Presence of *Steinernema scapterisci* Nguyen & Swart parasitizing the mole cricket *Scapteriscus borellii* in Argentina. *Nematol. medit.*, 20: 163-165.
- STOCK, S. P. (1993a). A new species of the genus *Heterorhabditis* Poinar, 1976 (Nematoda: Heterorhabditidae) parasitizing *Graphognatus* sp. larvae (Coleoptera: Curculionidae) from Argentina. *Res. Rev. Parasit.*, 53: 103-107.
- STOCK, S. P. (1993b). Description of an Argentinian strain of *Steinernema feltiae* (Filipjev, 1934) (Nematoda: Steinernematidae). *Nematol. medit.*, 21: 279-283.
- STOCK, S. P. (1993c). *Micoletzkyia vidalae* n. sp. (Nematoda: Diplogasteridae), a facultative parasite of *Diabrotica speciosa* (Coleoptera: Chrysomelidae) from Argentina. *Res. Rev. Parasit.*, 53: 109-112.
- STOCK, S. P. & CAMINO, N. B. (1991). *Pellioiditis pellioidis* (Schneider) (Nematoda: Rhabditidae) parasitizing *Scaptia* (*Scaptia*) *lata* (Guerin-Meneville) (Diptera: Tabanidae). *Mems Inst. Osvaldo Cruz*, 86: 219-222.
- STOCK, S. P. & CAMINO, N. B. (1992a). *Hexameris ovistriata* n. sp. (Nematoda - Mermithidae) a parasite of the grasshopper *Staurorhectus longicornis* Giglio-Tos (Orthoptera: Acridiidae) in Argentina. *Fundam. appl. Nematol.*, 15: 15-18.
- STOCK, S. P. & CAMINO, N. B. (1992b). *Hexameris cochlearius* n. sp. (Nematoda-Mermithidae) a parasite of *Dichroplus elongatus* Giglio-Tos (Orthoptera: Acridiidae) in Argentina. *Nematol. medit.*, 20: 167-169.
- URBIETA SALVARREDI, A. (1972). Sintomatología del ataque del nematodo *Ditylenchus dipsaci* (Kühn) Goodey y su distribución en la planta de ajo. *Revta Investig. agropec.*, 9: 1-14.
- URBIETA SALVARREDI, A., CUCCHI, N. & TORRIGLIA, I. (1971). Nuevas experiencias sobre tratamientos químicos de ajo para controlar nematodos. *Infivivo Investig. agropec.*, 287: 87-96.
- VEGA, E. (1971). Review of the nematological problems that affect the central-western zone of Argentina. *Nematropica*, 1: 17-18.
- VEGA, E. (1978). Replantation des vignobles et désinfection des sols en Argentine. *Bull. O.I.V.*, 51: 250-262.
- VEGA, E. (1981). Rotación de cultivo para tomate y pimiento. Efecto sobre la población de nematodos del suelo. *4th Jorn. Fitosan. Arg., Córdoba, 19-21 August 1981*: 117-118. [Abstr.]
- VEGA, E. & GATICA, J. (1968). Variedad de tomate resistente a *Meloidogyne incognita* y control químico del parásito. *Infivivo Investig. agropec.*, 245: 60-64.
- VEGA, E. & GATICA, J. (1972). Ensayo de control químico de nematodos y fertilización en vivero de durazneros. *Infivivo Investig. agropec.*, 290: 9-14.

VEGA, E., MOLINA, M. N., CASTRO, O. J. & GARCÍA, L. A. (1986). Resultados obtenidos en el control de nematodes en papa semilla producida en Malargüe (Mendoza). *6th Jorn. fitosan. Arg., Neuquén, 8-11 April 1986*: 122. [Abstr.]  
 VIRSOO, E. V. (1967). Comportamiento de especies silvestres tuberíferas y de formas autóctonas del género *Solanum* en las montañas de Catamarca. *Infivto Investig. agropec.*, 231: 16-31.

WEYENBERGH, D. H. (1876). *Gordius acridiorum* Weyenbergh. *Informe Dept. nac. Agric. Rep. Argentina, año 1875*: 466.

WEYENBERGH, D. H. (1878). Descripciones de nuevos gusanos. *Periód. zool. Córdoba*, 3: 106-111.

WEYENBERGH, D. H. (1879). Descripciones de nuevos gusanos. *Boln Acad. nac. Cienc. Córdoba*, 3: 212-218.

## Appendix

List of soil nematodes reported in continental Argentina (doubtful records are marked with an asterisk [\*]).

### Class Adenophorea

#### Subclass Enoplia

#### Order Enoplida

##### Family Tripylidae

*Trischistoma monohysteroides* Altherr, 1963  
*Trischistoma monohystera* (de Man, 1880) Schuurmans Stekhoven, 1951  
*Tripyla* sp.  
*Tobrilus macrospiculum* Altherr, 1963  
*Tobrilus papillicaudatus* Altherr, 1963  
*Tobrilus setosus* Altherr, 1963  
*Tobrilus* sp.

##### Family Ironidae

*Ironus filiformis* Altherr, 1963

#### Order Mononchida

##### Family Cobbonchidae

*Cobbonchus regulus* Altherr, 1963

##### Family Mononchidae

*Clarkus papillatus* (Bastian, 1865) Jairajpuri, 1970  
*Coomansus gerlachei* (de Man, 1904) Jairajpuri & Khan, 1977  
*Mononchus aquaticus* Coetzee, 1968  
*Mononchus bellus* Andrassy, 1985  
*Mononchus* sp.  
*Prionchulus muscorum* (Dujardin, 1845) Wu & Hoeppli, 1929

##### Family Mylonchulidae

*Mylonchulus brachyuris* (Bütschli, 1873) Altherr, 1953  
*Mylonchulus hawaiiensis* (Cassidy, 1931) Andrassy, 1958  
*Mylonchulus minor* (Cobb, 1893) Andrassy, 1958  
*Mylonchulus parabrachyurus* (Thorne, 1924) Andrassy, 1958  
*Mylonchulus sigmaturus* (Cobb, 1917) Altherr, 1953  
*Mylonchulus* sp.  
*Sporonchulus ibitiensis* (Carvalho, 1951) Andrassy, 1958

##### Family Iotonchidae

*Iotonchus arenicola* Altherr, 1963  
*Iotonchus geminis* Heyns & Lagerwey, 1965  
*Iotonchus monhystera* (Cobb, 1917) Jairajpuri, 1970

### Order Mermithida

#### Family Mermithidae

*Agameremis decaudata* Cobb, Steiner & Christie, 1926  
*Amphimeremis bonariensis* Miralles & Camino, 1983  
*Bathymermis simuliae* Camino, 1993  
*Ditremameremis simuliae* Camino & Poinar, 1989  
*Divisipiculimeremis mirus* Doucet, 1986  
*Gastromermis cordobensis* Camino, 1991  
*Gastromermis doloresi* Camino, 1993  
*Gastromermis fidelis* Doucet, 1982  
*Gastromermis kolleonis* Doucet & Poinar, 1984  
*Gastromermis vaginiferous* Camino, 1985  
*Hexameremis cochlearius* Stock & Camino, 1992  
*Hexameremis hortensis* Camino & Stock, 1989  
*Hexameremis macrostoma* Camino & Stock, 1994  
*Hexameremis ovistriata* Stock & Camino, 1992  
*Hydromermis* sp.  
*Hydromermis doloresi* Camino, 1993  
*Isomeremis* sp.  
*Isomeremis sierrensis* Camino, 1994  
*Limnomermis bonariensis* Camino, 1989  
*Limnomermis* sp.  
*Mermis* sp.  
*Mesomeremis crassivaginae* Camino, 1985  
*Mesomeremis dissimilis* Camino, 1985  
*Mesomeremis delponteiana* Camino, 1990  
*Mesomeremis nortensis* Camino, 1991  
*Mesomeremis ochrae* Camino, 1985  
*Mesomeremis subandina* Camino, 1985  
*Mesomeremis* sp.  
*Octomyomeremis albicans* Camino, 1985  
*Octomyomeremis arecoensis* Camino, 1991  
*Octomyomeremis bonariensis* Camino, 1988  
*Octomyomeremis longispiculae* Camino, 1992  
*Pseudomeremis* sp.  
*Strelkovimeremis spiculatus* Poinar & Camino, 1986  
*Strelkovimeremis* sp.

### Order Dorylaimida

#### Family Actinolaimidae

*Actinolaimus* sp.

#### Family Aporcelaimidae Heyns, 1965

*Aporcelaimellus obscurus* (Thorne & Swanger, 1936) Heyns, 1965



- Aporcelaimellus* sp.  
*Aporcelaimus* sp.
- Family Belonidiridae  
*Dorylaimellus monticolus* Clark, 1963  
*Dorylaimellus virginianus* Cobb, 1913
- Family Charcharolaimidae  
*Carcharolaimus formosus* Lordello, 1957  
*Carcharolaimus rotundicauda* (de Man, 1880) Thorne, 1939  
*Carcharolaimus* sp.
- Family Leptonchidae  
*Leptonchus granulatus* Cobb, 1920  
*Leptonchus* sp.  
*Tylencholaimellus montanus* Thorne, 1939
- Family Longidoridae  
*Paraxiphidorus michelluci* Coomans & Chaves, 1995  
*Paraxiphidorus* sp.  
*Xiphidorus achalae* Luc & Doucet, 1984  
*Xiphidorus balcarceanus* Chaves & Coomans, 1984  
*Xiphidorus tucumanensis* Chaves & Coomans, 1984  
*Xiphidorus saladillensis* Chaves & Coomans, 1984  
*Xiphidorus yepesara* Monteiro, 1976  
*Xiphidorus* sp.  
*Xiphinema americanum* Cobb, 1913  
*Xiphinema diversicaudatum* (Micoletzky, 1927) Thorne, 1939  
*Xiphinema index* Thorne & Allen, 1950  
*Xiphinema krugi* Lordello, 1955  
*Xiphinema setariae* Luc, 1958  
*Xiphinema rivesi* Dalmasso, 1969  
*Xiphinema surinamense* Loof & Maas, 1972  
*Xiphinema* sp.
- Family Mydonomidae  
*Dorylaimoides elongatus* Husain & Khan, 1968
- Family Nordiidae  
*Pungentus monhystera* Thorne & Swanger, 1936
- Family Dorylaimidae  
*Dorylaimus* sp.  
*Mesodorylaimus aberrans* Loof, 1969  
*Mesodorylaimus adalberti* Andrásy, 1963  
*Mesodorylaimus argentinus* Altherr, 1963  
*Mesodorylaimus dorni* Loof, 1969  
*Mesodorylaimus meridianus* Andrásy, 1963  
*Mesodorylaimus pseudobastiani* Loof, 1969  
*Mesodorylaimus puellae* Andrásy, 1963  
*Mesodorylaimus szekessyi* Andrásy, 1960  
*Mesodorylaimus* sp.
- Family Nygolaimidae  
*Nygolaimus* sp.
- Family Qudsianematidae  
*Eudorylaimus monhystera* (de Man, 1880) Andrásy, 1959
- Eudorylaimus obtusicaudatus* (Bastian, 1865) Andrásy, 1959  
*Eudorylaimus profestus* Andrásy, 1963  
*Eudorylaimus quadramphidius* Andrásy, 1963  
*Eudorylaimus* sp.  
*Labronema* sp.
- Order Triplonchida**
- Family Diphterophoridae  
*Diphterophora communis* de Man, 1880  
*Diphterophora* sp.
- Family Trichodoridae  
*Paratrichodorus minor* (Colbran, 1956) Siddiqi, 1974  
*Paratrichodorus* sp.  
*Trichodorus obscurus* Allen, 1957  
*Trichodorus* sp.
- Subclass Chromadoria**
- Order Araeolaimida**
- Family Plectidae  
*Anaplectus* sp.  
*Plectus (Plectoides) patagonicus* de Man, 1904  
*Plectus* sp.
- Order Chromadorida**
- Family Cyatholaimidae  
*Odontolaimus aquaticus* W. Schneider, 1937
- Order Monhysterida**
- Family Monhysteridae  
*Monhystera lepidura* Andrásy, 1963  
*Monhystera* sp.
- Class Secernentea**
- Subclass Rhabditida**
- Order Rhabditida**
- Family Cephalobidae  
*Acrobeles* sp.  
*Acrobeles emmatus* Shahina & De Ley, 1997  
*Acrobeloides* sp.  
*Cephalobus* sp.  
*Cervidellus* sp.  
*Eucephalobus* sp.  
*Nothacrobeles lunensis* Shahina & De Ley, 1997  
*Zeldia* sp.
- Family Rhabditidae  
*Cephaloboides* sp.  
*Cruznema tripartitum* (Linstow, 1906) Zullini, 1982  
*Cruznema* sp.  
*Distolabrellus veechi* Anderson, 1983  
*Oxycerca brevispina* (de Man, 1895) Andrásy, 1983  
*Pellioditis pellio* (Schneider, 1866) Timm, 1960

- Rhabditis* spp.
- Family Heterorhabditidae
- Heterorhabditis argentiensis* Stock, 1993  
*Heterorhabditis bacteriophora* Poinar, 1975  
*Heterorhabditis* sp.
- Family Steinernematidae
- Steinernema carpocapsae* (Weiser, 1955) Poinar, 1990  
*Steinernema feltiae* (Filipjev, 1934) Poinar, 1990  
*Steinernema glaseri* (Steiner, 1929) Poinar, 1990  
*Steinernema rarum* (Doucet, 1986) Poinar, 1990  
*Steinernema ritteri* Doucet & Doucet, 1990  
*Steinernema scapterisci* Nguyen & Smart, 1991  
*Steinernema* sp.
- Family Panagrolaimidae
- Panagrolaimus* sp.
- Order Ascaridida**
- Family Thelastomatidae
- Leidynema appendiculatum* (Leidy, 1890) Chitwood, 1932  
*Pseudonymus* sp.  
*Thelastoma dessetae* Adamson, 1985
- Order Diplogasterida**
- Family Diplogastridae
- Micoletzkyia vidalae* Stock, 1993
- Order Aphelenchida**
- Family Aphelenchidae
- Aphelenchus avenae* Bastian, 1865  
*Aphelenchus* sp.
- Family Aphelenchoididae
- Aphelenchoides bicaudatus* (Imamura, 1931) Filipjev & Schuurmans Stekhoven, 1941  
*Aphelenchoides blastophthorus* Franklin, 1952  
*Aphelenchoides fragariae* (Ritzema Bos, 1890) Christie, 1932  
*Aphelenchoides parietinus* (Bastian, 1865) Steiner, 1932  
*Aphelenchoides ritzemabosi* (Schwartz, 1911) Steiner, 1932  
*Aphelenchoides* sp.  
*Laimaphelenchus cocucci* Doucet, 1992  
*Laimaphelenchus* sp.  
*Seinura* sp.
- Family Ektaphelenchidae
- Ektaphelenchus tenuidens* (Thorne, 1935) Rühm, 1956
- Order Tylenchida**
- Family Tylenchidae
- Atylenchus* sp.  
*Atylenchus decalineatus* Cobb, 1913  
*Boleodorus* sp.
- Cephalenchus* sp.  
*Coslenchus* sp.  
*Filenchus* sp.  
*Lelenchus* sp.  
*Psilenchus bahiablancae* Doucet, 1996  
*Psilenchus hilarus* Siddiqi, 1963  
*Psilenchus pratensis* Doucet, 1996  
*Psilenchus* sp.  
*Tetylenchus* sp.  
*Tylenchus* sp.  
*Tylenchus davanei* Bastian, 1985  
*Tylenchus filiformis* Micoletzky, 1922
- Family Anguinidae
- Anguina agrostis* (Steinbuch, 1799) Filipjev, 1936  
*Anguina* sp.  
*Ditylenchus dipsaci* (Kühn, 1857) Filipjev, 1936  
*Ditylenchus intermedius* (de Man, 1880) Filipjev, 1936  
*Ditylenchus* sp.  
*Orrina phyllobia* (Thorne, 1934) Thorne, 1961
- Family Belonolaimidae
- Belonolaimus gracilis* Steiner, 1949 \*  
*Belonolaimus brevicaudatus* (Doucet, 1983) Fortuner & Luc, 1987  
*Tetylenchus* sp.  
*Triversus festonatus* (Doucet, 1985) Fortuner & Luc, 1987  
*Tylenchorhynchus acti* Hooper, 1959  
*Tylenchorhynchus annulatus* (Cassidy, 1930) Golden, 1971  
*Tylenchorhynchus capitatus* Allen, 1955  
*Tylenchorhynchus cylindricus* Cobb, 1913  
*Tylenchorhynchus dubius* (Bütschli, 1837) Filipjev, 1936  
*Tylenchorhynchus* sp.
- Family Dolichodoridae
- Dolichodorus aquaticus* Doucet, 1985  
*Dolichodorus heterocephalus* Cobb, 1914 \*  
*Dolichodorus longicaudatus* Doucet, 1981  
*Neodolichodorus leiocephalus* Doucet, 1981
- Family Pratylenchidae
- Pratylenchus agilis* Thorne & Malek, 1968  
*Pratylenchus brachyurus* (Godfrey, 1929) Filipjev & Schuurmans Stekhoven, 1941  
*Pratylenchus delattrei* Luc, 1958  
*Pratylenchus goodeyi* Oostenbrink, 1953  
*Pratylenchus hexincisus* Taylor & Jenkins, 1957  
*Pratylenchus microstylus* Bajaj & Bahatti, 1984  
*Pratylenchus neglectus* (Rensch, 1924) Filipjev & Schuurmans Stekhoven, 1941  
*Pratylenchus penetrans* (Cobb, 1917) Filipjev & Schuurmans Stekhoven, 1941  
*Pratylenchus pratensis* Meyl, 1954  
*Pratylenchus pseudopratensis* Seinhorst, 1968  
*Pratylenchus scribneri* Steiner, in Sherbakoff & Stanley, 1943  
*Pratylenchus thornei* Sher & Allen, 1953  
*Pratylenchus vulnus* Allen & Jensen, 1951  
*Pratylenchus zaeae* Graham, 1951

*Pratylenchus* sp.  
*Radopholus similis* (Cobb, 1893) Thorne, 1949  
*Nacobbus aberrans* (Thorne, 1935) Thorne & Allen, 1944

## Family Hoplolaimidae

*Aorolaimus longistylus* (Doucet, 1980) Fortuner, 1987  
*Aorolaimus perscitus* (Doucet, 1980) Fortuner, 1987  
*Aorolaimus conicori* (Doucet, 1984) Fortuner, 1987  
*Aorolaimus brevicaudatus* (Doucet, 1984) Fortuner, 1987  
*Aorolaimus triticeus* (Doucet, 1984) Fortuner, 1987  
*Aorolaimus* sp.  
*Helicotylenchus nannus* Steiner, 1945  
*Helicotylenchus digonicus* Perry, in Perry, Darling & Thorne, 1959  
*Helicotylenchus multicinctus* (Cobb, 1893) Golden, 1956  
*Helicotylenchus pseudorobustus* (Steiner, 1914) Golden, 1956  
*Helicotylenchus* sp.  
*Hoplolaimus galeatus* (Cobb, 1913) Thorne, 1935  
*Hoplolaimus* sp.  
*Rotylenchus buxophilus* Golden, 1956  
*Rotylenchus goodeyi* Loof & Oostenbrink, 1958  
*Rotylenchus robustus* (de Man, 1876) Filipjev, 1936  
*Rotylenchus uniformis* (Thorne, 1949) Loof & Oostenbrink, 1958  
*Rotylenchus* sp.

## Family Heteroderidae

*Cactodera amaranthi* (Stoyanov, 1972) Krall' & Krall', 1978  
*Cactodera* sp.  
*Globodera rostochiensis* (Wollenweber, 1923) Behrens, 1975 \*  
*Globodera tabacum tabacum* (Lownsberry & Lownsberry, 1954) Behrens, 1975  
*Globodera* sp.  
*Heterodera schachtii* Schmidt, 1871 \*  
*Heterodera galeopsidis* Goffart, 1936  
*Heterodera* spp.  
*Meloidogyne acrita* Chitwood, 1949  
*Meloidogyne arenaria* (Neal, 1889) Chitwood, 1949  
*Meloidogyne chitwoodi* Golden, O'Bannon, Santo & Finley, 1980  
*Meloidogyne cruciani* García-Martínez, Taylor & Smart, 1982  
*Meloidogyne decalineata* Whitehead, 1968  
*Meloidogyne hapla* Chitwood, 1949  
*Meloidogyne incognita* (Kofoid & White, 1919) Chitwood, 1949  
*Meloidogyne javanica* (Treub, 1885) Chitwood, 1949  
*Meloidogyne naasi* Franklin, 1965  
*Meloidogyne ottersoni* (Thorne, 1969) Franklin, 1971

*Meloidogyne* sp.

## Family Criconematidae

*Criconema duplicivestitum* (Andrássy, 1963) Raski & Luc, 1985  
*Criconema* sp.  
*Mesocriconema curvatum* (Raski, 1952) Loof & De Grisse, 1989  
*Mesocriconema douceti* (Siddiqi, 1986) Ebsary, 1991  
*Mesocriconema ornatum* (Raski, 1958) Loof & De Grisse, 1989  
*Mesocriconema peruensiforme* (De Grisse, 1967) Loof & De Grisse, 1989  
*Mesocriconema ritteri* (Doucet, 1980) Loof & De Grisse, 1989  
*Mesocriconema sphaerocephala* (Taylor, 1936) Loof & De Grisse, 1989  
*Mesocriconema talense* (Chaves, 1983) Loof & De Grisse, 1989  
*Mesocriconema xenoplax* (Raski, 1952) Loof & De Grisse, 1989  
*Mesocriconema* sp.  
*Nothocriconema mutabile* (Taylor, 1936) De Grisse & Loof, 1965  
*Nothocriconema* sp.  
*Caloosia paradoxa* (Luc, 1958) Brzeski, 1974  
*Hemicycliophora arenaria* Raski, 1958  
*Hemicycliophora filicauda* Doucet, 1982  
*Hemicycliophora fragilis* Doucet, 1982  
*Hemicycliophora gracilis* Thorne, 1955  
*Hemicycliophora penetrans* Thorne, 1955  
*Hemicycliophora poranga* Monteiro & Lordello, 1978  
*Hemicycliophora rara* Doucet, 1983  
*Hemicycliophora rionegrensis* Doucet, 1982  
*Hemicycliophora ripa* Van den Berg, 1981  
*Hemicycliophora tenuistriata* Doucet, 1982  
*Hemicycliophora thienemanni* (Schneider, 1925) Loos, 1948  
*Hemicycliophora zuckermani* Brzeski, 1963  
*Hemicycliophora* sp.  
*Ogma alternatum* (Doucet, 1986) Raski & Luc, 1987  
*Ogma comahuense* Brugni & Chaves, 1994  
*Ogma multisquamatum* (Kirjanova, 1948) Mehta & Raski, 1971

## Family Tylenchulidae

*Gracilacus colina* Huang & Raski, 1986  
*Paratylenchus neoamblycephalus* Geraert, 1965  
*Paratylenchus* sp.  
*Tylenchulus semipenetrans* Cobb, 1913

## Family Allantonematidae

*Metaparasytylenchus* sp.