



*incognita*, pointing to the need to search other variable regions of the genome to distinguish these species. While Blok *et al.* (1997) also found no size differences between the ITS-1 PCR products of *M. incognita*, *M. arenaria* and *M. javanica*, the PCR products of the less related *M. hapla* and *M. mayaguensis* were distinguishable from the other three species.

The intergenic spacer (IGS) has become another focal point in molecular diagnostics since this region of rDNA more easily yields polymorphisms between *Meloidogyne* species. Petersen and Vrain (1996) reversed the ITS primers of Vrain *et al.* (1992) to amplify a 5.8 kb fragment including the IGS of *M. chitwoodi*. They mapped, obtained partial nucleotide sequences, and designed new PCR primers to the 3' end of the 28S rRNA gene and to the 5' end of the 18S rRNA gene. These primers amplified the IGS region of *M. chitwoodi*, *M. hapla* and *M. fallax*, revealing distinct polymorphisms between these three species, and providing molecular evidence to confirm the status of the newly described *M. fallax* (Karssen, 1996).

A further advance in molecular identification of nematode populations is to synthesize oligonucleotide probes which specifically hybridize to a single species, but this requires species-specific DNA targets to be identified. Williamson *et al.* (1997) synthesized PCR primers derived from the sequence of cloned RAPD bands that distinguished single juveniles of *M. chitwoodi* from *M. hapla*. Zijlstra (1997) cloned and sequenced the ITS region of *M. chitwoodi*, *M. fallax*, *M. hapla*, and *M. incognita*, and identified a great deal more variability in the ITS2 region than in the ITS1. PCR primers were designed to the variable regions to separate these four species. The amplification product from *M. chitwoodi* was easily separated from *M. hapla* and *M. incognita*, but the size of the PCR product of *M. fallax* differed from that of *M. chitwoodi* by only 8 bp, making it difficult to distinguish between the two using standard gel electrophoresis.

Since selection pressure decreases mutation rates within protein coding DNA sequences, a higher degree of genetic variability is often found in non-coding regions. Indeed, diagnostic probes capable of distinguishing *Pratylenchus penetrans* and *P. scribneri* were designed from intron sequences in the major sperm protein (*msp*) gene (Setterquist *et al.*, 1996). The non-coding nature of the IGS, combined with size polymorphisms within PCR-amplified IGS fragments from closely related species (Petersen & Vrain, 1996), indicated a high potential for utilizing this region for producing species-specific probes. We have synthesized two probes, each targeting signature sequences within the IGS of either *M. chitwoodi* or *M. fallax*, and utilized them in multiplex PCR to distinguish and differentiate *Meloidogyne* species via a single-tube assay format.

## Materials and methods

### NEMATODES

The populations of *Meloidogyne* spp. used in this study are listed in Table 1. Genomic DNA was purified from nematode juveniles or eggs essentially as previously described (Vrain *et al.*, 1992; Zijlstra *et al.*, 1995).

### POLYMERASE CHAIN REACTION

PCR amplification of the entire IGS region utilized primers previously described (Petersen & Vrain, 1996). PCR reactions typically contained 3-5 ng genomic DNA template, 2 mM each dNTP, 1.5 mM MgCl<sub>2</sub>, 1 U *Taq* polymerase, and 1 X *Taq* buffer (60 mM Tris-SO<sub>4</sub> [pH 9.1], 15 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, Appligene-Oncor, France). Cycling parameters typically consisted of 95°C denaturation for 15 s, 60°C annealing for 30 s, and 72°C extension for 60 s. An initial denaturation step (95°C for 3 min) was performed prior to thirty amplification cycles, and a final extension step (72°C for 5 min) was performed immediately following amplification. PCR reactions utilizing the "pentaprimer" mix described below were conducted using an annealing temperature of 64°C.

### DNA SEQUENCING

Amplified fragments were separated from the genomic DNA template and primers by electrophoresis using 2% (w/v) low-melting point agarose (Gibco-BRL, Burlington, ON, Canada) gels, containing 0.25 µg/ml ethidium bromide (EtBr), in Tris-acetate EDTA buffer at 90 V for 1 h. Visualization of EtBr-stained DNA was accomplished using a UV transilluminator, and the size of each fragment calculated using the 100 bp marker (BRL) as a size standard. DNA bands were excised from gels with a sterile scalpel and purified from the agarose using Qiaquick spin columns (Qiagen Inc, Chatsworth, CA, USA), eluted in 10 mM Tris-HCl, pH 8.0, and stored frozen until further use. Sequencing was accomplished using the FSTaq Ready Reaction Kit (Perkin Elmer, Mississauga, ON, Canada) and analyzed by ABI310 capillary electrophoresis system according to the manufacturers directions. The sequencing data obtained for the IGS regions were aligned using Sequence Navigator so that additional primers could be designed to specifically target DNA from individual nematode species.

### OLIGONUCLEOTIDE SYNTHESIS

Cloned DNA fragments containing portions of the rDNA IGS (Petersen & Vrain, 1996) were sequenced and used as the basis for primer constructions. The DNA sequences of oligonucleotide primers used in

**Table 1.** Isolates and sources of species of *Meloidogyne* used in this study.

Code	Location	Isolate	Source	Pentaprimer bands <sup>a</sup>
<i>M. chitwoodi</i>				
Ca	The Netherlands	C3022	Pd <sup>b</sup>	C, I
Ck	The Netherlands	C5273g	PD	C, I
Cl	The Netherlands	C2960	PD	C, I
Co	The Netherlands	Horst	PAV <sup>c</sup>	C, I
Cba	Oregon, USA	ORMC12, race 1	USDA <sup>d</sup>	C, I
Cbd	Washington, USA	WAMC16, race 2	USDA	C, I
Cbf	Oregon, USA	ORMC8, race 2	USDA	C, I
Cbh	California, USA	CAMC2, race 3	USDA	C, I
Cbs	The Netherlands	IHO9	PAV	C, I
Cbu	The Netherlands	KBD4	PAV	C, I
Ccg	Portugal		INRA <sup>e</sup>	C, I
Cch	Argentina	CArg	INRA	C, I
<i>M. fallax</i>				
Fa	The Netherlands	CHB	PAV	F, I
Fb	The Netherlands	C4571	PD	F, I
Fc	The Netherlands	C6501	PD	F, I
Fd	The Netherlands	C6466	PD	F, I
Fe	The Netherlands	W834.769	BLG <sup>f</sup>	F, I
Ff	Belgium		Moens <sup>g</sup>	F, I
Fg	The Netherlands		CPRO-DLO <sup>h</sup>	F, I
Fh	France	CHK	INRA	F, I
<i>M. hapla</i>				
Hc	The Netherlands	C3064	PD	I
He	The Netherlands	Dr.91	PAV	I
Hh	The Netherlands	Wa.90 (raceB)	PAV	I
Hi	The Netherlands	Sm.Fer92	PAV	I
Hj	The Netherlands	Sm.Fer92	PAV	I
Hk	The Netherlands	Sl.92	PAV	I
Ham	Queensland, Australia	Q114	ORSTOM <sup>i</sup>	I
Han	South Korea	C2346	PD	I
Has	Washington, USA	WAMH2	USDA	I
Hau	Utah, USA	UTMH1	USDA	I
Hbj	Canada	Canada	Uw <sup>j</sup>	I
Hbq	Hungary	C6611	PD	I
Hbr	The Netherlands	C4900-H-1A	PD	I
Hbs	The Netherlands	C7388	PD	I
Hbt	The Netherlands	W834.347	BLGG	I
Hby	The Netherlands	C3029	PD	I
Hcb	The Netherlands	HBW	CPRO-DLO	I
L33	The Netherlands		SCRI <sup>k</sup>	I

End of Table 1 next page.

Table 1. (cont.)

Code	Location	Isolate	Source	Pentaprimer bands <sup>a</sup>
Pak1	Pakistan		SCRI	I
Cyp1	Cyprus		SCRI	I
<i>M. incognita</i>				
Ia	The Netherlands		IPO-DLO <sup>l</sup>	I
Ib	The Netherlands	Inc568-93	PD	I
If	The Netherlands	KnolE	De Ruiter <sup>m</sup>	I
Ih	The Netherlands	KnolG	De Ruiter	I
Ij	The Netherlands	KnolH	De Ruiter	I
I3			SCRI	I
L11	Louisiana		SCRI	I
L15	Thailand		SCRI	I
L17	Burkina Faso		ORSTOM	I
L19	French West Indies		ORSTOM	I
<i>M. arenaria</i>				
L10	Ivory Coast		ORSTOM	I
L26	Portugal		SCRI	I
L32	French West Indies		ORSTOM	I
<i>M. mayaguensis</i>				
L30	Burkina Faso		ORSTOM	I
L3	Ivory Coast		ORSTOM	I
<i>M. javanica</i>				
Ja	The Netherlands	C3059	PD	I
L25	Portugal		SCRI	I
L23	Burkina Faso		ORSTOM	I

<sup>a</sup> Amplified fragment sizes resulting from PCR using Pentaprimers are indicated as follows: 900 bp band = C (*chitwoodi*-specific), 1100 bp band = F (*fallax*-specific), 750 bp band = I (ITS-specific).

<sup>b</sup> Plant Protection Service, Wageningen, The Netherlands.

<sup>c</sup> Research Station for Arable Farming and Field Production of Vegetables, Lelystad, The Netherlands.

<sup>d</sup> United States Department of Agriculture, Prosser, Washington, USA.

<sup>e</sup> Institut National de Recherche Agronomique, Le Rheu, France.

<sup>f</sup> Laboratory for Soil and Crop Testing, Oosterbeek, The Netherlands.

<sup>g</sup> Centrum Landbouwkundige Onderzoek, Merelbeke, Belgium.

<sup>h</sup> Centre for Plant Breeding and Reproduction Research, Wageningen, The Netherlands.

<sup>i</sup> Office de la Recherche Scientifique et Technique Outre-Mer, Montpellier, France.

<sup>j</sup> Department of Plant Pathology, University of Wisconsin, Madison, WI, USA.

<sup>k</sup> Scottish Crop Research Institute, Invergowrie, Dundee, United Kingdom.


<sup>l</sup> DLO-Research Institute for Plant Protection, Wageningen, The Netherlands.

<sup>m</sup> De Ruiter Seeds, Bleiswijk, The Netherlands.

this study are listed in Table 2. Conserved primers included those designed to amplify the ITS (18S and 28S) and a conserved sequence primer which binds

within the IGS (1839). These three primers were mixed with primers specific for *M. chitwoodi* (C64) and *M. fallax* (F64) to produce the "pentaprimer"





**Fig. 2.** Agarose gel separation of species-specific amplicons. Multiplex PCR was conducted using the pentaprimer mix as described in Materials and Methods. Templates added to each reaction were either control DNAs (3 ng) from *Meloidogyne chitwoodi* (lane 2), *M. hapla* (lane 3), and *M. fallax* (lane 4), no DNA template (negative control, lane 5), or DNA extracted from each half of a sliced single *M. chitwoodi* juvenile (lanes 6, 7), or six *M. chitwoodi* eggs (lane 8) or a single *M. chitwoodi* egg (lane 9). Size confirmations were made by comparison with the 100 bp ladder (Gibco, BRL) in lane 1.

DNA from *M. chitwoodi* or *M. fallax* were accurately identified in all cases.

## Discussion

Difficulties in molecular diagnostics stem primarily from the lack of reproducible results between, and within, various laboratories. Subjective assessments of band intensities from agarose-separated RAPD or RFLP reaction products contributes further to the limitations of these methods for routine nematode identification purposes. Consequently, many researchers are seeking species-specific probes which yield positive/negative diagnostic results when used in combination with a non-specific primer in the PCR. Diagnostic markers have already been designed to target the *msp* gene of *Pratylenchus* spp (Setterquist *et al.*, 1996). These primers hybridize to variable sequences within an intron located between conserved, flanking coding regions. Primers designed to RAPD markers specific to *M. hapla* or *M. chitwoodi* were also constructed, but only successfully amplified 80-90% of J2s

tested (Williamson *et al.*, 1997). DNA sequencing has also assisted construction of primers which differentiate various *Meloidogyne* species using variations in the ITS. Species-specific primers for *M. hapla*, *M. javanica*, and *M. incognita* have been synthesized and demonstrated to yield amplicon polymorphisms in combination with a conserved reverse primer (Zijlstra, 1997; Powers *et al.*, 1998). However, the extensive homology observed between *M. chitwoodi* and *M. fallax* ITS sequences precluded species-specific DNA probes from being developed.

In contrast to the ITS, the high degree of variability identified within the IGS allowed us to expand the range of root-knot nematode species-specific probes to include *M. chitwoodi* and *M. fallax*. Primer sequences were chosen to have approximately 50% GC content, to lack obvious secondary structure, to lack homology with all other sequences in the Genbank database, and to provide a high annealing temperature (near that of the *Taq* extension cycle) to minimize non-specific amplification. Provided annealing temperatures are designed accordingly, additional primers can be included in future reactions since the multiplex PCR reaction conditions were the same as for the single primer pairs. Preliminary work with other species demonstrates the IGS may be generally applicable for species-specific probe development. We anticipate that such probes will facilitate rapid determination of nematode species identity.

Naturally, the universality of any species-specific primer needs to be proven. To this end, we tested the specificity of our pentaprimers upon nearly 60 different *Meloidogyne* spp. isolates from around the world. These tests were conducted in three laboratories (Canada, UK, and The Netherlands) so that the variations in extraction procedures, reaction techniques and hardware could be assessed simultaneously. The reliability of the method was clearly evidenced by accurate determination of all samples.

In addition to distinguishing *M. chitwoodi* and *M. fallax*, our pentaprimer mix provides several intrinsic features which advance upon previously published methods and which should prove useful for routine assessment of nematode identity: *i*) the species-specific markers included are complementary to the high copy-number rDNA IGS regions; the repetitive nature of these sequences increases the sensitivity of the PCR diagnosis, making possible the identification of a single juvenile or several eggs; *ii*) identification is based upon the presence or absence of species-specific bands so there is no cause for discrepancy arising from laboratory to laboratory variation in band intensity; *iii*) our pentaprimer mixture includes an internal control primer pair to verify the integrity of the detection system. The PCR can be extremely sensitive to poor DNA template quality or the presence of

inhibitors, a major problem when directly amplifying DNA of nematodes extracted from soil. Since the ITS primer pair generates an amplified fragment in all positive reactions, its visualization ensures that the other species-specific data (presence or absence) are valid; whereas if the ITS control band fails to appear then the absence of a particular nematode species DNA will not be assumed without reanalysis.

Having demonstrated the potential of the pentaprimer mixture, we look forward to incorporating additional species-specific probes into a multiplex PCR reaction mix to increase the range of species detected. Although no upper limit to the number of primers used in a multiplex reaction has been reported, limitations may arise due to competition between amplicons, limitations of substrates, and formation of chimaeras when too many primers are present. Reverse dot blot hybridizations (Iida *et al.*, 1993) may circumvent such problems. By immobilizing several species-specific probes in a filter array, amplified labelled-DNA from complex genomic samples such as those obtained from soil extracts or root washes can be hybridized to the filter to assess the species-biodiversity. This technique has been especially useful in detecting bacterial and viral pathogens on food destined for human consumption (Ehrmann *et al.*, 1994; Cromeans *et al.*, 1997), and has recently been modified for high through-put procedures using a microplate format (Soumet *et al.*, 1997). Application of this technique for detection or identification of plant-parasitic nematodes on imported agricultural commodities, or in field soils holds great potential for reducing the spread of quarantined organisms and the losses associated with nematode infestations.

#### Acknowledgements

We are grateful to G. Karssen, M. Moens, G. Janssen, L. P. G. Molendijk, D. Mugniéry, A. Schoemaker, A. van Eggermond, and G. Santo for providing samples or isolates used in this study, and to Robyn DeYoung for assistance with testing pentaprimer specificity.

#### References

- BLOK, V. C., PHILLIPS, M. S. & FARGETTE, M. (1997). Comparison of sequence differences in the intergenic region of the ribosomal cistron of *Meloidogyne mayaguensis* and the major tropical root-knot nematodes. *J. Nematol.*, 29: 16-22.
- BROWN, C. R., YANG, C. P., MOJTAHEDI, H., SANTO, G. S., & MASUELLI, R. (1996). RFLP analysis of resistance to Columbia root-knot nematode derived from *Solanum bulbocastanum* in a BC2 population. *Theoret. appl. Genet.*, 92: 572-576.
- CROMEANS, T. L., NAINAN, O. V. & MARGOLIS, H. S. (1997). Detection of hepatitis A virus RNA in oyster meat. *Appl. envir. Microb.*, 63: 2460-2463.
- CURRAN, J., MCCLURE, M. A., & WEBSTER, J. M. (1986). Genotypic differentiation of *Meloidogyne* populations by detection of restriction fragment length difference in total DNA. *J. Nematol.*, 18: 83-86.
- EHRMANN, M., LUDWIG, W. & SCHLEIFER, K. H. (1994). Reverse dot blot hybridization: A useful method for the direct identification of lactic acid bacteria in fermented food. *FEMS Microb. Letters*, 117: 143-149.
- FERRIS, V. R., FERRIS, J. M. & FAGHIHI, J. (1993). Variation in spacer region ribosomal DNA in cyst-forming species of plant-parasitic nematodes. *J. Nematol.*, 16: 177-184.
- GOLDEN, A. M., O'BANNON, J. H., SANTO, G. S. & FINLEY, A. M. (1980). Description and SEM observations of *Meloidogyne chitwoodi* n.sp. (Meloidogynidae), a root-knot nematode on potato in the Pacific Northwest. *J. Nematol.*, 12: 319-327.
- HARRIS, T. S., SANDALL, L. J., & POWERS, T. O. (1990). Identification of single *Meloidogyne* juveniles by polymerase chain reaction amplification of mitochondrial DNA. *J. Nematol.*, 22: 518-524.
- IIDA, K., ABE, A., NATSUI, H., DANBARA, H., WAKAYAMA, S. & KAWAHARA, K. (1993). Rapid and sensitive method for detection of *Salmonella* strains using a combination of polymerase chain reaction and reverse dot-blot hybridization. *FEMS Microb. Letters*, 114: 167-172.
- JANSSEN, G. J. W., JANSSEN, R., VANNOREL, A., VERKERK-BAKKER, B. & HOOGENDOORN, J. (1996). Expression of resistance to the root-knot nematodes, *Meloidogyne hapla* and *M. fallax*, in wild *Solanum* spp. under field conditions. *Europ. J. Plant Pathol.*, 102: 859-865.
- KARSSSEN, G. (1996). Description of *Meloidogyne fallax* n.sp. (Nematoda: Heteroderidae), a root-knot nematode from The Netherlands. *Fundam. appl. Nematol.*, 19: 593-599.
- NYCZEPIR, A. P., INSERRA, R. N., O'BANNON, J. H., & SANTO, G.S. (1982). Incidence and distinguishing characteristics of *Meloidogyne chitwoodi* and *M. hapla* in potato from the northwestern United States. *J. Nematol.*, 14: 347-353.
- PETERSEN, D. J. & VRAIN, T. C. (1996). Rapid identification of *Meloidogyne chitwoodi*, *M. hapla*, and *M. fallax* using primers to amplify their ribosomal intergenic spacer. *Fundam. appl. Nematol.*, 19: 601-605.
- POWERS, T. O., PLATZER, E. G. & HYMAN, B. C. (1986). Species-specific restriction site polymorphisms in root-knot nematode mitochondrial DNA. *J. Nematol.*, 18: 288-293.
- POWERS, T. O. & HARRIS, T. S. (1993). A polymerase reaction method for identification of five major *Meloidogyne* species. *J. Nematol.*, 25: 1-6.
- POWERS, T. O., TODD, T. C., BURNELL, A. M., MURRAY, P. C. B., FLEMING, C. C., SZALANSKI, A. L., ADAMS, B. A. & HARRIS, T. S. (1998). The rDNA internal transcribed spacer region as a taxonomic marker for nematodes. *J. Nematol.*, 30 (in press)
- SETTERQUIST, R. A., SMITH, G. K., JONES, R., & GOX, G. E. (1996). Diagnostic probes targeting the major sperm protein gene that may be useful in the molecular identification of nematodes. *J. Nematol.*, 28: 414-421.

- SOUMET, C., ERMEL, G., SALVAT, G. & COLIN, P. (1997). Detection of *Salmonella* spp. in food products by polymerase chain reaction and hybridization assay in microplate format. *Letters appl. Microbiol.*, 24: 113-116.
- VRAIN, T. C., WALKARCHUK, D. A., LÉVESQUE, A. & HAMILTON, R.I. (1992). Intraspecific rDNA restriction length polymorphism in the *Xiphinema americanum* group. *Fundam. appl. Nematol.*, 15: 563-573.
- WILLIAMSON, V. M., CASWELL-CHEN, E. P., WESTERDAHL, B. B., WU, F. -F., & CARYL, G. (1997). A PCR assay to identify and distinguish single juveniles of *Meloidogyne hapla* and *M. chitwoodi*. *J. Nematol.*, 29: 9-15.
- ZIJLSTRA, C. (1997). A fast PCR assay to identify *Meloidogyne hapla*, *M. chitwoodi* and *M. fallax*, and to sensitively differentiate them from each other and from *M. incognita* in mixtures. *Fundam. appl. Nematol.*, 20: 505-511.
- ZIJLSTRA, C., LEVER, A. E. M., UENK, B. J., & VAN SILFHOUT, C. H. (1995). Differences between ITS regions of isolates of the root-knot nematodes *Meloidogyne hapla* and *M. chitwoodi*. *Phytopathology*, 85: 1231-1237.