

## Acknowledgments

The author thanks Dr. Pinochet for suggesting this study and Dr. Inserra for useful procedure suggestions. Thanks are also given to Mrs. Casanovas for typing this manuscript.

## References

- BAINES, R. C., CAMERON, J. W. & SOOST, R. K. (1974). Four biotypes of *Tylenchulus semipenetrans* in California identified, and their importance in the development of resistant citrus root-stocks. *J. Nematol.*, 6 : 63-66.
- BELLO, A., NAVAS, A., BELART, C. & ALVIRA, P. (1986). *Los nematodos de los cítricos. Monografía*. Excelentísimo Ayuntamiento de Castellón de la Plana, 222 p.
- GOTTLIEB, Y., COHN, E. & SPIEGEL-ROY, P. (1986). Biotypes of the citrus nematode (*Tylenchulus semipenetrans* Cobb) in Israel. *Phytoparasitica*, 14 : 193-198.
- INSERRA, R. N., VOVLAS, N. & O'BANNON, J. H. (1980). A classification of *Tylenchulus semipenetrans* biotypes. *J. Nematol.*, 12 : 283-287.
- LAMBERTI, F., VOVLAS, N. & TIRRO, A. (1976). An Italian biotype of the citrus nematode. *Nematol. mediterr.*, 4 : 117-120.
- MARTINEZ BERINGOLA, M. L., CARCELES, A. & GUTIERREZ, M. P. (1987). Ensayos de nematocidas contra el nematodo de los agrios, *Tylenchulus semipenetrans*. *Boletín Sanid. Veget. Plagas*, 13 : 261-271.
- MCSORLEY, R., PARRADO, J. L. & DANKERS, W. H. (1984). A quantitative comparison of some methods for the extraction of nematodes from roots. *Nematropica*, 14 : 72-84.
- ORTUÑO MARTINEZ, A., GOMEZ GOMEZ, J. & CANOVAS CANDEL, F. (1969). Poblaciones nematológicas fitoparasitas en los suelos de la huerta de Murcia. *An. Edafol. Agrobiol.*, 28 : 389-398.
- TARJAN, A. C. (1972). Observations on extracting citrus nematodes, *Tylenchulus semipenetrans*, from citrus roots. *Pl. Dis. Repr.*, 56 : 186-188.

## A NEMATODE FEEDING MITE, *TYROPHAGUS PUTRESCENTIAE* (SARCOPTIFORMIS : ACARIDAE)

Anwar L. BILGRAMI \* and Qudsia TAHSEEN \*

\* Section of Nematology, Department of Zoology, Aligarh Muslim University, Aligarh-202002, India.

Accepted for publication 14 October 1991.

**Key-words :** *Tyrophagus*, Acaridae, nematodes.

There are few reports confirming nematodes as the diet of soil mites. Linford and Oliveira (1938) observed mites feeding on root knot nematodes. Murphy and Doncaster (1957) reported injury of *Heterodera* cysts by a mite. The most definite association came from the work of Rodriguez *et al.* (1962) who observed that when given equal choice, the adult mite *Macrocheles muscaedomesticae* preferred house fly eggs over nematodes while proto- and deutonymph under same conditions preferred nematodes. An oribatid mite *Pergalumna* sp., fed upon *Pelodera lambdiense* and *Tylenchorhynchus martini* in large numbers (Rockett & Woodring, 1965). Muraoka and Ishibishi (1976) described feeding by many species of mites which were identified as nematode predators. Recently, Imbriani and Mankau (1983) observed feeding of a neostigmatid mite, *Lasioseius scapulatus* on *Aphelenchus avenae* and *Cephalobus* sp.

In the present work observations were made on the predatory behaviour of *Tyrophagus putrescentiae* (Sarcoptiformis : Acaridae) using nematodes as prey.

## Materials and methods

Prey catching and feeding mechanisms were studied in culture dishes and special observation chambers. A

plastic ring (1 cm high; 2 cm diam.) glued to a coverslip at one end, was fixed in the middle of a metallic slide. The chamber, thus formed, was filled with 1 % water agar. The mites and prey nematodes were then inoculated and the ring was sealed with another coverslip to prevent air drying and escape of mites. Predation was observed on *Rhabditis* sp., *Cephalobus* sp., *Hirschmanniella oryzae* and *Tylenchorhynchus mashhoodi*. The rate of predation by *T. putrescentiae* was determined by using five adult mites against 50 individuals of prey. The effect of prey density on the rate of predation by *T. putrescentiae* was observed by placing 25, 50, 75, 100, 125, 150, 175 and 200 individuals of *H. oryzae* separately with five predators. The number of individuals killed or consumed by the mites was recorded after 24 h. All experiments were carried out at  $28 \pm 2$  °C and replicated five times.

## Predatory behaviour

*T. putrescentiae* feed on nematodes and other microorganisms in culture dishes. During routine observations the cultures of saprophagous species of nematodes viz., *Acrobeloides*, *Cephalobus*, *Rhabditis*, *Panagrellus* and predaceous nematodes viz., *Mononchus*

*aquaticus*, *Mononchoides fortidens*, *M. longicaudatus*, *Dorylaimus stagnalis*, *Aquatides thornei* were all found to be contaminated and consumed by *T. putrescentiae*. It restricted multiplication and reproduction of nematodes in culture dishes and as a result of predation brought down their population at a low level.

*T. putrescentiae* could hold its prey using its palp (used to identify prey organisms and other objects) and legs as soon as the prey came in contact. The legs are generally used for holding and wounding the prey. Chelicerae are the main killing and feeding organ, also used for grasping and crushing the prey. The ingestion of prey contents was intermittent (two or three feeding bouts) with short periods of resting activity at regular intervals. Mites also consumed its own faecal matter while moving randomly generally in absence of prey nematodes. No cannibalistic tendency was observed in *T. putrescentiae*. The nymphal stages of *T. putrescentiae* fed mostly on dead nematodes, their remains or upon the agar containing bacterial and fungal growth. *T. putrescentiae* killed *Rhabditis* sp. and *H. oryzae* most while *Cephalobus* sp. the least (Table 1). Maximum predation occurred in the population of 200 prey individuals which was double the number of prey killed when 25 individuals of *H. oryzae* were used as prey (Table 2).

## Conclusions

Observations on *T. putrescentiae* suggest its predaceous nature. These predatory mites appeared to possess greater predatory potentials mainly because of their ability to kill variety of nematodes in large numbers. *T. putrescentiae* fed most upon *Rhabditis* sp. and *H. oryzae*. Besides, mites also consumed bacterial and fungal populations grown over the agar. Increased predation by *T. putrescentiae* with increasing number of *H. oryzae* suggests density dependant predation as is the case with many predaceous nematodes (Bilgrami *et al.*, 1984; 1985; Bilgrami & Jairajpuri, 1990). The coprophagous behaviour of *T. putrescentiae* may be an added advantage as it could avoid intraspecific interactions (cannibalism).

## References

- BILGRAMI, A. L. & JAIRAJPURI, M. S. (1990). Predation by *Mononchoides fortidens* and *M. longicaudatus* (Nematoda : Diplogasterida). *Nematologica*, 35 : 475-488.
- BILGRAMI, A. L., AHMAD, I. & JAIRAJPURI, M. S. (1984). Observations on the predatory behaviour of *Mononchus aquaticus*. *Nematol. medit.*, 12 : 41-45.
- BILGRAMI, A. L., AHMAD, I. & JAIRAJPURI, M. S. (1985). Predatory behaviour of *Aquatides thornei* Nygolaimina : Nematoda). *Nematologica*, 30 (1984) : 457-462.
- IMBRIANI, J. I. & MANKAU, R. (1983). Studies on *Lasioseius*

**Table 1.** Rate of predation by *Tyrophagus putrescentiae* upon different prey nematodes.

Prey species	No. of prey killed
<i>Rhabditis</i> sp.	29-33 (34 ± 1.6)
<i>Cephalobus</i> sp.	21-27 (24 ± 2.4)
<i>Hirschmanniella oryzae</i>	28-34 (30 ± 2.6)
<i>Tylenchorhynchus mashhoodi</i>	24-31 (27 ± 2.5)

**Table 2.** Effect of prey density (*Hirschmanniella oryzae*) on the rate of predation by *Tyrophagus putrescentiae*.

Prey number	No. of prey killed	%
25	6-14 (10 ± 3.0)	40
50	23-29 (26 ± 2.0)	52
75	39-43 (41 ± 1.9)	55
100	58-62 (61 ± 3.0)	61
125	82-88 (85 ± 2.8)	68
150	108-113 (109 ± 2.3)	73
175	131-136 (133 ± 2.6)	76
200	158-164 (161 ± 2.3)	81

All figures are nearest to whole numbers.

*scapulatus*, a neostigmatid mite predaceous on nematodes. *J. Nematol.*, 15 : 523-528.

LINFORD, M. B. & OLIVIERA, J. M. (1938). Potential agent of biological control of plant parasitic nematodes. *Phytopathology*, 28 : 14.

MURAOKA, M. & ISHIBISHI, N. (1976). Nematode feeding mites and their feeding behaviour. *Appl. Ent. Zool.*, 11 : 1-7.

MURPHY, P. W. & DONCASTER, C. C. (1957). A culture method for soil microfauna and its application to the study of nematode predation. *Nematologica*, 2 : 202-214.

ROCKETT, C. L. & WOODRING, J. P. (1966). Biological investigation on a new species of ceratozetes and of *Pergalumna*. *Acarologia*, 8 : 14-18.

RODRIGUEZ, T. G., WADE, C. F. & WELLS, C. N. (1962). Nematodes as natural food for *Macrocheles muscaedomesticae* (Acarina : Macrochelidae) a predator of the house fly egg. *Ann. Ent. Soc. Am.*, 55 : 507-511.