

Scanning electronmicrographs of the anterior region of nematodes in the genus *Steinernema* (Steinernematidae : Rhabditida)

Solveig HAUKELAND

University of Reading, Department of Agriculture, Earley Gate, Reading RG6 2AT, Berks, UK.

Accepted for publication 10 April 1992.

Summary – Two isolates of *Steinernema feltiae* (= *bibionis*) and *S. carpocapsae* (UK) were examined for the number and arrangement of head sensilla on both adult and infective stages. On *S. carpocapsae* adult females, four cephalic and six labial papillae were consistently observed, conforming to previous reports. On the *S. feltiae* females however, six cephalic papillae were often observed indicating a variability in the number of these papillae. The adult males and infective stage juveniles of both species were similar with respect to the number of labial and cephalic papillae. There was also a clear difference in the morphology of the amphids between the adult males and the infective stages.

Résumé – *Observation en microscopie électronique à balayage de la région antérieure de nématodes du genre Steinernema (Rhabditida : Steinernematidae)*. – Sur deux isolats de *Steinernema feltiae* (= *bibionis*) et de *S. carpocapsae* (UK) ont été examinés le nombre et la disposition des sensilles céphaliques chez les adultes et les stades infestants. Quatre papilles céphaliques et six papilles labiales sont régulièrement observées, ce qui confirme les données antérieures. Chez *S. feltiae* par contre, six papilles céphaliques sont souvent présentes ce qui indique une certaine variabilité dans le nombre de ces papilles. Les adultes mâles et les juvéniles infestants des deux espèces sont semblables en ce qui concerne le nombre des papilles labiales et céphaliques. Il existe par contre entre adultes et stades infestants une nette différence dans la morphologie des amphides.

Key-words : Nematodes, *Steinernema*, scanning, electronmicrographs, morphology.

Nematodes in the family Steinernematidae are becoming increasingly well recognized as potential biological control agents against certain insect pests. The biology and lifecycle of many species has been studied in some detail by several workers (Bovien 1937; Poinar 1967 and references therein; Poinar 1975; Wouts 1980). Their classification however is less clear, and not made easier by an increasing number of isolates in the genus being found worldwide. The most recent nomenclature has been proposed by Poinar (1990), showing a total of nine species in *Steinernema*, and Doucet and Doucet (1990) give a key to ten nominal species in the genus.

The first steinernematid was described by Steiner (1923) as *Aplectana kraussei*, which was later placed in the genus *Steinernema* by Travassos (1927). In 1929 Glaser and Fox (1930) discovered a nematode parasitizing the Japanese beetle (*Popillia japonica*) and upon examination Steiner (1929) erected the genus *Neoaplectana* naming the new species *N. glaseri*. Although Steiner did not give clear differences between the two genera, both were accepted as valid and placed in the subfamily Steinernematinae by Filipjev (1934) later raised to family rank by Chitwood and Chitwood (1937).

The two genera remained separate until the early 1980's, when several workers suggested that they might be congeneric (Mráček *et al.*, 1981; Wouts *et al.*, 1982). They noted that in the original diagnosis one of the characters differentiating the genera was in the arrange-

ment and number of head sensilla. In the original description by Steiner, *Neoaplectana* shows six lips with one inner circle of six labial papillae, and an outer circle of six cephalic papillae, whereas *Steinernema* (*S. kraussei*) shows four cephalic papillae. In a review of nematode head sensilla by Coomans (1979) a typical arrangement of head sensilla indicates two circles of six labial papillae and a circle of four cephalic papillae. According to Coomans (1979) one circle of six labial papillae may be lost in some species. Wouts *et al.* (1982) examined the original type species *S. kraussei* (= *Aplectana kraussei*) as well as a re-isolated *Steinernema* species from the type host of *S. kraussei*. They compared these nematodes to species of the genus *Neoplectana*, using scanning electronmicrographs.

Their results showed that no differences could be found with regard to the number and arrangement of head sensilla, all those examined containing six labial papillae and four cephalic papillae, hence *Neoaplectana* was synonymized under *Steinernema*. This has been generally accepted, until an isolate that fits Steiner's original description of *S. kraussei* is found (Poinar, 1990).

In the present study scanning electronmicrograph photographs were taken of nematodes from two isolates of *S. feltiae* (= *bibionis*), and a UK strain of *S. carpocapsae* to examine the arrangement and number of head sensilla.

Materials and methods

Isolates R1.5 and V25A of *S. feltiae* (= *bibionis*) were isolated from soil samples collected in Reading (UK) and southern Norway respectively, using the *Galleria* trapping technique (Bedding & Akhurst, 1975). *S. carpocapsae* UK, was obtained from a laboratory culture at Reading University (originally isolated from sawfly larvae (Georgis & Hague, 1981)). The nematodes were cultured *in vivo* on late instar *Galleria* larvae.

For scanning electron microscopy first generation adult nematodes were dissected from previously infected *Galleria* larvae in 1/4 strength Ringer's buffer. They were rinsed several times in Ringer's before being fixed in 8 % gluteraldehyde (gluteraldehyde 25 % EM Grade, diluted in Ringer), for 2 h at room temperature. Five day

old infective stage juveniles (J3) were rinsed in 0.05 % NaCl three times at 30 min intervals and fixed in 8 % gluteraldehyde made up with distilled water. The fixed nematodes were then washed three times in Ringer (or distilled water J3) and post-fixed in 1 % osmium tetroxide for 1 h. They were rinsed again, dehydrated at 15 min intervals through 30 %, 50 % and 70 % acetone and placed at 4 °C overnight in 70 % acetone.

Subsequent dehydration continued through 90 %, 95 %, 100 % and absolute dry acetone also at 15 min intervals. The nematodes were then critically point dried in a "Samdri-780" critical point drier, mounted on stubs and finally sputter coated with gold. Twenty or more specimens were prepared for each isolate and stage examined. Photographs were taken of the head




Fig. 1. Anterior region of adult female *Steinernema*. A : *S. carpocapsae* (UK), showing cephalic papillae (arrows), and six labial papillae; B : *S. feltiae* showing six labial and the usual four cephalic papillae as well as the additional cephalic papillae (arrows); C, D : *S. feltiae* having less developed additional cephalic papillae (arrows).

region using a Hitachi S-570 scanning electron microscope at 15 or 25Kv.

Results

First generation *S. carpocapsae* females revealed a head sensilla pattern of six labial and four cephalic papillae on all specimens observed (Fig. 1 A). However, *S. feltiae* females appeared to be less consistent with respect to the number of cephalic papillae. In some specimens six cephalic papillae were observed, which were in most cases morphologically similar, i.e. "button like". The two sublateral cephalic papillae were located close to the lateral labial papillae with the remaining four cephalic papillae in the usual laterodorsal or lateroventral position (Fig. 1 B). This phenomenon was not always easily observed, in some cases these additional cephalic

papillae appeared morphologically similar to the usual four, whereas in others they appeared more like a bump in the cuticle (Fig. 1 C, D) but were always found in the same position, close to or at the base of the lateral labial papillae. No definite pore or slit-like amphids were observed on any of the females.

The males of both species clearly had six labial and four cephalic papillae, and amphids could be observed on good specimens (Fig. 2 A). These were in a similar position to the additional cephalic papillae found on the *S. feltiae* females.

Observations of the infective stage juveniles showed four distinct cephalic papillae, morphologically similar to the adult stage labial papillae, and the amphids were consistently very clear and larger, more pore-like than those found on the males. Labial papillae were not ob-




Fig. 2. Anterior region of male and the infective stage of *Steinernema*. A : Amphid (arrow) on a male *S. carpocapsae*; B : Pore-like amphid (arrow) on infective stage juvenile of *S. feltiae* (note also the four cephalic papillae; C, D : The beginning of the lateral field of *S. feltiae*, as well as the excretory pore (e).

served (Fig. 2 B). Eight lines in the lateral field was observed on some infective stage juveniles of *S. feltiae* specimens (Fig. 2 C, D). No lateral field was found on the adult stages.

Discussion

Observations on the head sensilla of *S. carpocapsae* and *S. feltiae* in this study showed two circles of papillae, an inner circle of six labial papillae and an outer circle of four, or apparently six, cephalic papillae for *S. feltiae*. The papillae in the two circles differed morphologically, the labial papillae protruding more from the base, whereas the cephalic papillae appeared flatter and "button-like". The labial papillae were always more obvious than the cephalic papillae.

The presence of two additional cephalic papillae on *S. feltiae* has not been reported previously. In some cases they were not distinctly papillae-like, but were always located in the same position, suggesting that the number of cephalic papillae may be variable in the genus *Steinernema*. Doucet (1986) reported no cephalic papillae in the original description of *Steinernema rara* from Argentina, but a re-examination of this species by Poinar *et al.* (1986) showed the presence of the usual four cephalic papillae, although they were not particularly distinct. The presence of cephalic papillae therefore may either be difficult to observe and hence missed, or some may be indistinct and not counted. In the present study there appeared to be some variability in the number of cephalic papillae on the *S. feltiae* adult females, hence the presence or number of cephalic papillae will not only be due to the observers interpretation, but also to the variability of the cephalic papillae on the nematodes themselves.

Amphids were not observed on the female specimens, although on *S. feltiae* the additional cephalic papillae were located in the usual amphid position indicating that these cephalic papillae have not completely developed to amphids (amphids being derived from cephalic papillae). Figure 1 B clearly shows that one of the sublateral cephalic papillae is very similar to the usual four, whereas the other is less developed, perhaps indicating a transitional phase in which these cephalic papillae have not developed into amphids. It may be postulated that the females, in relation to their micro habitat in an insect cadaver, being surrounded by their nutritional requirements have no particular need for amphids, which are chemosensory (Coomans, 1979). In most descriptions of *Steinernema* species (adult forms) the presence of amphids are shown and located at the base of the lateral labial papillae (eg. Doucet & Doucet, 1990; Poinar, 1988), however their function may be questionable, since their development can be partial or apparently absent, based on the SEM observations of *S. feltiae* and *S. carpocapsae* in this study. Males on the other hand, on which small slit-like amphids were found, may have amphids to assist in finding females in order to mate.

The adult males from both species showed no variability in the number of head sensilla, both the inner circle of six labial papillae and the outer circle of four cephalic papillae being distinct and easily observed. Small slit-like amphids were quite distinct in most cases and located in a similar position to the extra cephalic papillae seen on some of the *S. feltiae* females.

In contrast to the adult stages, the infective stage juveniles showed only one circle of four cephalic papillae, the labial papillae not being visible. Most striking were the relatively large pore-like amphids (Fig. 2 B). Infective juveniles are present in soil naturally, and in order to infect a suitable host they must first be able to sense its presence, it has been shown that infective stage juveniles are attracted to insect excretory products and carbon dioxide (Schmidt & All, 1979; Gaugler *et al.*, 1980; Pye & Burman, 1981) which probably accounts for the clear presence of amphids.

The lateral field for the *S. feltiae* infective stage juveniles examined had eight lines, which is in accordance with Mráček and Bednarek (1991), for this species.

Acknowledgments

I would like to thank Dr. D. J. Hunt from IIP (International Institute of Parasitology) for critical reading of the manuscript. The assistance of Dr. Lynda Bonner (University of Reading, School of Plant Sciences) and Sue Hines IIP for advice on specimen preparation was also greatly appreciated. Finally I must thank Rena for initialising interest in this work.

This work was funded by The Norwegian Agricultural Research Council (NLVF, Oslo).

References

- BEDDING, R. A. & AKHURST, R. J. (1975). A simple technique for the detection of insect parasitic rhabditid nematodes in soil. *Nematologica*, 21 : 109-110.
- BOVIEN, P. (1937). Some types of association between nematodes and insects. *Vidensk. Medd. dansk naturh. Foren.*, 101 : 1-114.
- COOMANS, A. (1979). The anterior sensilla of nematodes. *Revue Nématol.*, 2 : 259-283.
- DOUCET, M. M. A. (1986). A new species of *Neoplectana* Steiner, 1929 (Nematoda : Steinernematidae) from Córdoba, Argentina; *Revue Nématol.*, 9 : 317-323.
- DOUCET, M. M. A. & DOUCET, M. E. (1990). *Steinernema ritteri* n.sp. (Nematoda : Steinernematidae) with a key to the species of the genus *Nematologica*, 36 : 257-265.
- FILIPJEV, I. N. (1934). Miscellanea nematologica. I. Eine neue Art der Gattung *Neoplectana* Steiner nebst Bemerkungen über die systematische Stellung der letzten. *Parazit. Sb.*, 4 : 229-239.
- GAUGLER, R., LEBECK, L., NAKAGAKI, B. & BOUSH, G. M. (1980). Orientation of the entomogenous nematode *Neoplectana carpocapsae* to carbon dioxide. *Envir. Ent.*, 9 : 649-652.

- GEORGIS, R. & HAGUE, N. G. M. (1981). A neoaplectanid nematode in the web-spinning larch sawfly *Cephalcia lariciphila*. *Ann. appl. Biol.*, 99 : 171-177.
- GLASER, R. W. & FOX, H. (1930). A nematode parasite of the Japanese beetle (*Popillia japonica* Newm.). *Science*, 71 : 16-17.
- MRÁČEK, Z., WEISER, J. & GERDIN, S. (1981). Head and cuticular structures of some species in the family Steinernematidae (Nematoda). *Nematologica*, 27 : 443-448.
- POINAR, G. O. Jr. (1967). Description and taxonomic position of the DD-136 nematode (Steinernematidae, Rhabditoidea) and its relationship to *Neoaplectana carpocapsae* Weiser. *Proc. helminth. Soc. Wash.*, 34 : 119-209.
- POINAR, G. O. Jr. (1975). *Entomogenous nematodes. A manual and host list of insect-nematode associations*, Leiden E.J. Brill, 317 p.
- POINAR, G. O. Jr. (1988). Redescription of *Neoaplectana affinis* Bovien (Rhabditida : Steinernematidae). *Revue Nématol.*, 11 : 143-147.
- POINAR, G. O. Jr. (1990). Taxonomy and biology of Steinernematidae and Heterorhabditidae. In: Gaugler R., & Kaya, H. K. (Eds). *Entomopathogenic nematodes in biological control*. Boca Raton USA, CRC Press, 23-61.
- POINAR, G. O. Jr., MRÁČEK, Z. & DOUCET, M. M. A. (1988). A reexamination of *Neoaplectana rara* Doucet, 1986 (Steinernematidae : Rhabditida). *Revue Nématol.*, 11 : 447-449.
- PYE, A. E. & BURMAN, M. (1981). *Neoaplectana carpocapsae* : Nematode accumulations on chemical and bacterial gradients. *Exp. Parasit.*, 51 : 13-20.
- SCHMIDT, J. & ALL, J. N. (1979). Attraction of *Neoaplectana carpocapsae* (Nematoda : Steinernematidae) to common excretory products of insects. *Envir. Ent.*, 8 : 55-61.
- STEINER, G. (1923). *Aplectana kraussei* n. sp., eine in der Blattwespe *Lyda* sp. parasitierende Nematodenform, nebst Bemerkungen über das Seitenorgan der parasitischen Nematoden. *Zentbl. Bakt. ParasitKde*, 59 : 14-18.
- STEINER, G. (1929). *Neoaplectana glaseri* n.g., n.sp. (Oxyuriidae), a new nemtic parasite of the Japanese beetle. *J. Wash. Acad. Sci.*, 19 : 436-440.
- TRAVASSOS, L. (1927). Sobre o genera *Oxysomatium*. *Bolm. biol. Sao Paolo*, 5 : 20-21.
- WOUTS, W. M. (1980). Biology, lifecycle and redescription of *Neoaplectana bibionis* Bovien, 1937 (Nematoda : Steinernematidae). *J. Nematol.*, 12 : 62-72.
- WOUTS, W. M., MRÁČEK, Z., GERDIN, S. & BEDDING, R. A. (1982). *Neoaplectana* Steiner, 1929 a junior synonym of *Steinernema* Travassos, 1927 (Nematoda, Rhabditida). *Syst. Parasit.*, 4 : 147-154.