

Field trial evaluation of nematode susceptibility within *Musa*

Nigel S. PRICE

I.R.A. Ekona Centre, P.M.B. 25, Buea, South-West Province, Cameroon.

Accepted for publication 11 October 1993.

Summary – Fifty-two clones of *Musa* were evaluated for their susceptibility to the plant-parasitic nematodes *Radopholus similis*, *Hoplolaimus pararobustus* and *Meloidogyne incognita* in two field trials. As a group the AAB Plantains showed the greatest susceptibility to both *R. similis* and *H. pararobustus*. *Musa* AAA and ABB types tended to show lower susceptibility to *R. similis* and some types exhibited near resistance. *H. pararobustus* and *M. incognita* occurred in lower numbers than *R. similis* and intervarietal differences in susceptibility were less marked.

Résumé – Évaluation par essais en champ de la sensibilité aux nématodes chez le genre *Musa* – Au cours de deux essais en champ, cinquante-deux espèces et variétés de *Musa* ont été testées pour leur sensibilité envers les nématodes *Radopholus similis*, *Hoplolaimus pararobustus* et *Meloidogyne incognita*. Le groupe des bananiers plantains AAB présente la plus grande sensibilité tant envers *R. similis* que *H. pararobustus*. Les types AAA et ABB marquent une tendance vers une moindre sensibilité à *R. similis*, quelques variétés étant même proches de la résistance. *H. pararobustus* et *M. incognita* sont moins représentés que *R. similis* et dans leur cas les différences variétales de sensibilité sont moins marquées.

Key-words : *Musa*, plantains, bananas, nematodes, *Radopholus similis*, resistance, susceptibility.

Of the plant-parasitic nematodes known to cause damage to bananas and plantains, *Radopholus similis*, is considered the most important (Gowen & Quénéhervé, 1990) in particular due to its status as the most important nematode pest of commercial plantations. In Africa this nematode also poses a major and increasing threat to smallholder agriculture (Sarah, 1989). With increased awareness of the important role bananas and plantains occupy in domestic consumption there is a need for a better understanding of the host parasite relationships between plant-parasitic nematodes (in particular *R. similis*) and *Musa* varieties other than the AAA Cavendish cultivars upon which the commercial sector relies (Anon., 1988). Such information is needed both because breeders have so far successfully transmitted resistance to *R. similis* from only source (Pinochet, 1988 a) and because of suggestions that known naturally occurring cultivars with greater disease resistance may be promoted until plant breeding is able to provide improved varieties (Champion, 1976). Following a reorganization of the *Musa* germplasm collection at the Institut de la Recherche Agronomique Njombe Station, Littoral Province, Cameroon, planting material comprising 52 clones representing various genetic groups of *Musa* became available and was used in two field trials.

Materials and methods

Two adjacent trials (A and B) identical in design were planted in April 1989. The site is in the humid lowland tropical zone of the country 80 m above sea level in a

region of leached sandy soils of volcanic origin. Soils tend to be acid (pH 5.1-6.1) and deficient in potassium (Delvaux, 1989). One month prior to planting these trials the land had been growing the AAA Cavendish banana variety Poyo and was known to be infested with *R. similis*.

Planting material (trimmed "suckers") was planted in rows 3.2 m × 1.5 m. Each trial comprised 27 different clones and both included the two AAA Cavendish bananas Poyo and Grande Naine as reference varieties. Both trials were in randomized block designs of eight replicates, each plot consisting of a single plant, each variety representing a treatment and surrounded by a single outside border of Poyo. No nematocides or insecticides were applied, aerial spraying against Black Sigatoka (*Mycosphaerella fijiensis* var. *difformis* Mulder & Stover) occurred as necessary. Fertilization followed normal procedures and a limited amount of irrigation took place during the dry season.

Trials were sampled in September and October 1990. Root sampling and nematode extraction followed established procedures (Quénéhervé & Cadet, 1986; Pinochet, 1988 b). Each plant provided one sample, 25 g of root material being washed, blended and passed through screens of 250 µm, 50 µm and 32 µm apertures. Material retained by the 250 µm screen was discarded, material collected on the other screens was made up to 100 ml with water to form the sample. In cases of very high nematode numbers the sample was diluted to 500 ml. Nematodes in three 1-ml aliquots from each

sample were counted. A complete block was processed in a single day, blocks being sampled sequentially on consecutive days.

Counts of the three most commonly found plant-parasitic nematode species are reported here as numbers of nematodes per 100 g root fresh weight (RFW). Numbers of *R. similis* consist of adult females plus immature stages (ie. excluding adult males). Numbers of *Meloidogyne incognita* consist of all endoparasitic stages plus adult males but does not include vermiform second-stage juveniles. All *Hoplolaimus pararobustus* were counted.

After the log-transformation ($\log_e(n+1)$) results were subjected to Analysis of Variance using GENSTAT (Genstat, 1987) and the Duncans' Multiple Range Test was applied.

Results

Details of *Musa* clones used and nematode numbers obtained for trials A and B are shown in Tables 1 and 2, respectively.

Greater numbers of *R. similis* occurred than of the other two nematode species. Maximum observed values were 299 330 *R. similis* per 100 g RFW for the AAB Plantain Batard in Trial A and 296 670 *R. similis* per 100 g RFW for the AAB Plantain Obel in Trial B. Highest mean values (de-transformed from the $\log_e(n+1)$ means) were 32 859 *R. similis* per 100 g RFW for the AAB variety Laknao and 13 359 *R. similis* per 100 g RFW for the AAB Plantain Esang for trials A and B respectively (log-transformed means 10.4 ± 0.7 and 9.5 ± 0.7 respectively). The least susceptible variety in Trial A was the AAA Ibota group banana Yangambi with 4.0 *R. similis* per 100 g RFW (log-transformed mean 1.6 ± 0.7), and in Trial B the AAB Pisang Kelat with 6.0 *R. similis* per 100 g RFW (log-transformed mean 1.9 ± 0.7).

The maximum observed numbers of *H. pararobustus* found in trial A was 14 000 per 100 g RFW on *M. acuminata* and in trial B 18 670 per 100 g RFW on the AAB Plantain Obel. Maximum means (de-transformed from the $\log_e(n+1)$ means) were 298.0 on the AAA Cavendish Grande Naine in Trial A and 3293.0 on the AA Pisang Trimulin in Trial B (log-transformed means, 5.7 ± 0.6 and 8.1 ± 0.7 g RFW, respectively). *H. pararobustus* numbers were generally low and few clones differed significantly in susceptibility.

Root-knot nematode, *Meloidogyne incognita*, was the least abundant nematode. Maximum observed numbers were 10 000 per 100 g RFW on the AAB Plantain French Sombre in trial A and 9330 per 100 g RFW on the ABB Bluggoe Christine in trial B. Maximum means (de-transformed from the $\log_e(n+1)$ means) were 98.0 *Meloidogyne incognita* per 100 g RFW on *Musa basjoo* in trial A and 89.0 *Meloidogyne incognita* per 100 g RFW on *Musa acuminata burmannica* in Trial B (log-trans-

formed means 4.6 ± 0.6 and 4.5 ± 0.6 respectively). As with *H. pararobustus* inter-clonal differences were not marked. No *Meloidogyne incognita* were recorded on the AAB variety Laknao although vermiform J2s were observed in some samples.

Discussion

Total resistance (immunity) was not confirmed for any of the three nematode species considered and, as in other work, only differences between clones in susceptibility were observed (Gowen, 1976; Pinochet, 1988 a). The predominant nematode species present was *R. similis*, as found in Cavendish banana plantations elsewhere in West Africa (Quénéhervé *et al.*, 1991). *Helicotylenchus multicinctus* which can outnumber *R. similis* in some areas (Gowen, 1979) was found infrequently and was not evaluated.

These results show a high level of susceptibility to *R. similis* among the AAB Plantains. In both trials, six of the ten most susceptible clones were AAB Plantains (i.e. twelve out of the total of seventeen AAB Plantain varieties evaluated) and the closely related AAB variety Laknao was the most susceptible clone in Trial A. This is believed to be the first extensive evaluation of varietal susceptibility conducted in Africa and is clear evidence of the threat this nematode poses to this important crop (Sarah, 1989).

By contrast the AAA Cavendish varieties, mainstay of the commercial (banana export) industry, on which much money and effort is spent in chemical control (Gowen & Quénéhervé, 1990) were more resistant than the AAB Plantains. These results agree with those of Perez *et al.* (1986) in Cuba, reporting greater susceptibility, to both *R. similis* and *Pratylenchus coffeae*, of AAB Plantains compared with AAA Bananas. Maximum *R. similis* numbers observed on the reference variety Grande Naine, 86 000 and 56 000 *R. similis* per 100 g root for Trials A and B respectively, were approximately a quarter of the maxima recorded on AAB Plantains. Means numbers of *R. similis* found in the reference varieties (Tables 1 and 2), correspond well with field populations reported on Cavendish bananas elsewhere (Sarah & Vilardebó, 1979; Davide, 1980; Loridat, 1989).

Bananas of the Gros Michel group have long been considered less susceptible to *R. similis* than Cavendish varieties (Simmonds, 1966). Using a root-lesion index Wehunt *et al.* (1978) found roots of the Gros Michel var. Cocos to be significantly less damaged than roots of the Cavendish varieties they evaluated. Davide and Marasigan (1985) also reported lower numbers of *R. similis* in roots of Gros Michel than Cavendish and Trial A (Table 1) showed Gros Michel to be significantly less susceptible than two of the three Cavendish varieties included. Mateille (1992), in a study using tissue cultured plants, attributes this lower susceptibility of Gros Michel (compared to the Cavendish variety Poyo) to

Table 1. Numbers¹ of *Radopholus similis*², *Hoplolaimus pararobustus* and *Meloidogyne incognita*³ per 100 g Fresh Weight Root on 27 field grown Musa clones. Njombe Field Trial A.

Clone	Genome	Genetic Group	<i>R. similis</i>		<i>H. pararobustus</i>		<i>M. incognita</i>	
			Numbers	DMR	Numbers	DMR	Numbers	DMR
Laknao	AAB	Laknao	32 858.6	a	4.5	e	0.0	e
Ebanga	AAB	Plantain	24 342.0	ab	39.4	abcdef	1.5	cde
French Clair	AAB	Plantain	24 342.0	ab	23.5	cdefg	5.0	bcde
Batard	AAB	Plantain	13 358.0	abc	11.2	cdefg	32.1	ab
Mujuba	AAA	Lujugira	10 937.0	abcd	2.0	gh	5.0	bcde
Corne N° 5	AAB	Plantain	10 937.0	abcd	243.7	ab	43.7	ab
Popoulou	AAB	Popoulou	8 102.1	abcd	35.6	abcdef	13.9	abcd
French Sombre	AAB	Plantain	6 634.2	abcd	35.6	abcdef	2.0	cde
Corn Type	AAB	Plantain	6 001.9	abcd	11.2	cdefgh	1.2	de
Grande Naine	AAA	Cavendish	4 446.1	abcd	297.9	a	11.2	abcd
<i>Musa basjoo</i>			3 639.9	abcde	19.1	cdefgh	98.5	a
<i>Musa acuminata</i> Type II			2 996.3	bcde	17.2	cdefgh	3.9	bcde
Bluggoe	ABB	Bluggoe	2 439.6	cdef	3.9	fgh	1.2	de
Figue Pomme Adju	AAA	Lujugira	1 635.0	cdef	4.5	efgh	1.0	de
Madre del Platano	AAB	Plantain	1 479.3	cdef	243.7	ab	10.0	bcd
Pisang Mas	AA		1 211.0	defg	5.0	efgh	17.2	abcd
Poyo	AAA	Cavendish	1 095.6	defg	10.0	defgh	4.5	bcde
Americani	AAA	Cavendish	445.9	efgh	39.4	abcdef	5.0	abc
<i>Musa textilis</i> ⁴			269.4	fgh	48.4	fgh	5.0	bcde
Pelipita	ABB	Pelipita	163.0	ghi	3.9	fgh	11.2	abcd
<i>Musa coccinea</i>			108.9	hi	29.0	bcdefg	0.0	e
<i>Musa balbisiana</i> (CMR)			98.5	hi	1.5	h	4.5	bcde
Gros Michel	AAA	Gros Michel	80.4	hi	65.7	abcd	1.2	de
Rajapuri India	AAB	Nendra Padaththi	53.6	hi	108.9	abc	5.0	bcde
1877	AAAA		48.4		48.4	efgh	12.5	abcd
<i>Musa laterita</i>			23.5	ij	6.4	efgh	6.4	bcde
Yangambi	AAA	Ibota	3.9	j	12.5	cdefgh	15.4	abcd

¹Nematode numbers per 100 g Fresh Weight Root, de-transformed from the $\log_e(n+1)$ mean of 8 replicates; ²*R. similis* adult females + juveniles; ³*M. incognita* endoparasitic stages + males. ⁴Possibly a *textilis* × *balbisiana* hybrid. For each nematode species the $\log_e(n+1)$ transformed means of Musa clones followed by the same letter do not differ significantly ($P = 0.05$) according to the Duncan's Multiple Range Test (DMR).

both reduced and delayed initial invasion and lower nematode multiplication.

H. pararobustus has been reported attacking both AAA Cavendish bananas and AAB plantains in Côte d'Ivoire in similar numbers to those reported here (Adiko, 1988; Mateille *et al.*, 1988). The AAB Plantains as a group (as with *R. similis*) again appear to be more susceptible than other types. Five of the seven AAB Plantains evaluated in Trial A and six of the ten evaluated in Trial B ranked among the ten most susceptible varieties in each trial. Two non-parametric tests failed to show statistically significant rank correlations between susceptibility to these two nematode species over all varieties evaluated. However, one may speculate that the greater susceptibility to both nematode species observed

within the AAB Plantains may have a similar basis. Mateille (1992) found Gros Michel banana to be significantly more susceptible to *H. pararobustus* than the Cavendish variety Poyo. This was also seen in Trial A (Table 1) though the difference failed to reach statistical significance.

Root-knot nematodes are widely distributed and commonly found on bananas (Gowen & Quénehervé, 1990) although their importance in causing yield loss may be more restricted, and complicated by possible interactions with other nematode species. Davide (1980) showed how *R. similis* numbers may increase at the expense of *M. incognita*. Santor and Davide (1982) described the histological incompatibility between the two species. A possible example of this is provided by the

AAB Laknao, rated root-knot susceptible by Davide and Marasigan (1985). In Trial A this variety was found free of *M. incognita* but with the highest population of *R. similis*. By contrast in Trial B the AAB Pisang Kelat was the variety most resistant to *R. similis* but had the second highest number of *M. incognita*. An antagonism of *R. similis* to *M. incognita* may explain the generally much lower numbers of *M. incognita* found in this work compared with numbers reported when using pure root-knot nematode populations (Davide & Marasigan, 1985). As found by these workers most varieties screened, including both *M. balbisiana* and *M. acuminata* showed some degree of susceptibility.

Most hopes for breeding resistance to *R. similis* are based on *Musa acuminata* varieties of the Pisang Jari Buaya (PJB) group and their progenies (Pinochet, 1988 a, 1992). Although no PJBs were included here some of the varieties tested appear, even allowing for possible differences in extraction efficiencies and popu-

lation assessments, to show levels of *R. similis* resistance similar or greater to those reported for PJB and its progenies (Pinochet & Rowe, 1978). In Trial A the AAA Ibota group banana Yangambi showed resistance to *R. similis*, with a (de-transformed) mean population of six *R. similis* per 100 g RFW, an observation supported by greenhouse work (Sarah *et al.*, 1992). This variety appears to be resistant to another major pest of *Musa* in West Africa, the Banana Borer Weevil *Cosmopolites sordidus* Germar (Fogain & Price, 1993) and is also resistant to Black Sigatoka *Mycosphaella fijiensis* var. *difformis* (Fouré, 1982). However both *M. acuminata* accessions used in this work, including the subspecies *burmannica* Type Calcutta 4 used in Trial B (the source of Black Sigatoka resistance in most breeding schemes) appear relatively susceptible to *R. similis* (Tables 1 and 2). In Trial B both the AB Safet Velchi and the AAB Pisang Kelat also appeared to support populations as low as reported on PJB and its progenies (Pinochet, 1992).

Table 2. Numbers¹ of *Radopholus similis*², *Hoplolaimus pararobustus* and *Meloidogyne incognita*³ per 100 g Fresh Weight Root on 27 field grown *Musa* clones. Njombe Field Trial B.

Clone	Genome	Genetic Group	<i>R. similis</i>		<i>H. pararobustus</i>		<i>M. incognita</i>	
			Numbers	DMR	Numbers	DMR	Numbers	DMR
Esang	AAB	Plantain	13 358.7	a	811.4	abcde	89.0	abcdef
Obel	AAB	Plantain	7 331.0	ab	402.4	abcde	4.5	bcdef
Pisang Trimulin	AA		3 640.0	abc	3 293.5	a	11.2	abcdef
Mbouroukou N° 3	AAB	Plantain	3 640.0	abc	329.3	bcdefg	3.9	cdef
Kedong Kekang	AAB	Plantain	2 696.3	abcd	811.4	abcd	13.9	abcdef
Psi-Psi	AAB	Plantain	2 696.3	abcd	1 211.0	abcd	5.0	bcdef
Grande Naine	AAA	Cavendish	1 997.2	abcd	147.4	cdefg	5.0	bcdef
Mbotoko Rouge	AAB	Plantain	1 635.0	abcd	811.4	abcd	48.4	ab
Guyod	AA		1 635.0	abcd	444.9	abcde	13.9	abcdef
Williams	AAA	Cavendish	1 479.3	abcd	329.3	bcdefg	35.6	abcd
Figue Rose	AAA	Red	1 211.0	abcde	43.7	fgh	1.2	ef
<i>M. acuminata burmannica</i> Calcutta 4			991.3	bcdef	89.0	defgh	89.0	a
Cachao	ABB	Bluggoe	664.1	cdef	15.4	h	12.5	abcdef
1 Hand Planty	AAB	Plantain	543.6	cdef	89.0	defgh	43.7	abc
Plantain N° 17	AAB	Plantain	491.7	cdefg	297.9	bcdefg	1.2	ef
Plantain N° 2	AAB	Plantain	364.0	cdefg	89.0	defgh	17.2	abcde
Manneah	AAA	Cavendish	364.0	cdefg	147.4	cdefg	3.5	def
Poyo	AAA	Cavendish	243.7	defgh	133.3	cdefgh	6.4	bcdef
Christine	ABB	Bluggoe	120.5	efghi	402.4	abcdef	7.2	abcde
Thong Dok Mak	AA		98.5	fghi	243.7	bcdefg	17.2	ef
Big Ebanga	AAB	Plantain	48.4	ghij	1 095.6	abc	1.2	ef
Lacatan	AAA	Cavendish	26.1	hij	39.4	gh	8.2	bcdef
Foconah	AAB	Pome	26.1	hij	1 807.0	ab	1.0	f
Figue Pomme Ekona	AAB	Silk	21.2	ij	13.9	h	1.5	ef
Pisang Papan	AAA	Undetermined	15.4	ij	48.4	fgh	1.5	ef
Safet Velchi	AB		12.5	ij	80.4	efgh	5.7	bcdef
Pisang Kelat	AAB	Pisang Kelat	5.7	j	133.3	cdefgh	48.4	ab

¹Nematode numbers per 100 g Fresh Weight Root, de-transformed from the $\log_e(n+1)$ mean of 8 replicates; ²*R. similis* adult females + juveniles; ³*M. incognita* endoparasitic stages + males. For each nematode species the $\log_e(n+1)$ transformed mean of *Musa* clones followed by the same letter do not differ significantly ($P = 0.05$) according to the Duncans Multiple Range Test (DMR).

Pinochet (1988 *a*) called for information on the nematode susceptibility of potential substitutes, among them the ABB variety Pelipita, for Black Sigatoka susceptible Plantain varieties. Davide and Marasignan (1985) rated Penipita (*sic*) as resistant to *R. similis*. This variety was significantly less susceptible to *R. similis* than any of the AAB Plantains evaluated in Trial A (Table 1). The (de-transformed) mean number of *R. similis* (164 per 100 g root) on Pelipita was less than 1% of the (de-transformed) means recorded on the two most susceptible AAB Plantains in this trial. Although the AAB Plantains as a group exhibited high susceptibility to *R. similis* four other AABs, namely the varieties Foconah, Figue Pomme d'Ekona, Pisang Kelat and Rajapuri India of the Pome, Silk, Pisang Kelat and Nendra Padaththi Groups respectively, showed significantly lower susceptibility. Of the AAB Plantains evaluated the False Horn variety Big Ebanga showed similar low susceptibility to *R. similis*, an impression also gained during a field survey (Bridge *et al.*, unpubl.).

In conclusion this work demonstrates that a high degree of variability in susceptibility to *R. similis* exists within *Musa*. Of particular interest is how certain varieties are able to support much higher *R. similis* populations than other, the mechanisms behind this greater susceptibility, and if such mechanisms are also associated with a degree of tolerance to this nematode. Although the search for resistance to *R. similis* has so far been disappointing (Pinochet, 1988 *a*) this work shows that mechanisms of much reduced susceptibility do exist. Reduced susceptibility could in itself be a major advance and this information may be of use, particularly with the possibilities offered by advanced breeding techniques (Stover & Buddenhagen, 1986).

Acknowledgments

The author is a Technical Co-operation Officer of the UK Overseas Development Administration (ODA). Dr. Graham McLaren, ODA Biometrician, Nkolbisson, Cameroon kindly helped with statistical analysis. Thanks to Christophe Jenny of the CRBP Njombe for helpful comments and discussion.

References

- ADIKO, A. (1988). Plant-parasitic nematodes associated with Plantain, *Musa paradisiaca* (AAB), in the Ivory Coast. *Revue Nématol.*, 11 : 109-113.
- ANON. (1988). Nematodes and the Borer Weevil in Bananas : present status of research and outlook. *Proc. INIBAP Worksh., Bujumbura, Burundi, 7-11 Dec. 1987* : 11-12.
- CHAMPION, J. (1976). Quelques problèmes de la production des bananes plantains. *Fruits*, 31 : 666-668.
- DAVIDE, R. G. (1980). Influence of cultivar, age, soil texture, and pH on *Meloidogyne incognita* and *Radopholus similis* on banana. *Pl. Dis.*, 64 : 571-573.
- DAVIDE, R. G. & MARASIGNAN, L. Q. (1985). Yield loss assessment and evaluation of resistance of banana cultivars to the nematodes *Radopholus similis* Thorne and *Meloidogyne incognita* Chitwood. *Philipp. Agriculturalist*, 68 : 335-349.
- DELVAUX, B. (1989). Rôle des constituants de sols volcaniques et de leurs propriétés de charge dans le fonctionnement de l'agrosystème bananier au Cameroun. *Fruits*, 44 : 309-319.
- FOGAIN, R. & PRICE, N. S. (1993). Varietal screening of some *Musa* cultivars for susceptibility to the banana borer weevil, *Cosmopolites sordidus* (Coleoptera : Curculionidae). *Proc. II-TA Worksh. Banana Borer Weevil, Cotonou, Benin, Nov. 1991*.
- FOURÉ, E. (1982). Les Cercosporioses du bananier et leurs traitements. Comportement des variétés. Étude de la sensibilité variétale des bananiers et plantains à *Mycosphaerella fijiensis* (Morelet) au Gabon (maladie des raies noires). I. Incubation et évolution de la maladie. *Fruits*, 37 : 749-770.
- GENSTAT (1987). *Genstat 5. Reference Manual*. Lawes Agricultural Trust, Oxford Science Publications. XVIII + 748 p.
- GOWEN, S. R. (1976). Varietal responses and prospects for breeding nematode resistant banana varieties. *Nematropica*, 6 : 45-49.
- GOWEN, S. R. (1979). Some considerations of problems associated with the nematode pests of bananas. *Nematropica*, 9 : 79-91.
- GOWEN, S. R. & QUÉNÉHERVÉ, P. (1990). Nematode parasites of bananas, plantains and abaca. In : Luc, M., Sikora, R. & Bridge, J. (Eds). *Plant parasitic nematodes in subtropical and tropical agriculture* : 431-460.
- LORIDAT, P. (1989). Étude de la microflore fongique et des nématodes associés aux nécroses de l'appareil souterrain du bananier en Martinique. Mise en évidence du pouvoir pathogène du genre *Cylindrocladium*. *Fruits*, 44 : 587-598.
- MATEILLE, T. (1992). Comparative development of three banana-parasitic nematodes on *Musa acuminata* (AAA Group) cvs Poyo and Gros Michel vitro-plants. *Nematologica*, 38 : 203-214.
- MATEILLE, T., FONCELLE, B. & FERRER, H. (1988). Lutte contre les nématodes du bananier par submersion du sol. *Revue Nématol.*, 11 : 235-238.
- PEREZ, J. A., VALDES, S. & MOLA, G. (1986). Comportamiento varietal del Platano (*Musa* sp.) al ataque de los nematodos *Radopholus similis* y *Pratylenchus coffeae*. *Cienc. Técn. Protec.*, 9 : 13-22.
- PINOCHET, J. (1992). Breeding bananas for resistance against lesion forming nematodes. In : Gommers, F. & Maas, P. W. (Eds). *Nematology from molecule to ecosystem*. Europ. Soc. Nematologists, Wageningen, the Netherlands : 157-169.

- PINOCHET, J. (1988 *a*). Comments on the difficulty in breeding bananas and plantains for resistance to nematodes. *Revue Nématol.*, 11 : 3-5.
- PINOCHET, J. (1988 *b*). A method of screening bananas and plantains to lesion forming nematodes. *Proc. IMIBAP Worksh., Bujumbura, Burundi, 7-11 déc. 1987* : 62-65.
- PINOCHET, J. & ROWE, P. R. (1978). Reaction of two banana cultivars to three different nematodes. *Pl. Dis. Repr.*, 62 : 727-729.
- QUÉNÉHERVÉ, P. & CADET, P. (1986). Une nouvelle technique d'échantillonnage pour l'étude des nématodes endoparasites du bananier. *Revue Nématol.*, 9 : 95-97.
- QUÉNÉHERVÉ, P., CADET, P., MATEILLE, T. & TOPART, P. (1991). Population of nematodes in soils under bananas, cv. Poyo, in the Ivory Coast. 5. Screening of nematicides and horticultural results. *Revue Nématol.*, 14 : 231-249.
- SANTOR, W. & DAVIDE, R. G. (1982). Interrelationship of *Radopholus similis* and *Meloidogyne incognita* in Banana. *Philipp. Phytopath.*, 18 : 22-33.
- SARAH, J. L. (1989). Banana nematodes and their control in Africa. *Nematropica*, 19 : 199-216.
- SARAH, J. L., BLAVIGNAC, F., SABATINI, C. & BOISSEAU, M. (1992). Une méthode de laboratoire pour le criblage variétal des bananiers vis-à-vis de la résistance aux nématodes. *Fruits*, 47 : 559-564.
- SARAH, J.-L. & VILARDEBÓ, A. (1979). L'utilisation du Miral en Afrique de l'ouest pour la lutte contre les nématodes du bananier. *Fruits*, 34 : 729-741.
- SIMMONDS, N. W. (1966). *Bananas*, 2nd ed. London, Longmans, 512 p.
- STOVER, R. H. & BUDDENHAGEN, I. W. (1986). Banana - breeding : polyploidy, disease resistance and productivity. *Fruits*, 41 : 175-191.
- WEHUNT, E. J., HUTCHINSON, D. J. & EDWARDS, D. I. (1978). Reaction of banana cultivars to the burrowing nematode (*Radopholus similis*). *J. Nematol.*, 10 : 368-370.