

Reproductive cycle and biometric relations in a population of *Actinopyga echinites* (Echinodermata: Holothuroidea) from the lagoon of New Caledonia, western tropical Pacific

C.CONAND

ORSTOM, Nouméa, New Caledonia

1 INTRODUCTION

Despite their importance in the benthic fauna, there is little information on tropical holothurians and their biology and ecology remain little known (Bakus, 1973). The reproductive cycles of three large species of holothurians, *Holothuria* (*Microthele*) *nobilis*, *H. (M.) fuscogilva* and *Thelenota ananas* found in New Caledonia, have been studied (Conand, 1981).

Actinopyga echinites (Jaeger, 1833), belonging to the family *Holothuriidae*, is common in New Caledonia (Intes and Menou, 1977) and widely distributed in the Indo-West Pacific (Clark and Rowe, 1971). This paper reports the reproductive cycle of *A. echinites*, and individual and population characteristics.

2 MATERIAL AND METHODS

Samples were collected from November 1978 to May 1980 in lagoon sites near Noumea (Fig. 1). Collections were made at monthly intervals by skin diving generally on the reef slope of Maitre Islet, except in November 1978 at Anse Vata site, February 1979 on Ricaudy Reef flat and for collection of a few individuals in May 1980 on Croissant Reef flat.

The abundance was estimated at first indirectly, by assessing the catch per unit effort (CPUE) or number of specimens counted per diver per hour. Density was later measured on transects where the observer, swimming with a board equipped with a flow meter and a compass, counted all individuals within a width of 2 meters, a method used to survey populations of *Acanthaster planci* (Kenchington and Morton, 1976).




Fig. 1. Sampling sites.

During each sampling, approximately twenty *A. echinites* were collected. Total length (TL), total weight (TW), drained weight (DW) and gutted weight (GW) of individuals relaxed with magnesium chloride, were measured (Conand, 1981).

Direct microscopic observation of formalin-preserved gonads was used to examine the gonads. Sex was determined, diameter and length of the gonadal tubules recorded and oocyte size distribution established. Gonad index (GI) was expressed as a ratio of gonad weight to drained weight. The monthly mean of the gonad index and confidence limits at a probability level of 0.90 were calculated.

Mean sea water temperatures were calculated from daily measurements at the Noumea-Anse Vata coastal station (Fig. 1).

The size at first sexual maturity was obtained by plotting the percentage of individuals with developed or developing gonads against classes of drained weight.




Table 1. Biometric study: (n : sample size; R : range of the values; \bar{m} : mean; d.f. : degree of freedom; r : correlation coefficient).

Character	n	R	\bar{m}
TL (mm)	303	65 - 365	222
TW (g)	301	16 - 1380	435
DW (g)	299	11 - 580	274
GW (g)	303	9 - 405	200
Relation		d.f.	r




Fig. 3. The relation between (A) length and drained weight and (B) drained weight and gutted weight of *Actinopyga echinites*.




Fig. 4. Monthly percentages of males, females and undetermined *Actinopyga echinites*.

characteristics of the maturity stages of *A. echinites* are given in Table 2. Stages I and II respectively correspond to immature and resting individuals. During these stages sex can not be determined microscopically. It is impossible or very difficult to distinguish resting gonads from immature gonads.

There is a slight sexual dimorphism: testicular tubules are slightly longer and narrower than ovarian ones and the mature ovaries are pink and heavier than the whitish testes.

At stage IV, the gonad index values were 6.1 for the males and 7.8 for the females. The ripe oocytes measured approximately 175 μ m. The coefficient of fecundity, estimated by dividing the mean weight of ovaries in stage IV by the cube of the diameter, equals 4700. *A. echinites* is characterized by the relatively large size of oocytes and a low fecundity.

Table 2. Maturity stages (\bar{m} : mean; n: number of specimens; SD: standard deviation; L: length of tubules; ϕ : diameter).

Maturity stage	Undetermined				Males		Females	
	Gonad tubules L(mm)	Gonad tubules ϕ (mm. 10^{-1})	Gonad weight (g. 10^{-1})	GI	Gonad tubules L(mm)	Gonad tubules ϕ (mm. 10^{-1})	Gonad weight (g. 10^{-1})	GI
Stage I								
Immature	\bar{m} 10.0	2.3	1.7	0.12				
Stage II	n 49	37	54	54				
Resting	SD 9.7	1.4	1.3	0.14				
Stage III								
Growing	\bar{m} 65.5	5.0	28.4	1.08	47.9	6.7	30.2	0.99
	n 22	22	22	22	30	30	30	29
	SD 27.7	2.0	26.9	0.88	26.2	2.7	40.9	1.30
Stage IV								
Mature	\bar{m} 114.6	8.0	196.1	6.07	106.8	11.1	251.1	7.84
	n 42	42	44	42	38	38	39	39
	SD 29.0	3.1	136.7	3.19	25.8	3.9	189.3	5.29
Stage V								
Post spawning	\bar{m} 52.1	4.4	23.1	0.77	44.1	5.3	28.7	0.94
	n 53	41	53	53	52	38	53	52
	SD 30.9	1.6	25.3	0.72	27.1	3.9	44.9	1.40




Fig. 5. Monthly percentages of maturity stages of *Actinopyga echinites*.

At stage V of post-spawning, spicules were apparent in the gonadal wall. Monthly percentages were calculated for males, females and individuals of undetermined sex (Fig. 4). The overall sex ratio

was 1.0:1.0. Undetermined individuals represented 19% of the total. The percentage of these individuals was low from September to January and higher from April to August. This variation was related to the reproductive cycle.

Monthly percentages of maturity stages (Fig. 5) were similar in both males and females. Stage III was preponderant from July to October, stage IV from November to January. The post-spawning stage V lasted from February to July.

The gonad index cycle (Fig. 6) showed a growing phase from September to November-December, where the values were at their maximum. A rapid decline occurred in January and February in 1979 and in 1980, followed by a phase of low values. The high percentage of undetermined individuals from April to August corresponds to resting individuals.

By considering the gonad index cycle, the variations of maturity stages and the percentages of undetermined individuals all together, it was possible to define more precisely the different phases of the re-




Fig. 6. Monthly changes of gonad index of *Actinopyga echinites* and surface water temperature. The dotted area represents the confidence interval (0.90 probability level) of the mean gonad index.

- Trustees of the British Museum, London.
- Conand, C. 1979, Bêche-de-mer in New Caledonia : weight loss and shrinkage in three species of holothurians. Fisheries Newsletter, South Pacific Commission 19:14-18.
- Conand, C. 1981, Sexual cycle of three commercially important holothurian species (Echinodermata) from the lagoon of New Caledonia, Bull. mar. Sci. 31:523-544.
- Engstrom, N.A. 1980, Reproductive cycles of *Holothuria* (*Halodeima*) *floridana*, *H. (H) mexicana* and their hybrids (Echinodermata : *Holothuroidea*) in southern Florida, U.S.A., Int. J. Inv. Reprod. 2:237-244.
- Franklin, S.E. 1980, The reproductive biology and some aspects of the population ecology of the holothurians *Holothuria leucospilota* (Brandt) and *Stichopus chloronotus* (Brandt), Unpublished Thesis, University of Sydney.
- Harriot, V.J. 1980, The ecology of holothurian fauna of Heron Reef and Moreton Bay. Unpublished M.Sc. thesis, University of Queensland, 153 pp.
- Intes, A. & J.L. Menou 1979, Quelques holothuries (Echinodermata) des environs de Nouméa et leur répartition. Rap. Scient. et techn. (3), ORSTOM, Nouméa, 23 p.
- Kenchington, R.A. & B. Morton 1976, Two surveys of the crown of thorns starfish over a section of the Great Barrier Reef, Aust. Govt. Publ. Serv., Canberra, 186 p.
- Krishnaswamy, S. & S. Krishnan 1957, A report on the reproductive cycle of the holothurian *Holothuria scabra*, Jaeger, Curr. Sci. 36:155-156.
- Rowe, F.W. & J.E. Doty 1977, The shallow water holothurians of Guam, Micronesica 13(2):217-250.

THE INTERNATIONAL ECHINODERMS CONFERENCE / TAMPA BAY /
14-17 SEPTEMBER 1981


Echinoderms: Proceedings of the International Conference, Tampa Bay

Edited by

JOHN M. LAWRENCE

University of South Florida, Tampa

OFFPRINT



A.A. BALKEMA / ROTTERDAM



B 3955 ex. 1