

## A. INTRODUCTION

New Caledonia with the adjacent islands, situated some 1200 km north-  
west of Australia, is an exposed section of a highly complex orogenic

be of Montian (basal Palaeocene) age. Then sedimentary intercalations containing *Inoceramus* were found, but it seems very unlikely that these are autochthonous. Recently H. Gonord has observed that these volcanics are discordantly overlain by Eocene greywackes. The fact that they were emplaced on the west coast of the island at the same time as the M. Eocene gliding nappes, leads this author to believe they are allochthonous. The relationship between the basalt outcrops of the east and west coasts cannot yet be determined exactly, but it is possible that the former represent the zone of origin of the overthrust units. With this hypothesis one can explain why at various points on the west coast the basalts are in direct contact with Cretaceous formations and the intervening siliceous and calcareous Eocene I is absent.

The basalts are very often overlain by the great allochthonous, peridotitic massifs, whose derived fragments only appear in the L. Miocene

of the Nepoui region (Fig. 1) but the peridotites also lie on sedimentary and metamorphic rocks. The emplacement of the peridotites occurred late compared with the major phases of orogenesis. The basal surface of the large massifs is never affected by intense or tight folds; the angular discordance between this superstructure and the infrastructure is therefore very noticeable.

Metamorphism affected the sedimentary rocks older than Eocene II over large areas. A first phase of metamorphism occurred during the Cimmerian emergence in the central region of the island and a second (lawsonite, pumpellyite, epidote, glaucophane) occurred in the late Eocene in the northern regions. Among the major questions concerning the geology of New Caledonia, the analysis of metamorphic zones, their structure and the ages of the different phases must be looked at afresh.

The structure of New Caledonia remains little understood; mapping




Figure 1. Outline geology of New Caledonia and the Loyalty Islands.

in the east side is not advanced, since the area is thickly forested. Attention should be drawn, however, to an important aspect of the structure not emphasized in previous publications. A line parallel to the axis of the island separates zones 1 and 2. To the south-west occur great amounts of basalt, from Poum to near Bourail, covered by large peridotitic massifs whose basal surface is slightly inclined towards the south-west (Fig. 2). This is the zone which Routhier (1953) called 'sillon' (furrow) because the most recent formations occur there. In the central chain (Figs. 1 and 2), north-east of the division line, the rocks are everywhere older, often metamorphic and overfolded towards the south-west or the south; serpentinites and ultramafic masses occur, notably the Oua-Tilou and Tchinguou massifs (Fig. 1). The division line is certainly an old structural feature. It is effectively on this line that the most definite indications of Cimmerian emergence lie (Senonian conglomerates of the Congo and of the Mecouagna). In this same region, and over a distance of more than 50 km, the basalts directly

### C. DETAILS OF THE GEOLOGY OF THE ZONES

New Caledonia is described in terms of three zones. The West Caledonian Basin (zone 1) is characterized by ultramafic massifs and basalts resting on Mesozoic greywackes and Eocene rocks (see Table 1). In the Central Chain (zone 2) basalts are absent and the Cretaceous is non-metamorphic (see Table 2). The south-eastern belt (zone 3) is characterized by a very large ultramafic massif resting on sediments broadly similar to those of zone 1 (see Table 3).

The Loyalty Islands (zone 4) are a group of raised atolls which mark the position of a submarine, volcanic ridge. This NW.-SE. ridge lies 100 km north-east of New Caledonia and extends parallel to the latter for at least 800 km. The outline stratigraphical column of the Loyalty

<i>Rock unit</i>	<i>Age and evidence for age</i>	<i>Lithology</i>	<i>Thickness m</i>	<i>Comments</i>
'Formation à charbon'	U. Cret, perhaps Senon;	Rhyolitic tuff; carbonaceous beds; sandstones and arkoses with <i>Inoceramus</i> ; conglomerate	200	
	?Portland	Sandstone with gastropods		Unconformity: <i>Cimmerian phase</i> (Hokonui orogeny of New Zealand)
				Possible non-sequence during Kimm
'Formation des grau-wackes' (unmeta-	Call to Oxf, belemnites	Diorite and micro-gabbro; sandy greywacke; black shale		

Table 2  
STRATIGRAPHY IN ZONE 2

<i>Rock unit</i>	<i>Age and evidence for age</i>	<i>Lithology</i>	<i>Thickness m</i>	<i>Comments</i>
Ultramafic massifs	U. Eoc or Olig	Harzburgite, serpentinite	50-500	
Cretaceous strata	Cret, certainly Senon, fossils	Limestone, clay, sand; conglomerate	>100	Flat lying, unfolded and unmetamorphosed*
	?Portland	Sandstone with gastropods		Unconformity: <i>Cimmerian orogeny</i>
'Formation des grau-wackes'	Sinem to Oxf?	Fine-grained grey-wacke; black, chloritic shale; micrites	>1000	Much folded and metamorphosed (chlorite, mica, etc.)

<i>Rock unit</i>	<i>Age and evidence for age</i>	<i>Lithology</i>	<i>Thickness m</i>	<i>Comments</i>
		Ferruginous silt deposits	60-90	
		Coral reefs	30	Two phases of uplift and transverse faulting (Plioc to Quaternary); uplift greater in north-west of massif than in south-east
				Penetration
				Lateritization (Olig and Mioc) and development of Karst morphology
				Faulting and tilting of massif to south-west
Calc-alkaline	?Olig; intrude ultramafic	Dykes and veins of: feldspathic horn-		

Table 4  
STRATIGRAPHY IN THE LOYALTY ISLANDS

<i>Rock unit</i>	<i>Age and evidence for age</i>	<i>Lithology</i>	<i>Thickness m</i>	<i>Comments</i>
Raised terraces	Quaternary, madreporarians	Coral limestone terrace	0 to many metres	Discontinuous uplift of the southern part of the submarine volcanic arc is marked on the ancient uplifted atolls by a sequence of belts of coral platforms or erosion levels; there is no evidence of uplift in the north
				Unconformity: uplift
Fossil coral atolls on Maré, Lifou, Ouvéa	Neogene, madreporarians	Dolomitized coral atoll with madreporarians and Lithothamnions	Many hundreds of metres towards the margins of	






Figure 3. Hypothetical cross-section of Maré Island in the Loyalty Islands (by J. Launay & J. Recy).

can be locally very tilted (as on the southern border of the Boulinda massif). The positive gravity anomalies shown by Crenn (1953) in certain of the massifs in the north of the island can be explained in the same way.

(iii) *The large ultramafic massif in the south-east*: this massif covers about 4000 km<sup>2</sup> and rests upon Cretaceous formations or basalts and sedimentary rocks of Eocene age with either horizontal or gently dipping contact towards the north-east. Within it two lithologic units can be distinguished (Guillon 1969). *The main ultramafic mass* is more than 2.5 km thick and is composed throughout of forsteritic olivine (Fo 88-92), enstatite (En 90) and chromium spinel. It is little differentiated:

stratum but they are never remote from the floor contact of the massif. The occurrence of these rocks can be explained by injection of granitic material, originating from the sialic crust, later than the setting of the ultramafic massif. Late granites and quartz diorites are known in other islands of the Melanesian arc, especially in Neogene sedimentary rocks of the New Hebrides and in New Zealand Miocene formations.

After intrusion of the granitic rocks, the part of the massif located south-west of the big western fault was overthrust on the sedimentary substratum (e.g. in île Ouen). A late motion of this fault brings about a purely mechanical displacement of a part of the massif towards south or south-west. This could be the origin of the separation of the ultramafic bodies of the West Caledonian basin.

#### ACKNOWLEDGEMENTS

This paper has been written using the studies of the 'mission géologique de l'ORSTOM' 1946-9 (P. Routhier, A. Arnould, J. Avias) and those of the geologists at ORSTOM (J. H. Guillon, J. Launay, J. Recy, J. J. Trescases) and C.N.R.S. (H. Gonord) performed since 1966. It takes notice of data and even hypotheses resulting from recent works, often unpublished. The collaboration of H. Gonord, J. Launay and J. Recy is gratefully acknowledged.

#### E. REFERENCES

- ARNOULD, A. 1953. Phénomènes de métamorphisme dans le Nord-Est de la Nouvelle-Calédonie. *Pacific Sci. Congr. (Wellington)*, 7 (2), 142-6.
- AVIAS, J. 1958. Sur l'existence d'une phase tectonique hercynienne tardive ayant affecté les formations antétriasiques de la côte ouest de la Nouvelle-Calédonie. *C.r. hebdomadaire. Séances Acad. Sci., Paris*, 246, 136-7.
- 1959. Subdivisions et corrélations des terrains crétacés de la Nouvelle-Calédonie. *Int. geol. Congr.* 20 (El Sistema Cretácico), 2, 393-401.

- CRENN, Y. 1953. Anomalies gravimétriques et magnétiques liées aux roches basiques de Nouvelle-Calédonie. *Annls Géophys.* 9, 291-9.
- FROMAGER, D., GONORD, H. & GUILLON, J. H. 1967. Sur l'enracinement de certaines structures dans la région sud-ouest du bassin de Nouméa (Nouvelle-Calédonie). *C.r. somm. Séanc. Soc. géol. Fr.* 6, 242-3.
- GEOLOGICAL MAP OF NEW CALEDONIA (scale 1/100,000). 1952-63. ORSTOM (Paris) Sheets nos. 1-6 and no. 10, with explanation books.
- GONORD, H. 1968. Remarques sur les séries métamorphiques de la Nouvelle-Calédonie centrale; conséquence structurale et paléogéographique. *C.r. somm. Séanc. Soc. géol. Fr.* 9, 355.
- GONORD, H. & TRESCASES, T. T. 1970. Observations nouvelles sur la formation post-Miocène de Mueo (côte ouest de la Nouv.-Calédonie). *C.r. hebd. Séanc. Acad. Sci., Paris*, 270, 584-7.
- GUILLON, J. H. 1969. Données nouvelles sur la composition et la structure du grand massif péridotitique du Sud de la Nouvelle-Calédonie. *Cah. ORSTOM, sér. géol.* 1 (1), 7-25.
- 1969. Sur la fréquence des sulfures métalliques dans les massifs péridotitiques de la Nouvelle-Calédonie. *C.r. hebd. Séanc. Acad. Sci., Paris*, 268, 3013-14.
- GUILLON, J. H. & ROUTHIER, P. 1971. Les stades d'évolution des massifs péridotitiques de Nouvelle-Calédonie. *Bull. B.R.G.M.* IV, no. 2, 5-38.
- LACROIX, A. 1940. Les roches basaltiques de l'île Maré. *Bull. Soc. géol. Fr.* Séance du 9 mai 1940.
- LAUNAY, J. & RECY, J. 1970. Nouvelles données sur une variation relative du niveau de la mer dans toute la région Nouvelle-Calédonie—Îles Loyauté. *C.r. hebd. Séanc. Acad. Sci., Paris*. Séance du 13 avril 1970.
- LILLIE, A. R. 1970. The structural geology of lawsonite and glaucophane schists of the Ouégoa district, New Caledonia. *N.Z. Jl Geol. Geophys.* 13, 72-116.
- LILLIE, A. R. & BROTHERS, R. N. 1970. The geology of New Caledonia. *N.Z. Jl Geol. Geophys.* 13, 145-83.
- ORLOFF, O. & GONORD, H. 1968. Note préliminaire sur un nouveau complexe sédimentaire continental situé sur les massifs du Goa N°Doro et de Kadjitra (régions côtières à l'est de la Nouvelle-Calédonie). *C.r. hebd. Séanc. Acad. Sci., Paris*, 267, 5-8.
- ROE, DE G. 1964. Rubidium-Strontium analysis of ultramafic rocks and the origin of peridotites. M.I.T. Twelfth annual progress report, pp. 159-90.
- ROUTHIER, P. 1950. Sur la présence de formations liasiques en Nouvelle-Calédonie. *Cah. géol. Thoisy*, no. 3, 30.
- 1951. Sur les péridotites de la Nouvelle-Calédonie. *C.r. hebd. Séanc. Acad. Sci., Paris*, 233, 1204-7.
- 1953. Étude géologique du versant occidental de la Nouvelle-Calédonie entre le col de Boghen et la pointe d'Arama. *Mém. Soc. géol. Fr.* N.S. no. 67.
- TARLING, D. H. 1967. Results of a Palaeomagnetic reconnaissance of the New Hebrides and New Caledonia. *Tectonophysics*, 4, 55-68.
- TRESCASES, J. J. 1969. Premières observations sur l'altération des péridotites de Nouvelle-Calédonie. *Cah. ORSTOM, sér. géol.* 1 (1), 27-57.