

# Comparison of the Hydrographic Structure of Equatorial Waters North of New Guinea and at 170° E<sup>1</sup>

Philippe Hisard, Yves Magnier,  
and Bruno Wauthy

Centre O.R.S.T.O.M.  
B.P. 4  
New Caledonia

## ABSTRACT

The Equatorial Undercurrent was observed from March through August 1967 on a transect along 170°E during five cruises of the R.V. CORIOLIS. The velocity structure changed markedly from cruise to cruise. The minimum volume transport,  $15 \times 10^6 \text{ m}^3/\text{sec}$ , was observed in May; the maximum volume transport,  $55 \times 10^6 \text{ m}^3/\text{sec}$ , was observed in July. These changes were accompanied by equally marked changes in other hydrographic features. An attempt is made to compare and relate the observed variations along 170°E with the variations north of New Guinea, particularly in the North Equatorial Countercurrent.

*Introduction.* Tsuchiya (1968) has observed that, in the western Pacific, it is usually difficult to distinguish clearly the Equatorial Undercurrent from the North Equatorial Countercurrent north of it. Throughout the western Pacific Ocean these two currents are connected by a continuous layer of eastward-flowing water, but downstream (eastward) these currents are progressively farther apart (Montgomery 1962). It is of interest, therefore, to compare the hydrographic structure of the waters north of New Guinea with that of the waters at 170°E.

*The Hydrography North of New Guinea.* The pattern and strength of the surface currents north of New Guinea vary markedly during the year (Fig. 1). These variations are induced by meteorological changes associated with the monsoon rhythm. In July, the North Equatorial Countercurrent reaches its maximum strength when it is fed (i) by the Mindanao Current, which originates in the North Equatorial Current, and (ii) by the South Equatorial Current,

1. Accepted for publication and submitted to press 24 January 1969.

O. R. S. T. O. M.




Figure 1. Surface currents north of New Guinea. Upper part: summer; lower part: winter. (From Morskoï Atlas).

which flows along the northern coast of New Guinea up to Halmahera. In January, however, the Countercurrent is minimal in strength, since it receives water from only the north; the South Equatorial Current deteriorates in the face of increasing northwest winds and the Countercurrent loses water to the New Guinea Coastal Current. Thus, in July the North Equatorial Counter-

current is influenced by both northern and southern waters (the latter with relatively high salinity), but in January the Countercurrent is influenced by only the northern water (with decreased salinity along Mindanao and in the Celebes Sea).

Tsuchiya (1967) believes that the water mass north of New Guinea is the northern boundary of the Coral Sea water that passes through the Vityaz Strait. However, the lack of data from a southern station in July makes it impossible to determine whether the South Equatorial Current along the northern coast of New Guinea helps in the spreading of Coral Sea water.

According to Yamanaka et al. (1965), the major seasonal and long-term variations in the Countercurrent, in addition to the variations in transport, are the geographical positions of the Countercurrent's northern and southern edges; between  $130^{\circ}\text{E}$  and  $140^{\circ}\text{E}$ , the southern edge fluctuates between  $2^{\circ}\text{N}$  and  $4^{\circ}30'\text{N}$ . Yamanaka et al. (1965) in determining that the Countercurrent reaches its maximum transport in June–July, has confirmed Wyrski's (1961) observations and has provided evidence of the relative regularity in the annual variation in the volume transport. Burkov et al. (1960) have shown that the southern edge of the Countercurrent plays an important part in the interaction of the Countercurrent and the Undercurrent as the edge of the former sinks equatorward. These workers have also noted that the eastward flow of the Countercurrent is 700 km wide at the surface and 1200 km wide at 500 m; the interaction of these two currents, which are strengthened by their relative positions with depth, is not limited to the western Pacific but must be significant across the entire ocean because of the convergence observed at the surface along the southern edge of the Countercurrent across the entire ocean (Burkov 1963, 1966).

By considering the two equatorial sections occupied by the VITYAZ along  $142^{\circ}\text{E}$  in April 1958 and along  $140^{\circ}\text{E}$  in July 1957 (cruises 27 and 25; Burkov and Ovchinnikov 1960a), it is possible to demonstrate the variations previously described. During April and July, which represent opposite seasons, the surface circulation is different and typical. In both April and July (Figs. 2 and 3, respectively), the salinity distributions show the extension of a high-salinity tongue of South Pacific subtropical water across the equator to the north at 150 m. The northern boundary of this tongue constitutes a strong salinity gradient enclosed approximately by the 35‰ isohaline. In April, at 150 m, this boundary extends to  $3^{\circ}\text{N}$  whereas at 250 m it is close to the equator. However, in July, at a depth of about 120 m, the high-salinity tongue reaches to only  $2^{\circ}\text{N}$ ; but farther north, between  $5^{\circ}\text{N}$  and  $6^{\circ}\text{N}$ , at 120 m, there is an isolated high-salinity core that provides a higher salinity in the Countercurrent waters.

Figs. 2 and 3 also show the oxygen distributions, which illustrate the mixing processes at the southern edge of the Countercurrent. North of the equator, in April, when the convergence is weak due to the lesser flow of the Counter-




Figure 2. Salinity distribution (left) and oxygen distribution (right) along 142°E, April 1958. (VIRYAZ: cruise 27). Dotted areas indicate the 3.40 ml/l oxygen-content cores.

current, water of high oxygen content does not sink but forms a single core that remains at the same latitude as the salinity front; this oxygen core, with a salinity lower than 35.0‰, appears to originate from Countercurrent water. South of the equator, in April, at a depth of 200 m to 250 m, there is another core of high oxygen content that is richer than 3.40 ml/l; this core contains water having a salinity higher than 35.0‰. In July, at about 2°N, it is obvious that the subsurface oxygen isopleths slope downward and southward.

Fig. 4 shows the current-velocity structure obtained by direct measurements during April along 142°E; this permits clear identification of the Undercurrent, whose core has a velocity up to 20 cm/sec at a depth of 200 m. At this time, the North Equatorial Countercurrent, extending north to 2°N, flowed at speeds as high as 30 cm/sec at depths between 100 m and 200 m. Evidence for the existence of convergence between the two currents has been presented by Burkov and Ovchinnikov (1960b).

The eastward-flowing Undercurrent, with speeds greater than 10 cm/sec, has salinity values that range from 34.70 to 35.30‰, and the upper part of




Figure 3. Salinity distribution (left) and oxygen distribution (right) along 140°E, July 1957. (VIRYAZ: cruise 25).

the Undercurrent is associated with the northern core of oxygen at 2°N while the lower part is associated with the deep southern oxygen core. Thus, although the oxygen content of these two cores is the same, about 3.40 ml/l, their salinities are very different. It is possible that these two oxygen sources, with different origins but with the same value, could account for the oxygen homogeneity of the Undercurrent described by Knauss (1960).

In summary, it appears that the Undercurrent in the vicinity of 140°E, north of New Guinea, is composed of two different water masses as judged by the oxygen and salinity values. On the one hand, the Undercurrent is associated with the Countercurrent water in July when this mass is more oxygenated by mixing processes along the convergence at the southern edge of the Countercurrent. On the other hand, at the same time, the Undercurrent is associated with Coral Sea water that reaches the equatorial zone north of New Guinea. It is possible that the seasonal hydrographic variations that determine changes in the Countercurrent water have an effect upon the oxygen distribution in the upper northern part of the Undercurrent, with this variation in oxygen content being related to changes in the volume transport.




Figure 4. East-west velocity cross section at  $142^{\circ}\text{E}$  in April 1958. After Burkov 1960. Plus values are eastward velocities. Dotted line indicates an axis of convergence.

*The Hydrography along  $170^{\circ}\text{E}$ .* In 1966 and 1967, respectively, the R.V. CORIOLIS made four cruises in the "Bora" series and five cruises in the "Cyclone" series along transects at  $170^{\circ}\text{E}$ . The hydrographic data collected on these cruises have been published (Lemasson et al. 1967a and b, Magnier et al. 1967, Rotschi et al. 1967, Hisard et al. 1968, Jarrige et al. 1968, Magnier et al. 1968, Rotschi et al. 1968). The results from current meters will be published by the O.R.S.T.O.M. Centre of Noumea. On each cruise, the vessel occupied 10 hydrographic stations from  $20^{\circ}\text{S}$  to  $4^{\circ}\text{S}$  and 17 stations from  $4^{\circ}\text{S}$  to  $4^{\circ}\text{N}$ . The sections between  $4^{\circ}\text{S}$  and  $4^{\circ}\text{N}$  were completed in eight days in 1966 and in only four days in 1967 to minimize time variations; the equatorial stations were occupied on 11 March, 26 April, 6 June, 12 July, and 25 August.

At each hydrographic station, 24 samples from 0 to 500 m were collected and the 24 bottles were spaced to obtain close samples of the characteristic bathythermic structure. Temperature, salinity, oxygen, phosphate, nitrate and nitrite content were measured at each depth. In 1967, between  $4^{\circ}\text{S}$  and  $4^{\circ}\text{N}$ , direct current measurements were obtained with two current meters having Savonius rotors. These meters were spaced 1000 m apart on the same wire,

the wire angle being kept minimal at the surface. The current was then calculated as the vector difference between the reading on the upper meter (between 0 and 500 m) and the reading on the lower meter (assumed to be located




Figure 5. East-west velocity cross sections for the five cruises in March, April, June, July, and August 1967 at  $170^{\circ}\text{E}$ . Unit is cm/sec. Plus values are eastward velocities. Dotted lines indicate north-south divergences in the zone of connection.






Figure 6. Salinity distributions for the five cruises in March, April, June, July, and August 1967 at 170°E. Isohalines every 0.20 ‰. Dotted lines are isanosteres for every 100 cl/t.





Figure 7. Oxygen distribution for the five cruises in March, April, June, July, and August 1967 at 170°E. Isohalines every 0.20 ml/l. Hatched areas indicate the 3.40-3.60 ml/l oxygen-content layer. Dotted lines indicate the isanosteres for 200 and 300 cl/t.





In March, the salinity in the velocity core was lower than 35.0‰ and the volume transport was  $20 \times 10^6 \text{ m}^3 \text{ sec}$ . In July, the salinity in the core was 35.70‰, the volume transport was about  $55 \times 10^6 \text{ m}^3 \text{ sec}$  (nearly three times that in March), and the velocity reached 90 cm/sec. Simultaneously, oxygenated water richer than 3.40 ml/l was spreading progressively equatorward at a depth of about 200 m. By July this spreading had reached its maximal extent, and a secondary core, highly oxygenated, was present at the equator at a depth of 250 m.

The variations at 170°E can be related to the mean change in the North Equatorial Countercurrent, the transport of which, in the northern hemisphere, is minimal in the winter and maximal in the summer. Furthermore, in the winter the Countercurrent consists mainly of water from the North Equatorial Current, the salinity of which is lowered along the course of the Countercurrent through the Celebes Sea. In the summer, on the other hand, the salinity of the Countercurrent is increased by contributions from the South Equatorial Current.

*Acknowledgments.* This paper presents the preliminary results of research on the equatorial current system carried out by the O.R.S.T.O.M. Centre of Noumea, under the direction of Mr. H. Rotschi. It is based on the results of five cruises representing nearly five months working at sea for all our colleagues. We are also indebted to Dr. George L. Pickard, University of British Columbia, who took part in one of these cruises and was also most helpful in subsequent discussions.

#### REFERENCES

- BURKOV, V. A.  
1963. Water circulation in the North Pacific. *Okeanology*, 3 (6): 761-776.  
1966. Structure and nomenclature of Pacific Ocean currents. *Okeanology*, 6 (1): 3-14 (English translation).
- BURKOV, V. A., V. S. ARSENTYEV, and I. M. OVCHINNIKOV  
1960. To the notion of northern and southern tropical fronts in the ocean. *Trud. Inst. Okeanol. Akad. Nauk. SSSR*, 40: 108-120 (Russian).
- BURKOV, V. A., and I. M. OVCHINNIKOV  
1960a. Structure of zonal streams and meridional circulation in the central Pacific during the northern hemisphere winter. *Trud. Inst. Okeanol. Akad. Nauk. SSSR*, 40: 93-107 (Russian).  
1960b. Investigations of equatorial currents to the north of New Guinea. *Trud. Inst. Okeanol. Akad. Nauk. SSSR*, 40: 121-134 (Russian).
- HISARD, PHILIPPE, FRANÇOIS JARRIGE, and PIERRE RUAL  
1968. Résultats des observations physico-chimiques de la croisière "Cyclone 2" du N.O. "CORIOLIS". *Rapp. Sci. O.R.S.T.O.M. Centre de Nouméa, Sect. Oceanogr.*, 20; 21 p. (mimeo).

- JARRIGE, FRANÇOIS, JACQUES MERLE, and JACQUES NOEL  
1968. Résultats des observations physico-chimiques de la croisière "Cyclone 4" du N.O. "CORIOLIS". Rapp. Sci. O.R.S.T.O.M. Centre de Nouméa, Sect. Océanogr., 25; 22 p. (mimeo).
- KNAUSS, J. A.  
1960. Measurements of the Cromwell Current. *Deep-sea Res.*, 6 (4): 265-286.  
1966. Further measurements and observations on the Cromwell Current. *J. mar. Res.*, 24 (2): 205-240.
- LEMASSON, LIONEL, YVES MAGNIER, and BERNARD PITON  
1967a. Résultats des observations physico-chimiques de la croisière "Bora 1" du N.O. "CORIOLIS". Rapp. Sci. O.R.S.T.O.M. Centre de Nouméa, Sect. Océanogr., 9; 32 p. (mimeo).
- LEMASSON, LIONEL, BERNARD PITON, and FRANÇOIS JARRIGE  
1967b. Résultats des observations physico-chimiques de la croisière "Bora 3" du N.O. "CORIOLIS". Rapp. Sci. O.R.S.T.O.M. Centre de Nouméa, Sect. Océanogr., 11; 28 p. (mimeo).
- MAGNIER, YVES, PHILIPPE HISARD, FRANÇOIS JARRIGE, PIERRE RUAL, and BRUNO VOITURIEZ  
1967. Résultats des observations physico-chimiques de la croisière "Bora 4" du N.O. "CORIOLIS". Rapp. Sci. O.R.S.T.O.M. Centre de Nouméa, Sect. Océanogr., 12; 23 p. (mimeo).
- MAGNIER, YVES, HENRI ROTSCHI, BRUNO VOITURIEZ, and JACQUES MERLE  
1968. Résultats des observations physico-chimiques de la croisière "Cyclone 3" du N.O. "CORIOLIS". Rapp. Sci. O.R.S.T.O.M. Centre de Nouméa, Sect. Océanogr., 21; 23 p. (mimeo).
- MONTGOMERY, R. B.  
1962. Equatorial Undercurrent observations in review. *J. oceanogr. Soc. Japan*, 20th Anniversary Vol.: 29-40.
- ROTSCHI, HENRI, PHILIPPE HISARD, YVES MAGNIER, and JACQUES NOEL  
1967. Résultats des observations physico-chimiques de la croisière "Bora 2" du N.O. "CORIOLIS". Rapp. Sci. O.R.S.T.O.M. Centre de Nouméa, Sect. Océanogr., 10; 35 p. (mimeo).
- ROTSCHI, HENRI, and LIONEL LEMASSON  
1967. Oceanography of the Coral and Tasman seas. *Oceanogr. mar. Biol. Ann. Rev.*, 5: 49-97. Harold Barnes, Ed.
- ROTSCHI, HENRI, GEORGES PICKARD, PHILIPPE HISARD, and JACQUES CANNEVET  
1968. Résultats des observations physico-chimiques de la croisière "Cyclone 5" du N.O. "CORIOLIS". Rapp. Sci. O.R.S.T.O.M. Centre de Nouméa, Sect. Océanogr., 27; 23 p. (mimeo).
- SVERDRUP, H. U., M. W. JOHNSON, and R. H. FLEMING  
1942. *The Oceans*. Prentice Hall, N.Y. 1087 pp.
- TSUCHIYA, MIZUKI  
1967. Distribution of salinity, oxygen content, and thickness at 160 cl/t of thermocline anomaly in the intertropical Pacific Ocean. *Studies in tropical Oceanography*. 5; Proc. intern. Conf. trop. Oceanogr. 1965, Miami Beach, Florida; p. 37-41.  
1968. Equatorial circulation of the South Pacific Ocean. Proc. Ninth Gen. Meet. SCOR, South Pacific Symposium; p. 1-22.

## WYRTKI, KLAUS

1961. Physical oceanography of the south-east Asian waters. Scientific results of marine investigations of the South China Sea and the Gulf of Thailand. 1959-1961; Naga Rep., 2; 195 pp.

## YAMANAKA, HAJIME, NOBORU ANRAKU, and JIRO MORITA

1965. Seasonal and long-term variations in oceanographic conditions in the western North Pacific Ocean. Rep. Nankai Reg. Fish. Res. Lab., 22: 35-70.

## YOSIDA, S., H. NITANI, and N. SUZUKI

1959. Report of multiple ship survey in the equatorial region (I.G.Y.) Jan.-Feb. 1958. Hydrog. Bull., Tokyo, 59: 1-30.