

STUDIES ON MANGANESE-ACCUMULATING ALYXIA SPECIES FROM NEW CALEDONIA

Robert R. Brooks¹, Jennifer M. Trow¹, Jean-Marie Veillon², and Tanguy Jaffré³

Summary

Thirty-one species of *Alyxia* from New Caledonia were analysed for their manganese content. Most of the species showed excessive uptake of manganese with a maximum of 1.15% Mn in dried leaves of *A. rubricaulis*. The manganese content alone was sufficient to distinguish a number of species from each other. Composite samples were in addition analysed for calcium, magnesium, potassium and sodium. Correlation analysis and calculation of the slopes of regression lines showed that manganese uptake was mainly at the expense of calcium, rather than magnesium. Similarly magnesium was preferred to potassium and potassium to sodium. It is possible that manganese may have some physiological role in *Alyxia*, compensating to some extent for reduced uptake of the nutrients calcium and potassium.

Introduction

The manganese content of dried leaves of most plant species not growing over non-manganiferous substrates is typically of the order of 100 $\mu\text{g/g}$, whereas that of the soil is usually about 800-1,000 $\mu\text{g/g}$. Recently, Jaffré (1977, 1979) has found extremely high manganese levels in New Caledonia taxa growing on a wide variety of substrates. He reported (Jaffré, 1977) up to 3.2% manganese in dried leaves of *Maytenus bureauvianus* (Loes.) Loes. (Celastraceae) growing on basic hypermagnesian soils and 1.2% in *Alyxia rubricaulis* (H. Br.) Guillaumin growing on weathered acid soils. The respective manganese concentrations in the soils were 1.5% and




Fig. 1. Map of New Caledonia showing collection localities of *Alyxia* species. For numbering key see Fig. 2.

calcium, magnesium, potassium and sodium in order to study intraplant elemental relationships.

Materials and Methods

A total of 243 *Alyxia* specimens from 31 species held at the O.R.S.T.O.M. her-




Fig. 2. Geometric means and standard deviations for manganese concentrations in *Alyxia* species. Numbers to the left of each plot are the numbering code used in Fig. 1. Numbers in parentheses refer to number of specimens analyzed.

Fig. 2 shows the geometric means and standard deviation ranges for each species studied. These values alone give some differentiation of species. For example *A. caletoides* is distinguishable from 19 other species on the basis of these data, and *A. podocarpa* is also separated from the same number of taxa. Composites of each of 24 *Alyxia* (7 of the 31 species did not have sufficient material for analysis) were prepared and analysed for calcium, magnesium, manganese, potassium and sodium. Correlation coefficients were calculated for each pair of elements within each group (i.e. 3 pairs per group). The findings are summarized in Table 1.

Table 1. Statistical data for elemental concentrations in *Alyxia*.

Group	Element correlated with Mn	Significance	Slope of regression line
1. Ca-Mg-Mn	Ca	S**	-0.91
	Mg	S*	-0.08
2. K-Mg-Mn	K	S**	-0.77
	Mg	S**	-0.63
3. Mg-Mn-Na	Mg	S**	-0.73
	Na	S**	-0.93
4. Ca-Mn-Na	Ca	S**	-0.80
5. Ca-K-Mn	Ca	S**	-0.78
6. K-Mn-Na	K	S*	-0.53
	Na	S**	-0.79

S** Very-highly significant, $P < 0.01$.

S* Highly significant, $0.01 > P \geq 0.001$.

The table shows only relationship involving manganese and only those which are at least highly significant ($P < 0.01$). It is obvious that an increase in the manganese level in plant material must at the same time result in the diminution of the concentration of another constituent. This is clearly shown in the table by the negative values of the regression lines. Differences in the gradients of these lines do, however, give information of the relative competition between manganese and any other pair of elements. In group 1, manganese uptake among the 24 species of *Alyxia* is apparently mainly at the expense of calcium rather than magnesium. Similarly magnesium is preferred to potassium (group 2) and to sodium (group 3). In group 6 potassium is preferred to sodium.

The fall in relative levels of the nutrients calcium and potassium compared with magnesium may be due in part to the available levels of these elements in the substrate which tends to become progressively more ultrabasic as the manganese levels in the plants increase. There is, however, the possibility that manganese fulfills a physiological role in these plants and is able to replace calcium and/or potassium to some extent.

Although this work has been centred around *Alyxia*, there is no reason why a similar procedure should not be applied to other genera, particularly those which present severe classification problems. A good example of this is *Maytenus* (Celastraceae) which contains a number of New Caledonia species with even higher manganese levels than *Alyxia*. To date our work on this genus has been hampered by problems of identification of suitable reference material, but when these have been overcome, our studies will be extended further.

References

- Brooks, R. R., J. Lee, R. D. Reeves and T. Jaffré. 1977. Detection of nickeliferous rocks by analysis of herbarium specimens of indicator plants. *J. Geochem. Explor.* 7: 49-57.
- Jaffré, T. 1977. Accumulation du manganèse par des espèces associées aux terrains ultrabasiques de Nouvelle-Calédonie. *Compt. Rend. Acad. Sci. Paris Sér. D*, 284: 1573-1575.
- . 1979. Accumulation du manganèse par les Protéacées de Nouvelle-Calédonie. *Compt. Rend. Acad. Sci. Paris Sér. D*, 289: 425-428.