

TWO NEW SESQUITERPENE-ESTER ALKALOIDS FROM MAYTENUS MYRSINOIDES REISS.
(CELASTRACEAE)

Geneviève Baudouin, François Tillequin, and Michel Koch

Département de Pharmacognosie de l'Université René Descartes, E.R.A. au
C.N.R.S. N°950, Faculté des Sciences Pharmaceutiques et Biologiques
4, avenue de l'Observatoire F. 75006 PARIS (France)

Marie-Elise Tran Huu Dau and Jean Guilhem

I.C.S.N. du C.N.R.S., F. 91190 GIF-SUR-YVETTE (France)

Henri Jacquemin

Centre O.R.S.T.O.M. de Guyane, B.P. 165, 97305 CAYENNE (French Guyana)

Abstract - Two new alkaloids belonging to the dihydroagarofuran series, maymyrsine (1) and acetylmaymyrsine (2) have been isolated from the fruits of Maytenus myrsinoides Reiss. Their structures have been elucidated by M.S., ¹H NMR and X-ray diffraction analysis.


Maytenus myrsinoides Reiss. (Celastraceae) is a small tree growing throughout Guyana¹ and the northeastern region of Brazil². In a continuation of our studies of alkaloid containing plants from French Guyana³, we wish to report here the structural elucidation of two new alkaloids isolated from the fruits of this species⁴ and named maymyrsine and acetylmaymyrsine.

Maymyrsine (1) has been obtained as colourless prisms from ethanol, mp 185-187°C, $[\alpha]_D = +93^\circ$ (CHCl₃, C=1) (contents : 0.14 % from the dried plant material). The molecular formula has been established by high resolution mass spectrometry as C₃₂H₃₇NO₁₀ (Found : 595.2426 ; Calcd. : 595.2417). Substantial fragmentation ions suggested the presence of nicotinate (m/z = 124 and 106), benzoate (m/z = 105) and acetate (m/z = 43) units in the structure. In good agreement with these statements, the U.V. spectrum showed typical absorptions at $\lambda_{\max}^{\text{EtOH}}$ nm (log ε) : 230(4.13), 258 (sh., 3.51), 265(3.55), 271 (sh., 3.47) and 282 (sh., 3.06)⁵ and the I.R. spectrum at ν_{\max}^{KBr} cm⁻¹ = 725, 750, 1245, 1285, 1600, 1730, 1760, 2945 and 3400-3600, the last of which indicated the presence of a free alcoholic hydroxyl group. Moreover, the

^1H N.M.R. spectrum exhibits the characteristic signals of one nicotinate and one benzoate in the aromatic region and of two acetate groups (two 3H singlets at $\delta = 1.45$ and 1.73 ppm). Maymyrsine was thus a tetraester of a $\text{C}_{15}\text{H}_{26}\text{O}_5$ sesquiterpene pentaol belonging to the dihydroagarofuran^{5,6,7} series as shown by a 6H-singlet at $\delta = 1.39$ ppm ($-\text{O}-\text{C}(\text{CH}_3)_2-$) and a 3H-doublet ($J = 7\text{Hz}$) at $\delta = 1.26$ ppm ($\text{CH}_3-\text{CH}<$) on the ^1H N.M.R. spectrum and by the fragmentations at $m/z = 580, 553, 520, 431$ and 308 typical to this series⁵. The presence of a strong fragment at $m/z = 202$ instead of $m/z = 233$ generally encountered in the M.S. of Celastraceae alkaloids⁵ indicated the existence of a free primary alcoholic function at C-15. The oxygenation pattern on the dihydroagarofuran skeleton (C-1 eq., C-2 ax., C-6 eq. and C-9 ax.) has been determined by ^1H N.M.R., using double irradiation experiments (Table I). Unfortunately, the nature of the esterifying acids on these four positions could not be elucidated on N.M.R. basis because the ring effects created by benzoate and nicotinate groups are quite similar⁸.

We therefore decided to elucidate the structure by X-ray analysis. The crystals are monoclinic, space group $\text{P}2_1$ with $a = 14.058(7)$, $b = 8.652(5)$, $c = 13.606(7)$ Å and $\beta = 108.37(7)^\circ$; $Z = 2$, $V = 1570 \text{ \AA}^3$, $d_x = 1.18$. The experimental data were collected with a Philips 4-circle diffractometer, using graphite monochromated Cu K radiation ($\lambda = 1.5418 \text{ \AA}$). From 3170 measured independent reflexions, the intensities of 2584 of them were significant ($I > 3\sigma(I)$). The crystal structure was solved using local Direct Methods programs⁹. All hydrogen atoms but 3 were located from Fourier-differences syntheses. The refinement was performed using the large blocks least-squares method¹⁰ to a discrepancy factor of 6.3%. The identification of the nitrogen atom was made by interchanging the diffusion factors of C and N in the possible positions. A water molecule was included in the refinement with half occupation factor. The figure shows a perspective view of the molecule.

Acetylmaymyrsine (2) has been obtained as a colourless amorphous solid, $[\alpha]_D^{20} = +92^\circ$ (CHCl_3 , $C=1$) (contents: 0.44% from the dried plant material). The molecular formula has been established by high resolution mass spectrometry as $\text{C}_{33}\text{H}_{36}\text{NO}_{11}$ (Found: 637.2492; Calcd.: 637.2523). The U.V. spectrum: $\lambda_{\text{max}}^{\text{EtOH}}$ (log ϵ) = 230(4.12), 258 (sh. 3.51), 265(3.54), 271 (sh. 3.47), 282 (sh. 3.06) is very similar to that of maymyrsine. The I.R. spectrum differs from that of maymyrsine by the absence of typical OH absorption at $3400-3600 \text{ cm}^{-1}$. The ^1H N.M.R. spectrum (Table I) is also closely related to that of maymyrsine but exhibits the signals of three acetate





groups (three 3H singlets at $\delta = 1.35, 1.61$ and 1.91 ppm). Downfield shifts of H-15_a and H-15_b (4.52 and 5.72 instead of 4.13 and 4.62 ppm) and a strong shielding by 1,3-diaxial interaction of the signal of H-6ax. (6.10 instead of 6.51 ppm) clearly indicate that this second alkaloid is the acetic ester of maymyrsine at the C-15 position. Furthermore, acetylation of maymyrsine 1 ($\text{Ac}_2\text{O}/\text{C}_5\text{H}_5\text{N}/48$ h) leads to acetylmaymyrsine 2 in almost quantitative yield.

Maymyrsine and acetylmaymyrsine are novel alkaloids related to a new $\text{C}_{15}\text{H}_{26}\text{O}_5$ sesquiterpene pentaol. From a chemotaxonomic point of view, it is interesting to note that two structurally related sesquiterpene alkaloids had been previously isolated; cathedulin B8 from *Catha edulis* Forsk. (Celastraceae)⁷ and salacinin C from *Salacia brachypoda* Peyr. (Hippocrateaceae)⁵. This emphasizes the chemical homogeneity of the family Celastraceae and its close relationship with Hippocrateaceae^{11,12}.

TABLE I

¹H N.M.R. spectra of maymyrsine (400 MHz) and acetylmyrsine (270 MHz) C₆D₆,TMS, δ (ppm).

	Maymyrsine <u>1</u>	Acetylmaymyrsine <u>2</u>
Me-14	1.26, 3H, d, J=7Hz	1.14, 3H, d, J=7Hz
Me-12 and Me-13	1.39, 6H, s	1.30 and 1.32, 2x3H, s
Ac	1.45 and 1.73, 2x3H, s	1.35, 1.61 and 1.91, 3x3H, s
H-3(ax.)	1.65, 1H, dd, J=17Hz, J'=1Hz	not individualized
H-4(eq.)	2.28, 1H, qdd, J=7Hz, J'=6Hz, J''= 1Hz	2.17, 1H, qdd, J=7Hz, J'=6Hz, J''= 1Hz
H-3(eq.)	2.46, 1H, ddd, J=17Hz, J'=6Hz, J''= 3Hz	2.39, 1H, ddd, J=17Hz, J'=6Hz, J''= 3Hz
H-8(eq.)	2.67, 1H, ddd, J=16Hz, J'=7Hz, J''= 3Hz	2.51, 1H, ddd, J=16Hz, J'=7Hz, J''= 3Hz
H-15 _a	4.13, 1H, d, J= 13Hz	4.52, 1H, d, J=13Hz
H-15 _b	4.62, 1H, d, J=13Hz	5.72, 1H, d, J=13Hz
H-9(eq.)	5.94, 1H, d, J=7Hz	5.83, 1H, d, J=7Hz
H-1(ax.)	6.12, 1H, d, J=3.5Hz	6.14, 1H, d, J=3.5Hz
H-2(eq.)	6.20, 1H, dd, J=3.5Hz, J'=3Hz	6.24, 1H, dd, J=3.5Hz, J'=3Hz
H-6(ax.)	6.51, 1H, s	6.10, 1H, s
Nicotinoyl H-5'	6.78, 1H, dd, J=8Hz, J'=5Hz	6.73, 1H, dd, J=8Hz, J'=5Hz
Benzoyl	7.15-7.37, 5H, m	7.12-7.38, 5H, m
Nicotinoyl H-4'	8.17, 1H, dt, J=8Hz, J'=1.5Hz	8.30, 1H, dt, J=8Hz, J'=1.5Hz
Nicotinoyl H-6'	8.38, 1H, dd, J=5Hz, J'=1.5Hz	8.44, 1H, dd, J=5Hz, J'=1.5Hz
Nicotinoyl H-2'	8.63, 1H, d, J=1.5Hz	8.53, 1H, d, J=1.5Hz



REFERENCES AND NOTES

1. A. Lemée, "Flore de la Guyane française", Lechevallier éditeur, Paris, 1952, 2, p.305.
2. S. Reissek, in C.F.P. de Martius, "Flora brasiliensis", Fried. Fischer, Lipsiae 1861, 11, 1, col.21, nr.40.
3. G. Baudouin, F. Tillequin, M. Koch, M. Vuilhorgne, J.Y. Lallemand and H. Jacquemin, J. Nat. Prod., 1983, 46, 681.
4. The plant material has been collected at Trois Sauts (French Guyana) in march 1976. Herbarium samples (HJ 1817) are kept in the herbaria of the Centre ORSTOM de Cayenne and of the Museum National d'Histoire Naturelle (Paris).
5. Ok-Pyo Chi, "Strukturaufklärung neuer Terpenesteralkaloide aus Salacia brachypoda Peyr., Hippocrateaceae!" Dissertation, München, 1981, pp.1-97.
6. R.M. Smith in R.H.F. Manske, "The Alkaloids", Academic Press, New York, 1977, 16, pp.215-248.
7. R.L. Baxter, W.M.L. Crombie, D.J. Simmonds and D.A. Whiting, J.Chem. Soc. Perkin Trans. I, 1979, 2972.
8. R.L. Baxter, W.M.L. Crombie, L. Crombie, D.J. Simmonds, D.A. Whiting and K. Szendrei, J. Chem. Soc. Perkin Trans. I, 1979, 2982.

9. C. Riche "7th European Crystallographic Meeting, Collected Abstract", Jerusalem, 1982, p.25.
10. G.M. Sheldrick, "Shelx 76, Program for Crystal Structure Determination", University of Cambridge (England), 1976.
11. Th. Lösener, "Celastraceae" in A. Engler and K. Prantl, "Die natürlichen Pflanzenfamilien", W. Engelmann, Leipzig, 1892, III/5, pp.189-222.
12. Th. Lösener, "Hippocrateaceae" in A. Engler and K. Prantl, "Die natürlichen Pflanzenfamilien", W. Engelmann, Leipzig, 1892, III/5, pp.222-230.

Received, 4th June, 1984

HETEROCYCLES
An International Journal
for Reviews and
Communications
in Heterocyclic
Chemistry

**TWO NEW SESQUITERPENE-ESTER ALKALOIDS FROM
MAYTENUS MYRSINOIDES REISS. (CELASTRACEAE)**

Geneviève BAUDOUIN, François TILLEQUIN, and Michel KOCH

Département de Pharmacognosie de l'Université René Descartes, E.R.A. au
C.N.R.S. N° 950, Faculté des Sciences Pharmaceutiques et Biologiques
4, avenue de l'Observatoire F. 75006 PARIS, France

Marie-Elise Tran Huu DAU and Jean GUILHEM

I.C.S.N. du C.N.R.S., F. 91190 GIF-SUR-YVETTE, France

Henri JACQUEMIN

Centre O.R.S.T.O.M. de Guyane, B.P. 165, 97305 CAYENNE (French Guyana)

Reprinted from HETEROCYCLES 22, 2221 (1984)

16 AVRIL 1985

C.N.S.T.O.M. Fonds Documentaire

N° : 17 324, ex 1

Cote : B