

## 5. Free-living blue-green algae in tropical soils

P.A. ROGER and P.A. REYNAUD

### 1. Introduction

One result of the fertilizer price increase during the last decade is a renewed interest in biological  $N_2$  fixation as a means of reducing the use of N fertilizer. However, biological  $N_2$  fixation requires energy generally obtained by the catabolism of photosynthetically fixed carbon (photosynthate). Among the  $N_2$ -fixing microorganisms, only blue-green algae (BGA) are able to generate their own photosynthate from  $CO_2$  and water. This trophic independence makes BGA especially attractive as a biofertilizer. The agronomic potential of BGA was recognized in 1939 by De [14], who attributed the natural fertility of tropical paddy fields to  $N_2$ -fixing BGA. Since rice forms the staple diet of a high proportion of the world's population in areas where N fertilizer is rarely available, research on BGA in tropical soils has been focussed mainly on the paddy field ecosystem. Relative to the amount known about paddy soils, very little is known about other soils.

### 2. Occurrence of BGA in tropical soils

Blue-green algae (BGA) are photosynthetic prokaryotic microorganisms, some of which are capable of  $N_2$  fixation. This resulting trophic independence which has already been stressed, and a great adaptability to environmental factors should enable BGA to be ubiquitous. However Watanabe [81] and Watanabe and Yamamoto [84] found that  $N_2$ -fixing BGA are not present in every environment: of 911 samples only 46 (5%) harbored  $N_2$ -fixing species. This surprisingly low value is probably due to unsuitable methodology and to the small size of the samples. Watanabe's results suggested that  $N_2$ -fixing BGA grow more abundantly in tropical and subtropical regions than in temperate and subtemperate regions.

#### 2.1. Upland soils

Upland soils in arid climates are probably very inhospitable to many microorganisms because the temperatures are high and water is severely limited. BGA are especially resistant to such adverse conditions; thus they are the dominant components of the microflora in many cases [17]. A study of the savanna soils in the Congo indicated that the flora was almost entirely composed of BGA [10];  $N_2$ -

*Y.R. Dommergues and H.G. Diem (eds.), Microbiology of Tropical Soils and Plant Productivity. ISBN 90 247 2624 7.*

© 1982 Martinus Nijhoff/Dr W. Junk Publishers, The Hague/Boston/London.

fixing BGA have been reported to develop profusely in sugar cane and maize fields in India [63]. Large populations of *Calothrix* sp. in pearl millet fields and of *Gloeo-trichia* sp. in sugarcane fields were found by the authors in Senegal.

A qualitative study of the algal flora of dried soil samples from experimental upland fields (pH 7.8–8.3) at IARI, New Delhi, indicated that BGA were dominant in all the soil samples, Chlorophyceae were poorly represented and Xantophyceae were absent. Among the BGA, numerous  $N_2$ -fixing forms were observed [16]. Soil algae from regions around the Gulf of Mexico and areas in Ecuador and Colombia were studied by Durrell [15]. In 120 samples he found 62 species of algae; 46 of these species belonged to BGA. About half of the samples contained  $N_2$ -fixing species; *Nostoc muscorum* was observed in 21% of the samples and *Nostoc paludosum* in 13%. Other  $N_2$ -fixing species were observed in less than 4% of the samples.

## 2.2. Paddy soils (submerged soils)

The paddy field ecosystem provides a favourable environment for the growth of BGA with respect to their requirements for light, water, high temperature and nutrient availability. This could be the reason BGA grow in higher abundance in paddy soils than in upland soils [84] as reported in the widely different climatic conditions of India [35] and Japan [36]. By pooling data obtained in Senegal it appears that  $N_2$ -fixing BGA were recorded in 86 out of 89 paddy soils [48]. However, Venkataraman [78] pointed out that 'contrary to general belief,  $N_2$ -fixing BGA are not invariably present in tropical rice soils, and that an all India survey showed that out of 2213 soil samples from rice fields, only about 33% harboured  $N_2$ -fixing forms'. The heterogeneous and sometimes limited distribution of  $N_2$ -fixing BGA is still not well understood because no systematic analysis has correlated the presence or absence of BGA with environmental factors [30].

## 2.3. Quantitative evaluations

The lack of satisfactory methods for estimating biomasses of the different algal groups [17] has certainly limited ecological studies of soil algae. Plating techniques, most frequently used, are advantageous in providing qualitative and quantitative results simultaneously; however the accuracy of the count depends on the reliability of the particular dilution method. Filamentous forms are hard to separate into individual cells whereas moniliform filaments, which are easily separated, may give inflated figures of abundance [17].

Algal enumerations are often limited by an inadequate sampling methodology. Most of the results are expressed as numbers of algae per gram of soil, which do not take into account algae present in the floodwater of submerged soils and do not permit any extrapolation at the field level (what is the dry weight of soil colonized

by algae in one hectare of a paddy field?). A more satisfactory way to evaluate algal population is to determine the number of algae per  $\text{cm}^2$  by using core samples with a well-defined diameter, each core sample including the first centimeter of soil and the corresponding floodwater column [52]. Such a procedure allows comparisons and extrapolations at the field level. Roger and Reynaud [55] found that the distribution of soil algae is log-normal (logarithms of numbers are normally distributed) and that many samples are required to obtain a significant evaluation. For example, the mean value of *Anabaena* sp. biomass based upon 40 samples (each sample being obtained by mixing 10 sub-samples) taken in a 0.25 ha paddy field still exhibited a confidence interval of +32% and -27% of the mean. One fact at least is well established: in paddy soils (see Roger and Kulasooriya [58]), BGA numbers vary within large limits, from a few to  $10^7$  units  $\text{g}^{-1}$  dry soil. In upland soils lower values ranging from a few to  $10^6$  units  $\text{g}^{-1}$  dry soil have been generally reported [3, 25, 69, 71]. Reports on biomass are scarce. In paddy fields the biomass of BGA can reach values of several tons per hectare [58]. The little data available suggest that BGA seem to develop more profusely in submerged soils than in other cultivated soils. An exceptionally high total algal biomass of about  $40 \text{ t ha}^{-1}$  was recorded by Reynaud and Roger [47] in a sandy soil in Senegal that was spontaneously watered by a permanent spring; such a soil exhibited a water regime related more to that of submerged soils than to that of upland soils.

### 3. Ecology

#### 3.1. Physical factors

3.1.1. *Light.* Algae, as phototrophic microorganisms, are restricted to the photic zone and usually located in the upper 0.5 cm horizon. Yet algae also exist in deeper horizons, in a dormant condition as spores or filament fragments [10]. Light availability for soil algae depends upon the season and latitude, the cloud cover, the plant canopy, the vertical location of the algae in the photic zone and the turbidity of the water. Light intensity reaching the soil may vary from too low to excessive levels (10 to 110,000 lux).

In cultivated soils the screening effect of a growing crop canopy appears to cause a rapid decrease of light reaching the algae. Thus the canopy of transplanted rice decreased light by 50% when plants were 15 days old, 85% after one month and 95% after two months [29]. In Senegal, diatoms and unicellular green algae developed first and BGA developed later when the plant cover was dense enough to protect them from excessive light intensities, higher than 80 klux at 13:00h (Fig. 1); the  $\text{N}_2$ -fixing algal biomass and the density of the plant cover were positively correlated [52].

In the laboratory, after one month of incubation of a submerged unplanted soil under a range of screens, BGA were dominant in the most heavily shaded one, and green algae and diatoms were dominant in the soil exposed to full sunlight [45]. A




Fig. 1. Variation in biomass of different components of the algal flora during a vegetative cycle of rice in Senegal [52].

beneficial effect of the plant canopy shading the algae was also reported by Singh [63] in sugarcane fields, maize fields and grasslands in India.

As BGA are generally sensitive to high light intensities, they develop various protective mechanisms against it, namely:

- vertical migrations in the water of submerged soil;
- preferential growth in more shaded zones like embankments, under or inside decaying plant material [27] or a few millimeters, below the soil surface [17];
- migration into shaded zones (photophobotaxis) and aggregation providing a self-shading effect (photokinesis [46]);
- stratification of the strains in algal mats where  $\text{N}_2$ -fixing strains grow under a layer of eukaryotic algae more resistant to high light intensities [47].

However, some strains of BGA seem more resistant to high light intensities. *Cylindrospermum* sp. developed large biomasses in a harvested paddy field in Mali where light intensity was higher than 100 klux at 13:00 [72]. *Oscillatoria princeps* was also reported to grow profusely in full sunlight [46].

On the other hand, light deficiency may also be a limiting factor. In Japan, available light under the canopy was below the compensation point of the phytoplankton

during the second part of the cycle [22]. In the Philippines, during the wet season when light was moderate acetylene reducing activity (ARA) was higher in bare soil than in planted soil [85].

**3.1.2. Temperature.** The optimal temperature for BGA growth is about 30–35 °C which is higher than that for the growth of eukaryotic algae. In submerged soils daily variations in the temperature are moderated by the buffering effect of flood-water; temperature is rarely a limiting factor for BGA in paddy fields, because the range of temperatures permitting the growth of BGA is larger than that required by rice; however, it influences both algal biomass composition and productivity. Low temperatures decrease productivity and favour eukaryotic algae. High temperatures favour both the phytoplankton productivity and BGA [56].

Daily changes in the temperature are more drastic in terrestrial habitats than in aquatic environments [42]. An inhibitory effect of high temperature was observed by Jones [24] in the Kikuyu grasslands in Africa where algal N<sub>2</sub> fixation was higher on overcast days than on hot sunny days. Stewart [65] indicated a correlation between the algal ARA response to temperature and the temperature of the habitats from which the algae were collected. For many tropical species, ARA is optimum between 30–35 °C, but a *Nostoc* sp. isolated from the algal crust on a sandy soil in Senegal still exhibited significant ARA at 60 °C (Reynaud, unpublished). High temperatures occurring in the surface of tropical upland soils may have a selective action on the algal flora, favouring BGA which are more tolerant to high temperatures than eukaryotic algae. For example, the dry spores of *Nostoc* sp. can tolerate 2 minutes of 100 °C, the wet spores 20 minutes at 60–70 °C, and the vegetative filaments 10 minutes at 40 °C [10].

**3.1.3. Desiccation and remoistening.** Algal growth is hindered by intermittent desiccation periods which occur during the dry season and even during drought periods that occur in the rainy season. BGA have a high capacity to withstand desiccation. *Nostoc muscorum* and *Nodularia harveyana* were isolated from a soil that had been dry for 79 years [10]. Resistance to desiccation has been attributed to various characteristics [41], namely, with respect to fatal plasmolysis, the lack of cell vacuoles, the ability of some genera to quickly take on an encysted form, the presence in some genera of a mucilaginous sheath that absorbs water quickly and retains it. This latter characteristic could explain the dominance of mucilaginous colonies of *Nostoc* spp. and *Cylindrospermum* spp. in the paddy field during the last part of the cultivation cycle when the soil dries [38, 45, 72]. The dominance of BGA comprised only about 30% of the algal flora [32], whereas in Senegal, where the dry season lasts about 8 months, spores of heterocystous BGA constituted more long periods of desiccation than to the ability to remain active at low humidities [8]. The floristic composition of desiccated soils can be related to the dryness of the biotope. In a paddy in Italy, where the dry period is relatively short, N<sub>2</sub>-fixing BGA comprised only about 30% of the algal flora [32], whereas in Senegal, where the dry season lasts about 8 months, spores of heterocystous BGA constituted more

than 95% of the algal flora at the end of the dry period. In Uttar Pradesh (India), a large number of Chlorophyceae occurred in low-lying fields, whereas BGA were found in larger numbers in paddies at higher elevations [38]. In arid soils, BGA have been reported as dominant species [31] and sometimes as the only species present [5, 10].

### 3.2. Biotic factors

Organisms that limit BGA growth are pathogens, antagonistic organisms and grazers. Of these, only grazers have been documented. The development of zooplankton populations, especially cladocerans, copepods, ostracods, mosquito larvae, etc. prevented the establishment of algal blooms within one or two weeks [73]. Snails form another group of algal grazers in submerged paddy fields: the biomass of snails can be as high as  $1.6 \text{ t ha}^{-1}$  in certain rice fields in the Philippines which explains the low population of algae [58].

### 3.3. Soil properties

Among the soil properties, pH is the most important factor determining the algal flora composition. In culture media the optimal pH for BGA growth seems to range from 7.5 to 10.0 and the lower limit is about 6.5 to 7.0 [20]. Under natural conditions BGA grow preferentially in environments that are neutral to alkaline; which explains that in paddies correlations occur between:

- water pH and BGA number [37];
- soil pH and number of spores of  $\text{N}_2$ -fixing BGA in the soil during the dry season [18];
- soil pH and BGA growth [36];
- soil pH and the  $\text{N}_2$ -fixing algal biomass (however, this relationship was conspicuous only in samples homogenous for stage of rice development, fertilization and plant cover density [53]).

The beneficial influence of high pH on BGA growth is further demonstrated by the fact that the addition of lime increases BGA growth and  $\text{N}_2$  fixation [58]. However, the presence of certain strains of BGA in soils with pH values between 5 and 6 have been reported. Durrel [15] demonstrated the presence of *Nostoc muscorum* and *Anabaena torulosa* in soils with pH ranging from 5 to 7. *Aulosira fertilissima* and *Calothrix brevissima* have been reported to be ubiquitous in Kerala rice fields with pH from 3.5 to 6.5 [1]. The development of a dense algal bloom on an acidic soil (pH 5.5) was observed after the surface application of straw [33]. Stewart [66] also reported that some tropical BGA exhibited ARA even at pH 4. The poor growth of  $\text{N}_2$ -fixing BGA, frequently observed in acidic soils, is probably due to the inability of BGA to compete with Chlorophyceae, which are favoured by acidic conditions.

### 3.4. Algal successions

The factors mentioned above are responsible for the algal successions that have been reported. An example of algal succession is provided by a study conducted in paddies in Senegal [52, 53]. Soils are acidic, and have an average pH value of 5.0 at the beginning of rice cultivation and 6.2 after 2 months of submersion. The rainy season is short (15 July–15 November) and rice fields are dry the rest of the year. High light intensities (70–80 klux) occur throughout the year. During the early part of the cultivation cycle (planting and tillering), the algal biomass increased and consisted mainly of diatoms and unicellular green algae (Fig. 1). From tillering to panicle initiation, the algal biomass reached its highest values, and filamentous green algae and non-N<sub>2</sub>-fixing BGA were dominant. After panicle initiation the total biomass decreased; if the plant cover was sufficiently dense, heterocystous BGA developed, but if it was thin, filamentous green algae and homocystous BGA remained dominant. The observed variations in the algal flora were attributed to a decrease in light intensity and the N level resulting from the rice growth, and to an increase in pH value favouring BGA growth.

## 4. N<sub>2</sub> fixation

BGA are the only N<sub>2</sub>-fixing microorganisms that exhibit a higher plant type of photosynthesis. Their N<sub>2</sub>-fixing ability was first related to the presence of specialized non O<sub>2</sub>-evolving cells called heterocysts in which the nitrogenase, a highly O<sub>2</sub>-sensitive enzyme, is protected from O<sub>2</sub>. It is now clearly demonstrated that the ability to fix N<sub>2</sub> is not confined to heterocystous BGA, but is also found in a number of BGA, which were not known to fix N<sub>2</sub> a few years ago, and which can fix N<sub>2</sub> under microaerobic or anaerobic conditions [67]; over 125 strains are presently known to fix N<sub>2</sub>.

ARA is a reliable method for quantitative studies provided that one takes into account the following facts:

- the distribution of ARA is log-normal [54];
- N<sub>2</sub> fixation by BGA occurs not only in aerobic but also in anaerobic conditions [67];
- there are diurnal variations in ARA and also variations in ARA throughout the cultivation cycle. In submerged soils the variations exhibit typical patterns with one or two maxima. The occurrence of two maxima results from the inhibitory effect of excessive light during the middle of the day [45]. Similar variations were observed in upland soils but the inhibitory decrease of ARA during the middle of the day was attributed to an increase in temperature of the surface soil [24, 44]. Moreover, one should note that the duration of the incubation should be short [13] and that the conversion factor C<sub>2</sub>H<sub>4</sub>:N<sub>2</sub> reduced should be measured in each situation [40].

In submerged soils, light intensity appears to be the main factor governing

variations in photosynthetic  $N_2$  fixation, whereas in upland soils water availability becomes the main factor. Studying photosynthetic  $N_2$  fixation in a sandy soil colonized by a thick algal mat, Reynaud and Roger [47] observed that the variations of both total algal biomass and ARA were governed not only by light but also by the distribution of precipitations.

The average of 38 quantitative evaluations of  $N_2$  fixation in paddies recorded in the literature was  $27 \text{ kg N ha}^{-1}$  per crop [58]. The highest recorded value was  $50\text{--}80 \text{ kg N ha}^{-1}$  per crop [72].

In the savanna lands of southwest Nigeria,  $N_2$ -fixing Cyanobacteria such as species of *Scytonema*, *Tolypothrix* and *Nostoc* develop during the rainy season, particularly in previously burnt areas and may contribute significantly to the input of total nitrogen. Similarly in the open 'campina' areas among parts of the rain forests of tropical Brazil,  $N_2$ -fixing growth of blackish Stigonematales may be found [66]. In some conditions  $N_2$ -fixing BGA may significantly contribute to the N status of cultivated upland soils. Singh [63] reported a profuse growth of  $N_2$ -fixing BGA in sugarcane fields in India from beginning of the rainy season. Among the strains *Cylindrospermum licheniforme* was assumed to add about  $90 \text{ kg N ha}^{-1}$  in about 75 days. The same species was reported to very successfully colonize the clean soil surface of the weed-free maize fields. In grasslands of the Banaras Hindu University a thick and sometimes continuous BGA mat developed during the rainy season. The thicker the grass cover, the better the growth of BGA was. In protected and enclosed grasslands the total N content of the soil increased 88% in 5 months, an increase attributed mainly to the growth of *C. licheniforme*. From the above results it appears that BGA may contribute to maintain the N level of natural and cultivated ecosystems, but generally the extent of this contribution is poorly documented.

## 5. Relations between BGA and higher plants

### 5.1. Availability of fixed $N_2$ to higher plants

$N_2$  fixed by BGA is released either through exudation or through microbial decomposition after the death of the cells, but the process involved in the transfer of fixed  $N_2$  to the plants is largely a mystery [30]. It has been shown in the laboratory, that BGA liberate large portions of their assimilated nitrogenous substances [17, 70]. However, the large amounts of nitrogenous substances recorded may result from methodological artifacts such as the osmotic shock occurring when resuspending the cells or any other physical damage of the algal material. No information on the exudation of fixed  $N_2$  by BGA under field conditions is available, but it is clear that only part of it is available to rice, other parts being reincorporated by the

microflora or volatilized. Nutrients released through microbial decomposition after the death of the algae appear to be the principal manner in which N is made available to the crop. Field experiments with *Tolypothrix tenuis* for 4 consecutive years indicated that only 1/3 of the field algae was decomposed and absorbed by rice plants in the first year: the rest remained as residual soil N, which could have been responsible for continued yield increases in the succeeding years [80]. The transfer of N from algae to higher plants has been investigated using  $^{15}\text{N}$ -tracer techniques [34, 64]. Wilson (personal communication) recently recovered from a rice crop 39% and 51% of the N from  $^{15}\text{N}$ -labeled *Aulosira* spp. spread on or incorporated into the soil, thus showing that BGA N is readily available to rice. A similar experiment is under way at the International Rice Research Institute (Philippines). Results of the analysis of the first crop show a recovery of 13% and 38% of the N from  $^{15}\text{N}$ -labeled *Nostoc* sp. spread on or incorporated into the soil. This indicates that the availability of algal N for the plant varies either with the strain or with the physiological state of the algal material; i.e. N from an algal material rich in akinetes, not so easily decomposable, will be less available than that from vegetative cells which are more susceptible to decomposition.

### 5.2. Growth-promoting effect

Besides increasing N fertility, BGA have been assumed to benefit higher plants by producing growth-promoting substances. This hypothesis is based on the additive effects of BGA inoculations in the presence of nitrogenous fertilizers. Most of these results have been obtained with rice but similar results were observed also with vegetables such as radishes and tomatoes [51].

More direct evidence of hormonal effects has come primarily from treatments of rice seedlings with algal cultures or their extracts. Presoaking rice seeds with BGA cultures or extracts enhances germination, promotes the growth of roots and shoots, and increases the weight and protein content of the grain [19, 23]. It has also been established that algal growth-promoting substances are beneficial to other crops besides rice [11, 26, 39] and that the production of such substances is not confined to BGA. Whether these substances are hormones [19], vitamins [73], aminoacids [7] or any other components is still unknown.

### 5.3. Harmful effects

Blooms caused by filamentous algae can be harmful to rice, mainly due to a mechanical effect on the young plants [6]. However detrimental effects [49,43,53,59] of BGA are incidental [12, 58]. Even when BGA produced a bloom at the beginning of the growing cycle, their effect on grain yield was rarely negative [9].

#### 5.4. Epiphytism

Epiphytic BGA have been observed on wetland rice [57] deepwater rice [28] and on weeds growing in rice fields [27]. In wetland rice fields, epiphytic BGA on rice and weeds make a limited contribution to the N input but are important in providing an inoculum for the regeneration of the algal blooms that are periodically affected by adverse conditions. In deepwater rice, which offers a much greater biomass for colonization, the addition of N by epiphytic BGA is of agronomical significance [28]. BGA were found to grow preferentially on submerged decaying tissues. An endophytic growth inside the leaf sheath was also observed in deepwater rice. The results obtained did not confirm the existence or absence of biotic relationships between the algae and their hosts, but indicated that a mechanical effect in relation to the roughness of the support was involved in algal epiphytism and endophytism.

### 6. Role of BGA in soil colonization

Despite a worldwide distribution of BGA in terrestrial habitats their ecology has been less studied than that of heterotrophic microorganisms such as soil fungi and bacteria. This has sometimes given rise to the fallacious impression that they are unimportant soil microorganisms [17]. Terrestrial algae in fact play a major role in soil genesis and soil conservation. They constitute the initial successional stage on substrata which are poor in plant nutrients such as recent volcanic deposits, sand, and barren infertile soils denuded of macrovegetation. The interwoven algal growth consolidates the surface, leading to the formation of a soil crust which improves infiltration, may limit sheet erosion and affords a substratum upon which seeds of higher plants germinate. Algae produce a surface humus after death and dissolve certain soil minerals maintaining a reserve supply of elements in a semi-available form for higher plants [61]. Experiments on algal crusts from the Botanical Garden at the University of India indicated that these crusts (i) do not slow the rate of water infiltration [7], (ii) have a very efficient protective effect on erosion by buffeting rain, (iii) increased the moisture content of the soil underneath the algal stratum 10–15% [63]. The resistance of algal crusts to erosion is apparently the result of binding the soil surface particles into a non-erosible layer, which is also effective in breaking the force of falling water. The favourable effect of BGA on aggregation of the soil was demonstrated by Roychoudhury *et al.*, [60] who observed a 50–70% increase in the water-stable aggregates after algal inoculation. This was attributed to the action of polysaccharides released by the algae and the pressure of filamentous BGA growing in the soil. In sugarcane fields in India, a profuse growth of *Porphyrosiphon notarisii* was reported to check the erosion of the soil and to help the crop stand erect on the intact ridges during the monsoon period [63]. The same author in a study of grasslands subjected to close and heavy grazing by sheep and cattle concluded that in such deteriorated grassland, where soils are light and sandy, a thick growth of BGA during the monsoon

is perhaps the only check for erosion. *Porphyrosiphon notarisii* and *Microcoleus chthonoplastes* [8] were reported to be the most efficient colonizing strains growing as a mat that provides a suitable substratum for the germination of grass seeds. Booth [7] conducted a detailed investigation on the importance of BGA in badly eroded soils in south-central US. He concluded that (i) soil losses from plots with an algal stratum were greatly reduced as compared with the losses from bare areas, (ii) BGA constituted an initial stage in plant succession, and (iii) BGA colonization lasted for several years until higher perennial plants were able to form abundant ground cover. Singh [63] described the characteristic and common algal cover caused by *Aphanothece pallida*. Although the fresh thalli were extremely mucilaginous and fragile, they formed a compact dark grey stratum firmly adherent to the soil when dried; 0.3 g of algae were sufficient to bind 34.6 g of soil.

The floristic composition of algal crusts developing on cultivated sandy soils in Senegal was recently studied by Reynaud. The L.P.P. (*Lyngbia*, *Phormidium*, *Plectonema*) group [50] was dominant in most of the soils studied. A similar observation was made by Marathe and Anantany [31] in Indian arid soils. The dominance of the L.P.P. group in arid soils can be related to its ability to develop in drastic conditions a gel-like protoplasm and a thick mucilaginous sheath able to readily adsorb water and retain it. Among  $N_2$ -fixing genera, *Nostoc* was the most frequently observed, confirming the observation that it is one of the most consistently present  $N_2$ -fixing forms in arid soils [61]. Examples of the strains isolated from algal crusts on sandy Senegalese soils are given in Figs. 2 and 3.

## 7. Agronomical use of BGA

Most information on the effects of agronomical practices on BGA is related to paddies. Land preparation and management seem to have only incidental effects [58] and were reported not to interfere with the establishment and activity of inoculated BGA [2]. Pesticides depending on their nature and their concentration, could have inhibitory, selective or stimulatory effects on BGA. Experiments mainly with flask cultures suggest that BGA are generally more resistant to pesticides than other algae and are capable of tolerating pesticide levels recommended for field application [77]. Insecticides are generally less toxic to BGA than other pesticides [42] and have a secondary beneficial effect by controlling the grazers.

Organic manure may favour or depress BGA growth depending on its nature and mode of application. Incorporation of plant residues has frequently been reported to temporarily depress the algal populations and to have a negative effect on inoculated BGA (refer to Roger and Kulasooriya [58]). On the contrary the surface application of straw very significantly enhanced BGA growth and ARA [33].

The nature and the quantity of inorganic fertilizers as well as application techniques have a considerable influence on the algal flora. The surface application of NPK generally results in a profuse green algae growth. To prevent such a growth the incorporation of fertilizers into the soil is recommended. The deep placement




Fig. 2. BGA isolated from algal crusts in Senegal, in a medium without N. (a) *Nostoc* sp.; (b, c, h) *Anabaena* spp.; (d) *Dermocarpa*; (i, e) *Calothrix* sp.; (f) *Hapalosiphon* sp.; (g) *Gloeocapsa* sp.; (j) *Anabaenopsis* sp. (the scale represents 50  $\mu$ m).

of urea supergranules prevents the dense growth of green algae that occurs when urea is surface broadcast; moreover it does not inhibit the growth and  $N_2$  fixation in BGA [59]. Since the growth of  $N_2$ -fixing BGA in paddy fields is generally limited by P deficiency, P application alone or together with lime should be recommended (refer to Roger and Kulasoorya [58]).

In N-deficient conditions,  $N_2$ -fixing BGA are not hindered by competition from the other algae and they can develop profusely if the other environmental factors are not limiting. When nitrogenous fertilizers are applied their ARA is inhibited or




Fig. 3. BGA isolated from algal crusts in Senegal, on a medium with N. (a) *Pseudanabaena* sp.; (h, d) *Oscillatoria* spp.; (b, c, e, f, g, i, j) different species belonging to the LPP (*Lyngbia*, *Phormidium*, *Plectonema*) group (the scale represents 50  $\mu\text{m}$ ).

at least affected. Moreover, N fertilizers appear to enhance the growth of non- $\text{N}_2$ -fixing algae thus increasing the competitive pressure against  $\text{N}_2$ -fixing forms [59]. However, this effect seems to be transient as suggested by pot experiments [49] and preliminary field data. Simultaneously the effect of N fertilizers on ARA is only partial. Venkataraman [79] reported that ARA was not depressed in a soil-rice-BGA system with less than 40 ppm ammonium-N in stagnant paddy water. Within mixed algal masses combined N may diffuse at a slower rate than the readily

available dissolved  $N_2$ , and a local depletion in combined N may favour  $N_2$ -fixing algae. Thus in the fields, the lack of competition with  $N_2$ -fixing algae in the presence of mineral nitrogen may not be as clear-cut as was first thought.

Molybdenum, because of its function in nitrogenase, is required by  $N_2$ -fixing BGA. Subrahmanyam *et al.* [68], suggested the addition of sodium molybdate ( $0.25 \text{ kg ha}^{-1}$ ) to the soil to improve  $N_2$ -fixing algal growth. This addition has been shown to be beneficial in some situations.

Other nutrients (Fe, Mg, K, etc.) are required for optimal growth of BGA but their ecological implications as limiting factors or as factors affecting the composition of the algal community in paddies have not been documented.

### 7.1. Reclamation of Usar lands [62]

Usar soils are saline (solonchak) and alkaline (solonetz) unproductive soils, found extensively throughout India. They are characterized by impermeability, extreme hardness and the occasional presence of undesirable salts on the surface, all of which adversely affect the plant growth. The pH value is usually high throughout the profile. The subsoil water table is generally found between 3.0 and 4.5 meters below the surface. These soils are usually very deep. The method of reclamation arose from the observation [62] that in the alkaline Usar soils of northern India, while other plants fail to grow, blue-green algae form a thick stratum on the soil surface during the rainy season (July–September) and the retreating monsoon (December–January). These soils, characterized by an alkalization process, can be reclaimed by replacing sodium with calcium through the addition of chemical correctives such as gypsum, a rather expensive method. Another possibility is to help the reaction of calcium carbonate with the sodium clay by waterlogging the soil and by adding organic matter and N. The abundant growth of  $N_2$ -fixing BGA on waterlogged soils fulfills these requirements.

The following reclamation process was tested. During May and June, before the rains started, the land was divided into plots of less than 0.4 ha. The plots were enclosed by an earth embankment. After the first showers,  $N_2$ -fixing BGA formed a thick and compact stratum. Later, when the soil was waterlogged,  $N_2$ -fixing forms characteristic of rice fields developed and continued their active growth as long as the soil was waterlogged. In field trials, after a year of reclamation, a transplanted paddy crop produced a yield of  $1,600\text{--}2,200 \text{ kg grain ha}^{-1}$  and allowed the successful growth of sugarcane after 3 years. Laboratory experiments over a 3-year period indicated tremendous changes in the soil characteristics after algal growth compared with the control covered with a black cloth: pH decreased from 9.2 to 7.5. Large increase in organic matter content (69%), total N (46%), water-holding capacity (35%), and exchangeable calcium (31%), were observed. The most significant changes concerned the different forms of P.  $CO_2$ -soluble P was 0.45 in the control and 8.15 ppm in the treated plot with BGA. The role of BGA in

converting the sodium clay into a type of calcium clay was not explained but a possible effect of oxalic acid excreted by the algae was suggested.

Subsequent improvements of the reclamation technique were proposed: doing the process several times in a year where irrigation water is available, contour farming (field operations such as plowing, etc. are done at right angles to the slope of the land) to retard water flow and increase water penetration; and introducing *Nostoc commune* where it does not occur naturally.

## 7.2. Algalization

BGA were among the first  $N_2$ -fixing agents recognized to be active in flooded rice soils. Since De [14] attributed the natural fertility of the tropical paddy fields to these organisms, many trials have been conducted to increase rice yield by inoculating the soil with BGA. This practice, also called algalization, a terminology introduced by Venkataraman [74], has been reported to have a beneficial effect on grain yield in different agroclimatic conditions. However, some reports also indicate failure of algalization. The conclusions of the review of Roger and Kulasooriya [58] on algalization are summarized thereafter.

**7.2.1. Methodology.** Most experiments on algalization were on a 'black box' basis, where only the last indirect effect (grain yield) of algalization was observed and the intermediate effects were not studied. There is little information on the qualitative and quantitative variations of the  $N_2$ -fixing algal flora and the N balance in inoculated paddy soils. Pot and field experiments have been conducted, usually on a single crop. The relative increase in grain yield over the control was an average 28% in pot experiments and 15% in field experiments. The better growth of BGA in pot experiments is probably attributable to the reduction of climatic disturbance and to the mechanical effect of the pot walls, where BGA frequently seem to grow preferentially and profusely. Pot experiments may therefore only be suitable for qualitative studies, since they overestimate the effects of BGA inoculation. Most of the field experiments on the other hand, were conducted in only one growing season and may underestimate the effects of algalization since the advantages of a slow N release from dead BGA may not be apparent in the first algalized crop.

**7.2.2. Effects of algalization on rice.** Algalization may affect plant size, its nitrogen content, and the number of tillers, ears, spikelets, and filled grains per panicle. The most frequently used criterion for assessing the effects of algalization has been better grain yield. Results of field experiments conducted mainly in India report an average yield increase of about 14% over the control, corresponding to about 450 kg grain  $ha^{-1}$  per crop where algal inoculation was effective. A higher grain yield increase was observed when algalization was in combination with lime, P and sometimes molybdenum application. Unfortunately it is not possible to separate the direct effect of PK fertilizers on rice from its indirect effect upon the growth of

indigenous or introduced algae. The effects of algalization used with N fertilizers are controversial. Since biological N<sub>2</sub> fixation is known to be inhibited by inorganic N the beneficial effect of algalization in the presence of N fertilizers was most frequently interpreted as resulting from growth-promoting substances produced by algae or also by a temporarily immobilization of added N followed by a slow release through subsequent algal decomposition permitting a more efficient utilization of N by the crop (see Section 5.2.).

*7.2.3. Effects of algalization on soil properties and microflora.* Grain-yield measurements suggest that algalization produces both a cumulative and residual effect. This was attributed to a build up of both the organic N content and the number of BGA propagules in the soil, facilitating the reestablishment of the BGA biomass. Several reports indicate an increase in organic matter and organic N; algalization was also reported to increase: aggregation status of the soil [61], water-holding capacity [63], available P, total microflora, *Azotobacter*, *Clostridia*, and nitrofiere [21].

*7.2.4. Limiting factors for algalization.* Among the limiting factors responsible for the failure of algalization only pH and available P content of the soil have been studied. Low pH is an important limiting factor for algalization as demonstrated by unsuccessful trials in acidic soils of India, Sri Lanka and Japan. In Sri Lanka and Japan algalization was effective only when the soil was supplemented with calcium carbonate, the effect of inoculation being related to the amount of lime added. Since in some soils, algalization is inefficient in spite of the addition of lime and phosphate [36], available P content should probably not be the only factor limiting the effect of algalization; texture, organic matter content, CEC of saturated extracts, total N are probably not important limiting factors [36, 68]. Among the biotic factors that can possibly limit BGA growth inoculum, grazing by the zooplankton has been already mentioned. Other possible mechanisms involved such as antagonism, competition, etc. have been cited, but their role is not clear. Low temperatures, heavy rains, and cloudy weather have also been reported to limit the establishment of the inoculum.

#### *7.2.5. Algalization technology*

*7.2.5.1. Inoculum production and conservation.* The methodology of BGA production has been reviewed by Watanabe and Yamamoto [83] and Venkataraman [76, 77]. Methods of inoculum production in artificially controlled conditions have been developed mainly in Japan. Different types of tank cultures and outdoor closed circulating systems have been described [82]. Algae are grown either in liquid culture, mixed with an inert material (support) and dried, or grown directly on the support and dried. Various supports such as sand, pumice stone, volcanic earth, and blocks of synthetic sponge have been tested to conserve and facilitate transportation of the algae. Such inocula reportedly maintained their capacity for growth unimpaired for at least two years.

Producing the inoculum in artificially controlled conditions is well defined but

relatively expensive. On the contrary, the open air soil culture, used in India, is more simple, less expensive and easily adoptable by the farmers. It is based on the use of a starter culture that is a multistrain inoculum of *Aulosira*, *Tolypothrix*, *Scytonema*, *Nostoc*, *Anabaena* and *Plectonema*, provided by the 'All India Coordinated Project on Algae' [2]. This inoculum is multiplied by the farmer in shallow trays or tanks with 5–15 cm water, about 4 kg soil m<sup>-2</sup>, 100 g triple superphosphate m<sup>-2</sup> and insecticide. If necessary, lime is added to correct the soil pH to about 7.0–7.5. In 1 to 3 weeks, a thick mat develops on the soil surface and sometimes floats. Watering is stopped and water in the trays is allowed to dry up in the sun. Algal flakes are then scraped off and stored in bags for use in the fields. With such a method, the ultimate proportion of individual strains in the algal flakes is unpredictable. It is assumed that, because the inoculum is produced in soil and climatic conditions similar to those in the field, dominant strains will be the most adapted to the local conditions. The recorded rates of production of algal flakes in the open air soil culture range from 0.4 to 1.0 kg m<sup>-2</sup> in 15 days, indicating that a 2m<sup>2</sup> tray can produce in 2–3 months enough algal material to inoculate 1 ha of rice field.

*7.2.5.2. Inoculation of the rice field.* The methods of field application have been reviewed by Venkataraman [77]. For transplanted rice the algal inoculum is generally applied 1 week after the transplanting. When rice is sown, seeds can be coated by mixing the algal suspension and 2–3 kg calcium carbonate per 10–20 kg seed and air-dried in the shade.

Recommendations for field application of dried algal inoculum (algal flakes) given by the 'All India Coordinated Project on Algae' [2] indicate that 8–10 kg of dry algal flakes applied 1 week after transplanting is sufficient to inoculate 1 ha; a large amount will accelerate multiplication and establishment in the field. Algalization can be used with high levels of commercial nitrogen fertilizer, but reduction of the N dose by one third is recommended. To benefit from the cumulative effect of algalization the algae should be applied for at least three consecutive seasons. Recommended pest-control measures and other management practices do not interfere with the establishment and activity of these algae in the fields.

*7.2.5.3. Pay-off of algal technology.* Trials in Indian station (in 1978), showed that adding 10 kg of algal culture ha<sup>-1</sup> which costs about 4 US \$, increases paddy yield; the average yield increase is worth 60 to 90 US \$ [43]. The pay-off of algal technology based on the results of a large number of field experiments was discussed by the 'All India Coordinated Project on Algae' [2] and it was inferred that if algal technology was introduced into even 50% of the Indian rice area, it would supposedly result in a saving of about  $3.8 \times 10^5$  t of N. In a large number of trials where the recommended level of N fertilizer (100 kg ha<sup>-1</sup>) was complemented with algal application (10 kg ha<sup>-1</sup>) an average increase of about 300 kg rice grains ha<sup>-1</sup> was reported.

## References

1. Aiyer, R.S. 1965 Comparative algological studies in rice fields in Kerala state. *Agric. Res. J. Kerala* 3, 100–104.
2. All India Coordinated Project on Algae. 1979 Algal biofertilizers for rice. India Agricultural Research Institute, New Delhi, 61 pp.
3. Araragi, M., Phetchawee, S. and Tantitanapat, P. 1979 Microflora related to the nitrogen cycle in the tropical upland farm soils. *Soil Sci. Plant Nutr.* 25, 235–244.
4. Arora, S.K. 1969 The role of algae on the availability of phosphorus in paddy fields. *Riso* 18, 135–138.
5. Barbey, C. and Coute, A. 1976 Croûtes à Cyanophycées sur les dunes du Sahel mauritanien. *Bull. IFAN* 38A, 732–736.
6. Bisiach, M. 1972 Laboratory algicidal screening for the control of algae in rice. *Riso* 21, 43–58.
7. Booth, W.E. 1941 Algae as pioneers in plant succession and their importance in erosion control. *Ecology* 22, 38–46.
8. Brock, T.D. 1975 Effect of water potential on a *Microcoleus* (Cyanophyceae) from a desert crust. *J. Phycol.* 11, 316–320.
9. Chapman, R.L., Bayer, D.E. and Lang, N.J. 1972 Observations on the dominant algae in experimental California rice fields. *J. Phycol. Suppl.* 8, 17.
10. Chapman, V.J. and Chapman, D.J. 1973 Soil algae and Symbiosis. In: *The Algae*, Ch. 17, pp. 381–387. Macmillan, London.
11. Dadhich, K.S., Varma, A.K. and Venkataraman, G.S. 1969 Effect of *Calothrix* inoculation on vegetable crops. *Plant and Soil* 31, 377.
12. Das, S.S. 1976 Algal weeds and their chemical control – a review. *Indian J. Plant Prot.* 4, 201–208.
13. David, K.A.V. and Fay, P. 1977 Effects of long-term treatment with acetylene on nitrogen fixing micro-organisms. *Appl. Environ. Microbiol.* 34, 640–646.
14. De, P.K. 1939 The role of blue-green algae in nitrogen fixation in rice fields. *Proc. R. Soc. Lond.* 127 B, 121–139.
15. Durrel, L.W. 1964 Algae in tropical soils. *Trans. Amer. Micros. Soc.* 83, 79–85.
16. Dutta, N. and Venkataraman, G.S. 1958 An exploratory study of the algae of some cultivated and uncultivated soils. *Indian J. Agron.* 3, 109–115.
17. Fogg, G.E., Stewart, W.D.P., Fay, P. and Walsby, A.E. 1973 *The blue-green algae*. Academic Press, London and New York, 459 pp.
18. Garcia, J-L., Raimbault, M., Jacq, V., Rinaudo, G. and Roger, P. 1973 Microbial activities in paddy fields in Senegal; influence of physico-chemical properties of soils and of the rhizosphere effect (in French, English summary) *Rev. Ecol. Biol. Sol* 11, 169–185.
19. Gupta, A.B. and Shukla, A.C. 1967 Studies on the nature of algal growth promoting substances and their influence on growth, yield, and protein content of rice plants. *Labdev. J. Sci. Technol. Kanpur* 5, 162–163.
20. Holm-Hansen, O. 1968 Ecology, physiology and biochemistry of blue-green algae. *Ann. Rev. Microbiol.* 22, 47–70.
21. Ibrahim, A.N., Kamel, M. and El-Sherbeny, M. 1971 Effect of inoculation with alga *Tolypothrix tenuis* on the yield of rice and soil nitrogen balance. *Agrokem. Talagtan* 20, 389–400.
22. Ichimura, S. 1954 Ecological studies on the plankton in paddy fields I. Seasonal fluctuations in the standing crop and productivity of plankton. *Jpn. J. Bot.* 14, 269–279.

23. Jacq, V. and Roger, P.A. 1977 Decrease of losses due to sulphate reducing processes in the spermosphere of rice by presoaking seeds in a culture of blue-green algae (in French, English summary). Cahiers ORSTOM. sér. Biol. 12, 101–108.
24. Jones, K. 1977 The effects of temperature on acetylene reduction by mats of blue-green algae in sub-tropical grassland. *New Phytol.* 78, 433–436.
25. Jurgensen, M.S. and Davey, C.B. 1968 Nitrogen-fixing blue-green algae in acid forest and nursery soils. *Can. J. Microbiol.* 14, 1179–1183.
26. Kaushik, B.D. and Venkataraman, G.S. 1979 Effect of algal inoculation on the yield and vitamin C content of two varieties of tomato. *Plant and Soil* 52, 135–137.
27. Kulasooriya, S.A., Roger, P.A., Barraquio, W.L. and Watanabe, I. 1980 Epiphytic nitrogen fixation on weeds in a rice field ecosystem. In: Proceedings of Nitrogen Cycling South-East Asian Wet Monsoonal Ecosystems. Wetselaar, R., Simpson, J.R. and Rosswall, T. (eds.), Australian Academy of Science, Canberra (in press).
28. Kulasooriya, S.A., Roger, P.A., Barraquio, W.L. and Watanabe, I. 1980 Biological nitrogen fixation by epiphytic microorganisms in rice fields. IRRI. IPRS No. 47. Feb. 1980, 10 pp.
29. Kurasawa, H. 1956 The weekly succession in the standing crop of plankton and zoobenthos in the paddy field, Part 1 and 2. *Bull. Res. Sci. Japan* 41–42, 86–98 and 45, 73–84.
30. Lowendorf, H.S. 1980 Biological nitrogen fixation in flooded rice. Agronomy paper 1305. Dpt. of Agronomy. Cornell Univ. Ithaca, N.Y.
31. Marathe, K.U. and Anantani, Y.S. 1972 Observations on the algae of some Indian arid soils. *The botanique* 3, 13–20.
32. Materasi, R. and Balloni, W. 1965 Some observations on the presence of autotrophic nitrogen-fixing microorganisms in paddy soils, (in French, English summary), *Ann. Inst. Pasteur* 109, 218–223.
33. Matsuguchi, T. and Ick-Dong Yoo. 1979 Stimulation of phototrophic N<sub>2</sub> fixation in paddy fields through rice straw applications. In: Proceedings of Nitrogen Cycling in South-East Asian Wet Monsoonal Ecosystems. Wetselaar, R., Simpson, J.R. and Rosswall, T. (eds.), Australian Academy of Science, Canberra (in press).
34. Mayland, H.F. and McIntosh, T.H. 1966 Availability of biologically fixed nitrogen-15 to higher plants. *Nature, London* 209, 420–421.
35. Mitra, A.K. 1951 The algal flora of certain Indian soils. *Indian J. Agric. Sci.* 21, 357.
36. Okuda, A. and Yamaguchi, M. 1952 Algae and atmospheric nitrogen fixation in paddy soils. II: Relation between the growth of blue-green algae and physical or chemical properties of soil and effect of soil treatments and inoculation on the nitrogen fixation. *Mem. Res. Inst. Food Sci.* 4, 1–11.
37. Okuda, A. and Yamaguchi, M. 1956 Nitrogen-fixing microorganisms in paddy soils. II: Distribution of blue-green algae in paddy soils and the relationship between the growth of them and soil properties. *Soil and Plant Food* 2, 4–7.
38. Pandey, D.C. 1965 A study of the algae from paddy soils of Ballia and Ghazipur districts of Uttar Pradesh, India. I. Cultural and ecological considerations. *Nova Hedwigia* 9, 299–334.
39. Pankratova, E.M. and Vakrushev, 1969 Utilization by higher plants of nitrogen fixed from the atmosphere by blue-green algae (in Russian, English summary). *Mikrobiologiya* 38, 1080–1084.
40. Peterson, R.B. and Burris, R.H. 1976 Conversion of acetylene reduction rates

- to nitrogen fixation rates in natural populations of blue-green algae. *Anal. Biochem.* 73, 404–410.
41. Prescott, G.W. 1968 Ecology Soil. In: *The Algae*, pp. 320–323, Steers, W.C. and Glass, H.B. (eds.), Houghton Mifflin, Boston.
  42. Raghu, K. and MacRae, I.C. 1967 The effect of the gamma isomer of benzene hexachloride upon the microflora of submerged rice soils. II. Effect upon algae. *Can. J. Microbiol.* 13, 173–180.
  43. Rao, T.R. 1978 Blue-green algae boost rice yields. *Intensive Agric.* 16, 19–20.
  44. Renaut, J., Sasson, A., Pearson, H.W. and Stewart, W.D.P. 1975 Nitrogen-fixing algae in Morocco. In: *Nitrogen fixation by free-living microorganisms*, pp. 229–246. Stewart, W.D.P. (ed.), Cambridge University Press, Cambridge.
  45. Reynaud, P.A. and Roger, P.A. 1978 N<sub>2</sub>-fixing algal biomass in Senegal rice fields. *Ecol. Bull. Stockholm* 26, 148–157.
  46. Reynaud, P.A. and Roger, P.A. 1978 Photophobotaxis and photokinesis among *Oscillatoria* sp. 77 S23 (in French, English summary). *Cah. ORSTOM, sér. Biol.*, 13, 157–164.
  47. Reynaud, P.A. and Roger, P.A. 1981 Seasonal variations of algal flora and of N<sub>2</sub>-fixing activity in a waterlogged sandy soil. *Rev. Ecol. Biol. Sol.* 18, 9–27.
  48. Reynaud, P.A. 1980 Cyanobacteria and *Azolla* nitrogen fixation: Agronomical potentialities in tropical Africa. In: *Proceedings of Recycling of Organic Matter in West Africa*. FAO/SIDA, LOME, Togo (in French, English summary).
  49. Reynaud, P.A. 1980 Fate of Cyanobacteria inoculums during the two first months of rice growth on a waterlogged soil (in French, English summary). *Cah. ORSTOM. sér. Biol.* 43, 53–60.
  50. Rippka, R., Deruelles, J., Waterbury, J.B., Herdman, M. and Stanier, R.Y. 1979 Generic assignments, Strain histories and Properties of pure Culture of Cyanobacteria. *J. Gen. Microbiol.* 111, 1–61.
  51. Rodgers, G.A., Bergman, B., Henriksson, E. and Udris, M. 1979 Utilization of blue-green algae as biofertilizers. *Plant and Soil* 52, 99–107.
  52. Roger, P. and Reynaud, P. 1976 Dynamics of the algal populations during a culture cycle in a Sahel rice field (in French, English summary). *Rev. Ecol. Biol. Sol.* 13, 545–560.
  53. Roger, P. and Reynaud, P. 1977 Algal biomass in rice fields of Senegal: relative importance of Cyanophyceae that fix nitrogen (in French, English summary). *Rev. Ecol. Biol. Sol* 14, 519–530.
  54. Roger, P.A., Reynaud, P.A., Rinaudo, G.E., Ducerf, P.E. and Traore, T.M. 1977 Log-normal distribution of acetylene-reducing activity *in situ* (in French, English summary). *Cah. ORSTOM, sér. Biol.* 12, 133–140.
  55. Roger, P.A. and Reynaud, P.A. 1978 Enumeration of the algae in submerged soil: law of the distribution of organisms and the density of sampling (in French, English summary). *Rev. Ecol. Biol. Sol* 15, 219–234.
  56. Roger, P.A. and Reynaud, P.A. 1979 Ecology of blue-green algae in paddy fields. In: *Nitrogen and rice*, pp. 289–309. International Rice Research Institute, Los Baños.
  57. Roger, P.A., Kulasoorya, S.A., Barraquio, W.L. and Watanabe, I. 1979 Epiphytic nitrogen fixation on lowland rice plants. In: *Proceedings of Nitrogen Cycling in South-East Asian Wet Monsoonal Ecosystems*. Wetselaar, R., Simpson, R.J. and Rosswall, T. (eds.), Australian Academy of Science, Canberra (in press).

58. Roger, P.A. and Kulasoorya, S.A. 1980 Blue-green algae and rice. The International Rice Research Institute, Los Baños, Philippines, 112 pp.
59. Roger, P.A., Kulasoorya, S.A., Tirol, A.C. and Craswell, E.T. 1980 Deep placement: a method of nitrogen fertilizer application compatible with algal nitrogen fixation in wetland rice soils. *Plant and Soil* 57, 137–142.
60. Roychoudhury, P., Kaushik, B.D., Krishnamurthy, G.S.R. and Venkataraman, G.S. 1979 Effect of blue-green algae and *Azolla* application on the aggregation status of the soil. *Current Science* 48, 454–455.
61. Shield, L.M. and Durrell, L.W. 1964 Algae in relation to soil fertility. *Bot. Rev.* 30, 93–128.
62. Singh, R.N. 1950 Reclamation of Usar lands in India through blue-green algae. *Nature*, London. 165, 325–326.
63. Singh, R.N. 1961 Role of blue-green algae in nitrogen economy of Indian agriculture. Indian Council of Agricultural Research, New Delhi, 175 pp.
64. Stewart, W.D.P. 1967 Transfer of biologically fixed nitrogen in a sand-duneslack region. *Nature*, London 214, 603–604.
65. Stewart, W.D.P. 1978 Nitrogen fixing Cyanobacteria and their association with eukaryotic plants. *Endeavour* 2, 170–179.
66. Stewart, W.D.P., Sampaio, M.J., Isichei, A.O. and Sylvester-Bradley, R. 1978 Nitrogen fixation by soil algae of temperate and tropical soils. In: Limitation and potentials for biological nitrogen fixation in the tropics, pp. 41–63. Döbereiner, J. *et al.* (eds.), Plenum Press, New York and London.
67. Stewart, W.D.P., Rowell, P., Ladha, J.K. and Sampaio, M.J.A. 1979 Blue-green algae (Cyanobacteria) – some aspects related to their role as sources of fixed nitrogen in paddy soils. In: Nitrogen and Rice, pp. 263–285. International Rice Research Institute, Los Baños, Philippines.
68. Subrahmanyam, R., Manna, G.B. and Patnaik, S. 1965 Preliminary observations on the interaction of different rice soil types to inoculation of blue-green algae in relation to rice culture. *Proc. Indian Acad. Sci. B* 62, 171–175.
69. Suzuki, T. and Kawai, K. 1971 Soil of the Sambor. Cambodia. *Bull. Natl. Agric. Sci. B* 22, 211–304.
70. Taha, E.E.M. and Al Refai, A.E. 1962 Physiological and biochemical studies on the nitrogen-fixing blue-green algae: 1. On the nature of cellular and extracellular substances formed by *Nostoc commune*. *Arch. Mikrobiol.* 41, 307–312.
71. Tchan, Y.T. and Beadle, N.C. 1955 Nitrogen economy in semi arid plant communities II. The non-symbiotic nitrogen-fixing organisms. *Proc. Linn. Soc. N.S.W.* 80, 97–104.
72. Traore, R.M., Roger, P.A., Reynaud, P.A. and Sasson, A. 1978 N<sub>2</sub>-fixation by blue-green algae in a paddy field in Mali (in French, English summary). *Cah. ORSTOM sér. Biol.* 13, 181–185.
73. Venkataraman, G.S. 1961 The role of blue-green algae in agriculture. *Sci. Cult.* 27, 9–13.
74. Venkataraman, G.S. 1966 Algalization. *Phytos.* 5, 164–174.
75. Venkataraman, G.S. and Neelakanthan, S. 1967 Effect of the cellular constituents of the nitrogen fixing blue-green alga *Cylindrospermum musicola* on the root growth of the rice seedlings. *J. Gen. Appl. Microbiol.* 13, 53–61.
76. Venkataraman, G.S. 1969 The cultivation of algae. Indian Council of Agricultural Research, New Delhi, 319 pp.
77. Venkataraman, G.S. 1972 Algal biofertilizers and rice cultivation. Today and Tomorrow's Printers, Faridabad (Haryana), 75 pp.
78. Venkataraman, G.S. 1975 The role of blue-green algae in tropical rice culti-

- vation. In: Nitrogen fixation by free-living microorganisms, pp. 207–218. Stewart, W.D.P. (ed.), Cambridge Univ. Press.
79. Venkataraman, G.S. 1979 Algal inoculation of rice fields. In: Nitrogen and rice, pp. 311–321. International Rice Research Institute, Los Baños.
  80. Watanabe, A. 1956 On the effect of the atmospheric nitrogen-fixing blue-green algae on the yield of rice (in Japanese). Bot. Mag. 69 (820/821), 530–535.
  81. Watanabe, A. 1959 Distribution of nitrogen-fixing blue-green algae in various areas of south and east Asia. J. Gen. Appl. Microbiol. 5, 21–29.
  82. Watanabe, A. 1959 On the mass culturing of a nitrogen fixing blue-green algae *Tolypothrix tenuis*. J. Gen. Appl. Microbiol. 5, 85–91.
  83. Watanabe, A. and Yamamoto, Y. 1970 Mass culturing preservation and transportation of the nitrogen fixing blue-green algae. In: Proc. 2nd Symposium on Nitrogen Fixation and Nitrogen Cycle Sendai, Japan, pp. 22–28.
  84. Watanabe, A. and Yamamoto, Y. 1971 Algal nitrogen fixation in the tropics. Plant and Soil (special volume), 403–413.
  85. Watanabe, I., Lee, K.K., Alimagno, B.V., Sato, M., Del Rosario, D.C. and De Guzman, M.R. 1977 Biological N<sub>2</sub>-fixation in paddy field studied by *in situ* acetylene-reduction assays. IRRI Res. Pap. Ser. 3, 1–16.

DEVELOPMENTS IN PLANT AND SOIL SCIENCES  
VOLUME 5

**MICROBIOLOGY  
OF TROPICAL SOILS  
AND PLANT PRODUCTIVITY**

Y.R. DOMMERGUES/H.G. DIEM  
(EDITORS)

MARTINUS NIJHOFF / DR W. JUNK PUBLISHERS

*Distributors:*

*for the United States and Canada*

Kluwer Boston, Inc.  
190 Old Derby Street  
Hingham, MA 02043  
USA

*for all other countries*

Kluwer Academic Publishers Group  
Distribution Center  
P.O. Box 322  
3300 AH Dordrecht  
The Netherlands

Library of Congress Cataloging in Publication Data

Main entry under title:

Microbiology of tropical soils and plant  
productivity.

(Developments in plant and soil sciences ;  
v. 5)

Includes index.

1. Soil microbiology--Tropics. 2. Micro-organisms, Nitrogen-fixing--Tropics. 3. Crops and soils--Tropics. 4. Soils--Tropics. 5. Optimal soil fertility--Tropics. I. Dommergues, Y. R. (Yvon R.) II. Diem, H. G. III. Series.

QR111.M4 631.4'6 81-22536

AACR2

ISBN 90-247-2624-7

(this volume)

ISBN 90-247-2405-8

(series)

*Copyright © 1982 by Martinus Nijhoff/Dr W. Junk Publishers, The Hague.*

*All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, mechanical, photocopying, recording, or otherwise, without the prior written permission of the publishers, Martinus Nijhoff/Dr W. Junk Publishers, P.O. Box 566, 2501 CN The Hague, The Netherlands.*

PRINTED IN THE NETHERLANDS