

1

Placer mining operations and modifications of the physical chemical nature of the waters of the Rio Kaka drainage basin (Andes, Bolivia)

Jean Louis Guyot & Gerard Herail
Orstom, C.P. 8714, La Paz, Bolivia

ABSTRACT Exploitation of gold placers in the Bolivian Andes leads to a modification of the rivers physico-chemistry. The impact of the mining activity is evaluated for the River Tipuani where extraction is intense in low water, compared to the River Challana which is lightly exploited.

INTRODUCTION

In Bolivia, the gold placers of the Royal Cordillera, located in the North of the country (Fig. 1), provides several tons of gold a year. The most important exploitations are located along the Rivers Challana, Tipuani and Consata-Mapiri, that join the River Coroico to form the River Kaka. Gold is produced from the treatment of alluvial deposits contained in Miocene paleovalleys, of alluvial deposits from Quaternary terraces, but mainly of alluvial deposits of present riverbeds or low terraces. Mining activity is particularly important during the low water season, between April-May and November-December. Indeed, the deposits contained in the bottom of the valley are inaccessible during the high water season.

The purpose of this paper is the evaluation of the influence of placer exploitation on water quality. Three rivers have then been studied: the River Challana, the River Tipuani and the River Consata-Mapiri. Along the River Challana, gold exploitation is yet little developed. This river is then used as a reference, because it intersects the same geological, geomorphological and biogeographical structures as the River Tipuani, where mining exploitation is very important. Along the River Consata-Mapiri, the extraction is located on terraces and effluents are not directly poured to the river. The part of the suspensions due to mining activity in the solid discharge of the River Kaka is discussed.

HYDROCHEMICAL CONTEXT

Geology

The rivers of the River Kaka drainage basin, where sandstone and pyritic slates from Ordovician build up the largest outcrops (Martinez, 1980), granites, granodiorites and orthogneiss that outcrop along the crest line (Zongo-Yani block, Illampu block), as well as the contact metamorphic aureole, that line them to the North-East (Bard & al., 1974) are only drained by the most upstream tributaries. These rocky set-ups are intersected by narrow and sloping valleys covered by vegetation below 3000 m. Below 1500 m topographical features are softer and the Ordovician

ORSTOM Fonds Documentaire 115

10 AVR. 1990


N° : 27.883 ex 1

Cote : B

M

P112

substratum is partially covered by Tertiary Cangalli conglomerates (Fornari & al., 1987) and Quaternary alluviums. Out of these mining areas, these deposits are entirely covered by forest, and form a buffer between pyritic slates and run-off waters. Along the River Consata-Mapiri, the same geological formations outcrop, but the vegetal coverage is very discontinuous on the slopes, and numerous landslides affect ordovician schists and alterites that cover them. Upstream of the Consata and Camata drainage basin, volcanic series and limestones outcrop, but on limited areas.



Mostly formed by the Rivers Mapiri, Tipuani, Challana and Coroico, the River Kaka shows a $1220 \text{ m}^3 \cdot \text{s}^{-1}$ mean discharge for the 1968-1983 period at the station of Nube. This mean discharge represents 52% of the River Alto-Beni discharge at the footsteps of the Andes (Angosto del Bala), whereas the area of the River Kaka drainage basin represents only 30% of the total area of the Alto-Beni basin (Espinoza, 1985). The hydrological regime of the River Kaka (Fig. 2) is characterized by a well defined high water period from December to April, and a low water period from May to November. The five months of the rainy season are responsible for 75% of the annual mean discharge, and the three rainiest months only, from January to March, provide more than half (55%) of the mean annual discharge. The same regime, of torrential type, is observed on the whole of the basin rivers, such as the River

The flow of total suspended sediments (TSS) has been measured by the SENAMHI (Bolivian National Climatological and Hydrological Service) on the River Coroico (1976-1977) and the River Mapiri (1975-1979). The regime of this sediment discharge can be compared to the hydrological regime, but the influence of the rainy season is much more important (Fig. 4). Indeed, most of the sediment discharge occurs during the rainy season from January to March, which provide up to 95% of the annual global volume (Guyot & al., 1988).

Table 1 Characteristics of the sampling water points (July 1987)

Num.	River	Place	Area (km ²)	Discharge (m ³ .s ⁻¹)	Temp. (°C)	pH	TDS (mg.l ⁻¹)	TSS
1	Challana	Upstream	1850	18.3	21.0	6.2	22.9	0.2
2	Sapukani		24	<0.2	22.7	6.7	20.7	0.4
3	Challana	Guanay	2070	24.9	21.7	6.3	26.0	6.2
4	Tipuani	Cfl.Tora	911	12.4	18.4	6.0	30.3	6.2
5	Tora	Cfl.Tipuani	109	3.2	19.4	6.6	27.0	0.3
6	Tipuani	Mine			18.0	6.1	30.6	8.5
7	Bartolo		17	<0.2	18.7	5.4	18.4	0.2
8	Tipuani	Tipuani			21.7	6.1	30.6	466
9	Tipuani	Tipuani	1350	17.4	20.7	6.2	26.2	826
10	S.Cristobal	Sorata	264		15.0	7.8	117	6.3
11	Gunduraya			<0.2	17.0	4.9	69.8	0.5
12	Consata	Consata			18.0	7.8	388	813
13	Consata	Cfl.Camata	3590	17.9	23.0	7.4	508	439
14	Camata	Cfl.Consata	2530	29.1	23.7	6.4	50.4	9.6
15	Tarapo			<0.2	22.7	6.1	26.8	5.3
16	Mapiri	Mapiri			22.9	7.8	131	100
17	Korijahuira		245	2.0	20.1	4.7	38.6	4.7
18	Merkhe		595	7.1	22.3	4.1	27.1	0.1
19	Chimate		107	3.7	19.8	5.2	28.1	1.4
20	Dinamarca		52	0.6	21.0	5.6	21.6	0.6
21	Mariapo		208	1.9	22.6	6.6	20.0	0.4
22	Charoplaya		34	<0.2	22.3	7.3	30.7	0.1
23	Mapiri	Guanay	11600	82.3	23.0	6.9	80.6	41.5
24	Mapiri	Cfl.Coroico	15000	125	22.7	6.9	56.9	382
25	Coroico	Cfl.Mapiri	5640	83.1	22.5	7.0	30.3	6.6
26	AltoBeni	Angosto Bala	67200	501	26.0	8.0	90.6	119




Fig. 5 TDS and TSS evolutions along the Rivers Tipuani and

Challana in low water season (July 1987).

discharge, 4.4% of the TDS, but 77.8% of the TSS. The River Challana, then represents 12.0% of the discharge, 0.8% of the TSS and 6.3% of the TDS of the River Kaka (tabl. 2). This very strong influence of mining exploitation in low water upon the sediment discharge is still noticeable at the footsteps of the Andes, at Angosto del Bala, where the River Tipuani alone is responsible for 24.1% of the sediment discharge, admitting that there is no deposit neither retake of sediments between these two stations. The influence of gold mining upon the suspended load of the River Mapiri is difficult to quantify because of the high natural concentrations that are observed in this river.

Table 2 Exports by the River Kaka (July 1987)

River	Place	Discharge ($m^3 \cdot s^{-1}$)	% Ang. Bala		Dissoolv. discharge ($t \cdot d^{-1}$)	% Ang. Bala		Suspend. discharge ($t \cdot d^{-1}$)	% Ang. Bala
			TDS ($mg \cdot L^{-1}$)	TSS ($mg \cdot L^{-1}$)		TDS ($mg \cdot L^{-1}$)	TSS ($mg \cdot L^{-1}$)		
Mapiri	Guanay	82.3	16.4	80.6	573	14.6	41.5	295	5.7
Tipuani	Guanay	17.4	3.5	26.2	39	1.0	826	1240	24.1
Challana	Guanay	24.9	5.0	26.0	56	1.4	6.2	13	0.3
Coroico	Cf. Mapiri	83.1	16.6	30.3	218	5.6	6.6	47	0.9
Kaka		208	41.5	—	886	22.6	—	1595	31.0
Alto-Beni	Ang. Bala	501	100.0	90.6	3920	100.0	119	5150	100.0

From physico-chemical data of the DUCAR (Climatological and

maximum TSS concentration in low water and a minimum in high water, presents an artificial sediment regime. The low contents that are observed in high water, although higher than the ones measured in the River Challana, correspond to a slower mining activity during the rainy season. On the contrary, in low water periods, when the exploitation in the river bed is intense, the maximum loads are measured. The gold exploitation influence on the TSS balance of the River Kaka drainage basin is then maximum in low water period, when the natural TSS are the lowest.

CONCLUSION

The exploitation of gold bearing deposits in the River Tipuani alluvium yields strong modifications of this andean river water quality. The importance of the contamination has been evaluated by the physico-chemical behavior of the River Tipuani and the River Challana, where gold extraction is very low, during the hydrological cycle. So, in low water periods, when mining activity