

Water and water activity in the solid state fermentation of cassava starch by *Aspergillus niger*

Eric Oriol, Maurice Rimbault, Sevastianos Roussos, and Gustavo Viniegra-Gonzales

Universidad Autonoma Metropolitana, Departamento de Biotecnologia, Apartado postal 55-535, 09340 Mexico D.F. Mexico

Summary. During the solid state fermentation (SSF) of cassava starch by *Aspergillus niger* estimations were made of total water, consumed water and the residual water remaining in small quantities after 23 h. A theoretical calculation based on the Ross equation showed that the water activity (a_w) of the substrate decreased to 0.85 towards the end of the culture. Such low values were assumed to be inhibitory to growth. The a_w of the substrate was increased when sugarcane bagasse was used as a high water retention capacity support. Higher growth rates and substrate conversion to biomass were obtained with this system, confirming that water availability is a critical factor in the SSF of starch substrates.

The importance of water in SSF has been widely studied under quantitative aspects (Nishio et al. 1979; Rimbault and Alazard 1980; Hoe-Kim et al. 1985) but very few reports deal with the concept of a_w to explain growth of molds on solid substrates (Narahara et al. 1982).

Water content in SSF ranges from 30% to almost 80%, depending on the material, but the retention capacity of starchy substrates (rice, cassava) is fairly poor and generally does not exceed 1.0 to 1.2 g water/g of solids (50% to 55% moisture content). In the production of proteins from solid cassava flour by fungal culture (Rimbault and Alazard 1980), growth of *A. niger* stopped, leaving 30% of starch in the medium. We investigated water availability as the possible limiting factor for growth under SSF conditions.

Introduction

One of the differences between solid-state and submerged cultures is that in the former the mois-

Material and methods

Microorganism. *Aspergillus niger* var. *hennebergi* (N° 10) de-

bled twice in water and the air flow was set to 4 l/h; two columns were removed at each sampling time. Composition of the solid substrate was: Cassava meal, 100 g; KH_2PO_4 , 5 g; $(\text{NH}_4)_2\text{SO}_4$, 9.8 g; Urea, 2.4 g. Ph of the salt solution with spores was adjusted to 2.7 which resulted in a value of 4.5 for the solid substrate after inoculation.

Analysis. The exit air of one column reactor was dried by passing through a condensation and a Silicagel column. The oxygen content was then determined by pumping it through a paramagnetic Servomex Oxygen Analyzer (Taylor Instruments). For sample analysis, 4 g of sample were homogenized in 96 ml of water with an Ultra-Turrax. After dilution, the residual starch was hydrolysed with HCl 1 N for 3 h and the resulting reducing sugars were measured by the Miller (1959) method. Protein content of samples of the cassava cultivation was assayed by the method of Lowry et al. (1951). Protein content of the bagasse cultivation was estimated by hydrolysing with HCl 6 N for 20 mn at 120°C; this was neutralized and the aminoacids of the supernatant were determined by colorimetric reaction with ninhydrin (Snell 1956) to avoid interference from lignin. Protein was related to biomass assuming that 40% of the biomass is constituted by protein.

Determination of water activity. Water activity was determined with a water activity detector Humidat IC II purchased from Novasina. Prior to the measurements, 3 to 4 g of sample were cut if necessary and exposed for 10 min to an U.V. radiation of 254 nm wavelength and 300 W intensity to avoid any evolution of the sample during the measurement. The relative humidity captor was placed in a temperature regulation chamber at 30°C and the state of equilibrium was reached after 3 h.

Results and discussion

Water activity (a_w) of the substrate

The two components of the initial substrate medium were the cassava flour and the mixture of salts. We thus determined the sorption isotherms




Fig. 1. Sorption isotherms of water for cassava flour (—●—) and for the salts mixture (—△—) as linearized from equation (1)

The initial moisture content of the substrate was 50%; that is, 1.0 g of water/g of solids and the composition of the solid substrate was 0.85 g of flour and 0.15 g of salts. The calculated value gave an a_w of 0.944 for the initial substrate while the measured one was 0.937. It is important to notice that the initial a_w of the substrate was low mainly because of the mixture of salts and that it can be satisfactorily approximated by using the Ross equation.

Growth of *A. niger*

Figure 2 shows the growth of *A. niger* in the

The maximum biomass was reached after about 25 h and as may be seen substrate consumption was not total.

Figure 2 also shows the oxygen uptake rate (OUR); although the maximum respiratory activity took place 3 h prior to maximum biomass production, the analysis of the gas evolution in the exhaust air was a useful tool for estimation of growth (Okazaki et al. 1980; Narahara et al. 1982).

Evolution of the water content during growth

During growth, the amount of total water (H_2O_T) in the medium is the only accessible by measure. The residual water (H_2O_R) in the substrate is given by:

$$H_2O_R = H_2O_T - H_2O_C \quad (3)$$

where H_2O_C is the amount of water consumed which may be estimated by the following equation:

$$H_2O_C = 0.1 S_H + (1/0.33) X \quad (4)$$

where S_H is the amount of sugars hydrolysed
 X is the biomass dry weight
0.33 is the stoichiometric biomass yield for water ($g X/g H_2O$) as determined by Raimbault (1980).

Figure 3 shows the time course of total water as measured during growth and the time course of consumed water and residual water in the medium as calculated from the data reported in Fig. 2.

Total water increased slightly as growth proceeded passing from 1.00 to 1.12 g water/g IDW. The device used in these experiments allowed a good regulation of the relative humidity of the circ-




Fig. 3. Evolution of water during growth of *A. niger* on cassava starch. (H_2O_T =total water, H_2O_C =consumed water, H_2O_R =residual water)

most equal to 1.00 after 22 h. This is, however, the result of two phenomena: an increase in the water-content of the mycelium and a decrease in the residual water in the substrate. The growth pattern of *A. niger* on cassava starch has been illustrated by electron micrographs (Aufeuve et Raimbault 1982). The hyphae gradually invaded the surface of the starch granule which was totally covered by the mycelium after 25 h. It may thus be presumed that if a_w were measured with a device which did not register water loss, as the microorganism colonized the substrate, the mea-



sured a_w would gradually become that of the mycelium, which was almost equal to 1.00.

Consequently, the a_w of the residual substrate was only accessible by calculation. Assuming that the rate of salt uptake was the same as the rate of carbohydrate consumption, the consumption of the salts during growth might be deduced. By inserting the values for residual water (Fig. 3) and for the two ingredients of the medium into the Ross equation (2) it was possible to estimate the a_w of the remaining substrate throughout the fermentation time. As can be seen in Fig. 4, the calculated value of the a_w of the substrate decreases during growth until it falls below 0.90 after 21 h. Such low a_w values are known to inhibit growth; Pitt and Hocking (1977) reported drastic reductions (10 times) in the growth rate of *A. flavus* when decreasing the a_w of an agar plate medium from almost 1.00 to 0.9.

Reduced a_w is supposed to result in lower mass transfers and little water availability for the microorganism and may thus be held responsible for the incomplete conversion of substrate to biomass.

This hypothesis might be verified by working with a higher initial a_w . This could be achieved by reducing the amount of salts or by changing the composition of the salt mixture (sulphate ions for instance are known to produce drastic decreases in the a_w of solutions). Another alternative was investigated by Sato et al. (1982) who introduced a support of high water absorption capacity (wood pulp) to enhance the production of amyloglucosidases and spores by *A. oryzae* on wheat bran. Extending this idea to the SSF of cassava, sugarcane bagasse which is widely available in tropical areas, was chosen for use as a lignocellulosic support since it has, a high water-absorption capacity.

Growth of *A. niger* on mixed media

Studies on water absorption in different mixtures of cassava and bagasse showed that with 20% dry wt. of bagasse, moisture content could be reduced at the outset to 70% without water drainage. Initial moisture contents ranging from 42% to 70% were used, resulting in initial water activities from 0.94 to 0.98. As shown in Table 1, good agreement was shown between the calculated values and the measured ones.

The oxygen uptake rate (OUR) was used as a growth parameter (Fig. 5). It was integrated by the




Fig. 5. Evolution of the oxygen uptake rate during growth of *A. niger* on mixed medium with different initial water contents

method of Sato et al. (1983) and allowed the average specific growth rate to be calculated and then plotted against the initial a_w of the medium (Fig. 6). An increase in the a_w of the substrate caused an increase in the specific growth rate and shortened the spore germination time. With a 0.98 a_w , cultivation took only 20 h and resulted in a high specific growth rate (0.45).




Fig. 6. Evolution of the specific growth rate (—●—) and of the germination time (—◇—) as a function of the initial water activity of the medium

Table 1. Parameters of growth of *A. niger* on cassava starch and bagasse with different initial water contents

IMC %	Initial a_w	Initial a_w *	Incubation time (h) **	X (g/g IDW)	Substrate Consumption (g/g IDW)	Sc*** (%)
42	0.933	0.937	28	0.123	0.233	33.2
48	0.940	0.950	26	0.172	0.344	48.9
55	0.958	0.963	23	0.251	0.417	59.3
65	0.975	0.977	20	0.324	0.477	68.0
70	0.980	0.982	20	0.339	0.558	79.5

* Calculated values

** Time of maximum biomass as calculated from the OUR

*** Substrate conversion: consumed substrate/initial substrate

IMC Initial moisture content

IDW Initial dry weight

As may be seen in Table 1, higher a_w also enhanced the substrate conversion to fungal biomass. A calculation of the evolution of the a_w , made on the same basis as previously described showed that it remained above 0.98 for the entire

creasing and controlling the a_w of the substrate during the process.

Acknowledgements. This work was done as part of the cooperation agreement between the Consejo Nacional de Ciencia y

- Nishio N, Tai K, Nagai S (1979) Hydrolase production by *Aspergillus niger* in solid-state cultivation. *Eur J Appl Microbiol Biotechnol* 8:263—270
- Okazaki N, Sugama S, Tanaka T (1980) Mathematical model for surface culture of koji mold. *J Ferment Technol* 58:471—476
- Pitt JI, Hocking AD (1977) Influence of solutes and hydrogen ion concentration on the water relation of some xerophilic fungi. *J Gen Microbiol* 101:35—40
- Raimbault M (1980) Croissance de champignons filamenteux

- Sato K, Nagatani M, Sato S (1982) A method of supplying moisture to the medium in a solid-state culture with forced aeration. *J Ferment Technol* 61:623—629
- Sato K, Nagatani M, Nakamura K, Sato S (1983) Growth estimation of *Candida lipolytica* from oxygen uptake in a solid state culture with forced aeration. *J Ferment Technol* 60:607—610
- Snell EE (1956) In: *Colorimetric Method of Analysis*. Vol. IV, 3rd Ed. Van Nostrand Co.