

Ethylene production as an indicator of chilling injury in oil palm (*Elaeis guineensis* Jacq.) somatic embryos

Françoise Corbineau^a, F. Engelmann^b and D. Côme^a

^aUniversité Pierre et Marie Curie, Laboratoire de Physiologie Végétale Appliquée, tour 53, 1er étage, 4 place Jussieu, 75252 Paris Cédex 05 and ^bORSTOM, Laboratoire des Ressources Génétiques et Amélioration des Plantes Tropicales, BP 5045, 34032 Montpellier Cédex (France)

(Received February 9th, 1990; revision received May 7th, 1990; accepted May 9th, 1990)

Oil palm (*Elaeis guineensis* Jacq.) somatic embryos are very susceptible to chilling injury. The critical temperature is found to be close to 18°C. The development of injury increases with decreasing temperature and with extending exposure to chilling temperatures. Ethylene production is inhibited at temperatures below 18°C, however it is enhanced after transfer of somatic embryos to warmer temperature (27°C). This chilling induced ethylene production decreases with extending chilling treatment. Results obtained suggest that ethylene release after warming chilled tissues could be a good indicator of chilling injury severity. Repairing of damage induced by chilling temperatures is associated with an increased activity of the enzymatic system (ethylene forming enzyme) which converts ACC to ethylene.

Key words: oil palm; *Elaeis guineensis*; somatic embryos; chilling injury; ethylene; ethylene-forming enzyme

Introduction

121 Dependence on the tissues on the species the

to a method developed by the ORSTOM (Institut Français de Recherche Scientifique pour le Développement en Coopération)/IRHO (Institut de Recherches pour les Huiles et Oléagineux) research team [14] in Bondy, France. The embryoid clumps used were cultured on a standard medium corresponding to the Murashige and Skoog medium devoid of growth regulators and containing 0.1 M sucrose [15]. Each clump comprised adventitious embryos of different sizes and at different developmental stages, which were linked together. Cultures were performed in 37-ml tubes (one embryoid clump per tube). The average weight of embryoid clumps was around 500 mg at the beginning of experiments.

Embryoid clumps were placed in darkness at 10°, 12°, 15°, 18° and 20°C for various durations and then transferred to 27°C, the optimal temperature for their growth [16]. Control cultures were maintained continuously at 27°C.

Growth measurements

Each embryoid clump was weighed at the beginning of experiments and after 6 weeks of culture at 10°, 12°, 15°, 18°, 20° and 27°C. Embryoid clumps that had grown were then divided into parts of about 500 mg. After weighing of each part all embryoid clumps were transferred to 27°C and weighed again 6 weeks later. The increase in weight after the first 6 weeks of culture represented the ability to grow at the different temperatures, whereas weight increase during the second period of 6-week culture quantified the potential recovery of growth at 27°C following the treatment at lower temperatures. Results were expressed as % of weight increase of each embryoid clump, and correspond to the mean of 12 measurements \pm S.D.

Survival of embryoid clumps was evaluated by the number of clumps among a population of 12, which were able to grow at 27°C after various periods of culture at lower temperatures.

Measurement of ethylene production

Ethylene production was measured with embryoid clumps placed at 12°, 18° and 27°C, or at 27°C after various durations at 12° and 18°C. For each measurement the tubes were tightly

closed for 24 h and 1 ml of gas sample was taken with a syringe and injected in a gas chromatograph (type 330, Girdel, France) using a flame ionisation detector, through an activated alumine column (6 mm in internal diameter, 50 cm long, 50–80 mesh). The carrying gas was nitrogen and the column temperature was 60°C. The minimal detectable ethylene quantity was 0.01 nl. Results were expressed as nl of ethylene produced by one embryoid clump in 24 h, and correspond to the mean of 12 measurements \pm S.D.

Treatment with 1-aminocyclopropane 1-carboxylic acid

To determine the capability of somatic embryos to produce ethylene from 1-aminocyclopropane 1-carboxylic acid (ACC), embryoid clumps were first cultured for 6 weeks at 12° and 18°C on the standard medium. They were then soaked for 1 h in a solution of 1 mM ACC and transferred to 27°C on the standard medium containing 1 mM ACC. Ethylene production was measured as previously described, using 12 embryoid clumps, and was compared to ethylene production in the absence of exogenous ACC. Measurements were performed during 14 days after transfer to 27°C. Control embryoid clumps were placed directly at 27°C.

Results

Effects of temperature on growth

The growth of embryoid clumps was very sensitive to temperature (Fig. 1). It decreased sharply from 27° to 18°C, and was almost nil at lower temperatures.

Moreover, a 6-week culture at temperatures lower than 18°C had a strong inhibitory effect on further growth of embryoid clumps after transfer to 27°C (Fig. 2).

Survival of embryoid clumps, evaluated by their ability to grow when transferred to 27°C, depended on the temperature and the duration of previous culture (Table I). All clumps were dead after 6 weeks at 10°C, and after 30 weeks at 12°C. A culture longer than 12 weeks at 18°C also induced the death of some clumps, but all clumps remained viable at 20°C even after 30 weeks.




Fig. 1. Variation with temperature, of weight increase (in % of initial weight) of embryoid clumps after 6 weeks of culture. Mean of 12 measurements \pm S.D.

Clumps which did not grow when transferred to 27°C, and which were then considered as dead, became progressively brown presumably because of oxidation of phenolic compounds.

Effects of temperature on ethylene production

Table II shows the production of ethylene by embryoid clumps cultured at 12°, 18° and 27°C. Ethylene production was relatively high at 27°C,




Fig. 2. Variation with temperature of a 6-week preculture, of weight increase of embryoid clumps placed at 27°C for 6 weeks (in % of the weight at the moment of transfer to 27°C). Mean of 12 measurements \pm S.D.

Table I. Number of surviving embryoid clumps, among populations of 12 clumps, after culture for 6–30 weeks at 10°, 12°, 15°, 18° and 20°C. Survival was evaluated by the ability of clumps to grow when transferred to 27°C.

Temperature of preculture (°C)	No. of surviving clumps after preculture for (weeks)				
	6	12	18	24	30
10	0	0	0	0	0
12	9	7	4	3	0
15	12	9	8	7	4
18	12	12	11	10	10
20	12	12	12	12	12

and it increased with duration of culture probably because of growth of embryoid clumps. At 18°C, ethylene production was much lower and remained almost at the same value (20–30 nl/day by one embryoid clump) during the 11 weeks of experiment though embryoid clumps were able to grow slowly (cf. Fig. 1). At 12°C, the embryoid clumps lost progressively their ability to produce ethylene.

Changes in ethylene production by embryoid clumps placed at 27°C after different durations of culture at 12°C and 18°C, are shown in Fig. 3. A burst of ethylene appeared after transfer of embryoid clumps from 18° to 27°C (Fig. 3A). The increase in ethylene release occurred within the first week at 27°C. It was very marked after 2–6

Table II. Variation of ethylene production (nl/24 h by one embryoid clump) by embryoid clumps during their culture at 12°, 18° and 27°C. Mean of 12 measurements \pm S.D.

Duration of culture (weeks)	Ethylene production (nl/24 h/embryoid clump) at		
	12°C	18°C	27°C
1	20.2 \pm 7.3	28.2 \pm 7.4	365.9 \pm 39.2
2	3.7 \pm 3.8	23.7 \pm 7.9	644.2 \pm 50.3
4	1.1 \pm 0.4	19.5 \pm 6.2	702.7 \pm 78.3
6	0.7 \pm 0.4	19.1 \pm 8.1	868.4 \pm 103.3
8	0.4 \pm 0.3	30.6 \pm 9.2	—
11	0.4 \pm 0.4	32.8 \pm 10.6	—




Fig. 3. Variation during culture at 27°C, of ethylene production (nl/24 h by one embryoid clump) by embryoid clumps previously placed for 1, 2, 4, 6, 8, 11 and 13 weeks at 18°C (A) or 12°C (B). Control unchilled embryoid clumps (C) were placed directly at 27°C. Mean of 12 measurements. Vertical bars denote the largest S.D. Arrows indicate at which times the embryoid clumps were transferred from 18° or 12°C to 27°C. The data points below the arrows represent ethylene production before transfer to 27°C.

weeks of culture at 18°C, and was then reduced when exposure to 18°C was prolonged. The same phenomenon was observed with embryoid clumps transferred from 12° to 27°C (Fig. 3B). However, the burst of ethylene was much less intense and practically disappeared after 6 weeks of culture at 12°C.

Effects of temperature on ACC conversion to ethylene

Figure 4 shows the development of ethylene production at 27°C in the presence of 1 mM ACC, of control embryoid clumps and of embryoid clumps previously cultured for 6 weeks at 12° and 18°C. Control embryoid clumps placed directly at 27°C produced much more ethylene in the presence of ACC than in the absence of this compound (compare with Table II), and they converted more and more ACC during culture




Fig. 4. Variation during culture at 27°C in the presence of 1 mM ACC, of ethylene production (nl/24 h by one embryoid clump) by embryoid clumps previously placed for 6 weeks at 12° or 18°C. Control embryoid clumps (C) were placed




Fig. 5. Variation during culture at 27°C, of the stimulation of ethylene production by 1 mM ACC (expressed as the ratio of ethylene production in the presence of ACC to ethylene production in the absence of ACC) for embryo clumps previously placed for 6 weeks at 12° or 18°C. Control embryo clumps (C) were placed directly at 27°C. The ratio was calculated from mean values of 12 measurements.

produced more ethylene from ACC than the control ones. This production of ethylene occurred immediately after transfer to 27°C, and became maximal after about 1 week. Production of ethylene by embryo clumps previously placed at 12°C was very low during the first week at 27°C. It increased afterwards, but remained lower than with control embryo clumps.

The stimulation of ethylene production by ACC, quantified by the ratio of ethylene production in the presence of ACC to ethylene production in the absence of ACC, is shown in Fig. 5. The value of this ratio was around 4–5 for control embryo clumps placed directly at 27°C, and reached 9–11 for embryo clumps previously placed at 18°C. In the case of embryo clumps precultured at 12°C the effect of ACC on ethylene production was weak during the first 5 days of culture at 27°C, and then increased sharply.

Discussion and Conclusion

Like many organs of tropical and subtropical plants [1,2,13], *Elaeis guineensis* somatic embryos are very susceptible to chilling injury. A minimal

temperature for growth (Fig. 1) and survival (Fig. 2 and Table I) of embryo clumps has been found to exist around 18°C. Similar results were reported by Mok and Hor [17] for the seeds of the same species. The severity of injury increases with decreasing temperature and with extending exposure to chilling temperatures (Table I).

Production of ethylene by oil palm embryo clumps is high at 27°C, whereas it remains low at 18°C and decreases markedly at 12°C (Table II). However, it is strongly enhanced after transfer from 18° to 27°C (Fig. 3A). Such an enhanced release of ethylene is also observed, though at a lesser extent, with embryo clumps transferred from 12° to 27°C (Fig. 3B). A burst of ethylene production after transfer to warmer temperature has been observed with various organs of chilling sensitive plants such as *Citrus* [7], *Cucumis sativus* [11,18,19], *Carica papaya* [12] and *Phaseolus vulgaris* [19]. In oil palm somatic embryos, this enhanced ethylene synthesis decreases with extending chilling treatment (Fig. 3), and is suppressed faster for somatic embryos exposed to 12°C (Fig. 3B) than for those exposed to 18°C (Fig. 3A). It seems, therefore, that the ability of somatic embryos to produce ethylene at 27°C reflects the severity of damage induced by chilling temperatures, and the possible recovery of the cultures after chilling stress. These results suggest that, as previously shown by Chen and Patterson [20], the release of ethylene after warming chilled tissues could be accepted to be a good indicator of chilling severity. A burst of ethylene after transfer to warm temperature is indicative of weak chilling stress, whereas a too severe stress induces an inability to produce ethylene.

In higher plants, ethylene is mainly synthesized through the pathway from *S*-adenosylmethionine (SAM) to ACC and from ACC to ethylene [4], and the enzymatic system ethylene-forming-enzyme (EFE) which converts ACC to ethylene has been shown to be associated with cell membranes [4]. The results obtained by applying exogenous ACC to oil palm somatic embryos show that EFE is affected by chilling temperatures. EFE is activated after treatment for 6 weeks at 18°C, since chilled embryos transferred to 27°C produce more ethylene than control unchilled ones, in the

presence of ACC (Fig. 4). Moreover, the stimulation of ethylene production by ACC is higher in embryos precultured at 18°C than in embryos placed directly at 27°C (Fig. 5). Exposure for 6 weeks at 12°C results in a marked reduction of ACC conversion to ethylene during the first days following transfer to 27°C (Fig. 4). Such an effect of too low a temperature, which was observed with various chilling sensitive materials [11,18,20], suggests that the EFE system is impaired, probably because chilling affects membrane integrity or lipid-protein composition of membranes [1,21]

- 8 R.J. Field, A relationship between membrane permeability and ethylene production at high temperature in leaf tissue of *Phaseolus vulgaris* L. *Ann. Bot.*, 48 (1981) 33—39.
- 9 C.Y. Wang, Changes of polyamines and ethylene in cucumber seedling in response to chilling stress. *Physiol. Plant.*, 69 (1987) 253.
- 10 C.Y. Wang, Relation of chilling stress to ethylene production, in: P.H. Li (Ed.), *Low Temperature Stress Physiology in Crops*, CRC Press, Boca Raton, Florida, 1989, pp. 167—189.
- 11 C.Y. Wang and D.O. Adams, Ethylene production by chilling cucumbers (*Cucumis sativus* L.). *Plant Physiol.*, 66 (1980) 841—843.