

EVIDENCE OF TEXTURAL AND CHEMICAL EVOLUTIVE PATHS BY MEANS OF IMAGE PROCESSING IN WEATHERING AND DIAGENETIC PROCESSES

PARRON C.*, SIMON B.*,**, OLTRA P.H.** and COLIN F. *

* Laboratoire de Géosciences de l'Environnement, URA CNRS 132, Université Aix-Marseille III, 13397 Marseille Cedex 13, France.

** Service de Télédétection et de Sciences de l'image (STSI), Université Aix-Marseille III, 13397 Marseille Cedex 13, France.

I - INTRODUCTION

Both the texture and the chemical composition of rocks and minerals change when affected by low-rate weathering and/or by diagenetic processes. Such changes cannot be accurately defined because the detailed study is impossible by means of classical methods such as optical microscopy or more simply by naked eye. Nevertheless, these changes in texture and in chemical composition reflect the changes of the organization of the matter and thus are good pathfinders to the genesis, and the development of both the weathering and the diagenetic processes. The purpose of this work is to perform a statistical and mathematical algorithm to analyse chemical and textural information from geologic samples.

II- METHOD

The studied samples are glauconite grains and gold particles ranging in size from a few microns to a millimeter. The starting images are obtained by optical or electron microscope. The mathematical treatment of these images is based on the co-occurrence matrix theory, applied to the grey levels of the images, from which it is possible to calculate the contrast, the anisotropy and the entropy (Oltra, 1988). The contrast parameter allows, for a given digitized image, determination of the texture classification, and cartography of textural anisotropies. The images of minerals are obtained from thin sections by a camera through an optical microscope under transmitted light. The mappings of chemical elements are obtained from polished thin sections by microprobe analysis. The images of these mappings are obtained by a camera under

reflected light. This camera is interfaced with a pericolor Numelec 2000 system, generating 256x256 pixel images. The treatment of the digitized images is complemented with a 4381-12 IBM computer.

III - APPLICATIONS

I - Textural anisotropy study of glauconite grains : Evidence of glauconitization process.

A study of crystallochemical and structural mechanisms involved in the genesis of glauconite minerals has been carried out on glauconite-rich rocks from the Paleocene series of the Ivory Coast (Parron, 1989). Unusual glauconite grains show well rounded shapes, internal spherulitic structure and have a diameter ranging from a few microns to 200 microns. Within a given sample, at the scale of the thin section, those grains occur with larger grains (500 to 700 microns), characterized by ovoid and botryoidal shapes and unspherulitic structures. In addition, the smaller grains have a darker green color and a higher amount of K_2O than the larger ones. Such features indicate that the smaller grains result from a more evolved stage of glauconitization (Parron, 1989). The image processing has been carried out on transitional series of grains, in order to draw a parallel between the glauconitization process and the textural changes of the grains. The result demonstrates that all the grains show a textural heterogeneity. Within a given grain, several domains occur, characterized by a decreasing anisotropy from the core to the rim of each domain. The textural

heterogeneity is correlated to the number of textural domains. This number decreases from the larger ovoid grains (7), to the botryoidal grains (3 to 5) and then to the small rounded grains (1). From ovoid to botryoidal grains, the domains of textural anisotropy coincide with protuberances of the botryoidal morphology of the grains. Those protuberances seem to be protograins for the small rounded grains. As a result, the evolution of both the color and the chemical composition of the grains during the glauconitization process simultaneously evolve with size, morphology and internal texture of such grains. The end-members of this evolution by glauconitization process are the smaller grains characterized by rounded shape and textural homogeneity. These grains result from an in situ division of the larger ones.

This result contrasts with those previously reported in the literature which suggests that the smaller spheroidal grains originate from the larger ones by physical abrasion (Boyer, 1977) or generate botryoidal grains by packing (Triplehorn, 1966).

2- Study of chemical evolution by image

silver. To study in details how silver distribution evolves within gold particles during weathering, microprobe analysis mapping has been performed on 21 gold particles. As the silver concentrations are below 9 percent, the mappings are difficult to interpret. To improve the mappings, image processing has been used. About the fresh particles, the results demonstrate that the silver distribution is homogeneous and undependant of the shape of the particles.

This results contrasts with those obtained for the weathered particles. In addition, the preferential leaching of silver, which takes place in the weathered particles, develops either from the rims of the particles or from the intergranular limits inside the particles. The heterogeneity of the silver distribution within gold particles progressively increases from the protore to the surface.

IV - CONCLUSIONS

Two new methods of image processing have been perfected and carried out on rock and mineral samples, which result from low-rate textural and