

Site fidelity, home range and homing behaviour in some species of birds captured at the Ornithological Station of Mbour (Senegal)

F. BAILLON¹ and S. BENVENUTI²

¹ ORSTOM, Institut Français de Recherche Scientifique pour le Développement en Coopération, Station d'Ornithologie et de Mammalogie, B.P. 105, Mbour, Senegal

²Dipartimento di Scienze del Comportamento Animale e dell'Uomo, Università di Pisa, Via A. Volta 6, 56126 Pisa, Italy

and

Centro di Studio per la Faunistica ed Ecologia Tropicali del C.N.R., Via Romana 17, 50125 Firenze, Italy

Received 4 November 1989, accepted 21 December 1989

The authors report data on the site fidelity and home range in some species of birds captured and banded at the Ornithological Station of Mbour (Senegal). In addition, the orientational strategies of individuals displaced to sites 13 and 25 km from the home site were studied in two species of doves (the Laughing dove and the Black-billed Wood dove).

The results show that many of the species investigated display a tendency to spend the whole winter (the non-breeding period for most of the birds considered in our study) in a small home plot of very limited size. Other species, on the other hand, apparently range over a much wider area or are found in our study station only during migratory movements. As regards the initial orientation of the displaced doves, the Laughing dove [*Streptopelia senegalensis* (Linnaeus 1776)] shows a clear tendency to fly in a fixed compass direction with no apparent relationship with the home direction. In the Black-billed Wood dove [*Turtur abyssinicus* (Sharpe 1902)], on the other hand, this stereotyped directional preference tends to be corrected by a weak but significant homeward orientation.

KEY WORDS: home range, homing, orientation, birds, Senegal.

Introduction	58
Materials and methods	58
Results	60
Experiments on site fidelity and home range	60
Homing experiments with Laughing doves and Black-billed Wood doves (<i>Streptopelia senegalensis</i> and <i>Turtur abyssinicus</i>)	62
Discussion	64
Experiments on site fidelity and home range	64
Homing experiments with doves	66
Acknowledgements	67
References	67

ORSTOM Fonds Documentaire

N° : 35 183 ex 1

Cote : b M p49

22 AVR. 1992

INTRODUCTION

The attachment of birds to the places where they live is demonstrated not only by the frequent recaptures of individuals in the original ringing area, but also by the marked homing ability usually observed after birds have been experimentally displaced. This site attachment is pertinent not only to the breeding area, but also to the wintering site (see BENVENUTI & IOALÈ 1980a, 1980b, and IOALÈ & BENVENUTI 1983) and even to a series of places where birds stop during their migratory flights (see WILTSCHKO & WILTSCHKO 1978).

Even if the home fidelity and homing ability of birds have been documented in a variety of species (references for earlier papers are found in MATTHEWS 1968, ABLE 1980), there has been little recent systematic research on this interesting behaviour, with the exception of longstanding investigations on pigeons (references in PAPI 1986; WALLRAFF 1986a), a study on breeding colonies of Sand martins (*Riparia riparia*) (references in BALDACCINI et al. 1989), and a series of studies on birds wintering in the USA (RALPH & MEWALDT 1975, 1976) and in the Mediterranean area (BENVENUTI & IOALÈ 1980a, 1980b, 1983; IOALÈ & BENVENUTI 1982, 1983; IOALÈ et al. 1988).

To make a new contribution to this subject, we performed experiments on the site attachment and homing behaviour of some species of birds wintering or permanently residing in a study area within the Ornithological Station of Mbour in Senegal.

MATERIALS AND METHODS

Study area and capture methods. Our study plot was in the area of the ORSTOM Center (Observatoire géophysique, Station d'Ornithologie et de Mammalogie), 2 km SE of Mbour, Senegal (14°23'N, 16°58'W). The data given here were mainly collected from November 14th to December 5th, 1988; between these dates only, birds were banded and subjected to passive displacements and to other experimental procedures. Additional captures were performed in the following periods: January 10th-14th and 18th-19th, and February 21st-24th, 1989. During these winter capture-sessions, we recorded data on trapped birds, but neither banding activity nor displacements were carried out.

The birds were trapped in mistnets 15 m long and 3 m high, which were in place from sunrise to late afternoon in four «capture stations» (see Fig. 1). Most of the doves and some of the other birds were captured in other sites in the same area where they were sometimes attracted by water and food placed in cage-traps.

Release experiments. Most of the captured birds, which were banded for individual identification and checked for the wing length and body weight, were released in the immediate vicinity of the capture site. Each of the remaining birds, on the other hand, was placed in a separate cloth bag and carried by car to a release site, usually within a few hours of their capture. We used four different release sites: 5 and 25 km NNW, 5 km SE and 13 km SSE of the capture station (home). The only proof of homing ability was taken to be the recapture of the displaced birds at the trapping site. As regards the recapture distance, we have considered the distance between the median points of the two nets in which each single bird was captured and subsequently recaptured. When a bird was trapped twice in the same net, the recapture distance was considered to be zero.

As far as the doves are concerned, we made attempts to study their orientational strategies by releasing them according to the procedures used for the studies on the homing behaviour of pigeons. Releases were accomplished in sunny conditions, with light or moderate wind; the birds were released singly, alternating the two species of doves (the Laughing dove, *Streptopelia senegalensis* and




Fig. 1. — Map of the study area, 2 km SE of Mbour in Senegal. The buildings are offices, laboratories and staff residences of the ORSTOM Observatoire géophysique, Station d'Ornithologie et de Mammalogie. In the four capture stations A, B, C and D, the solid circles represent the sites where mistnets were routinely set up. The nets were not operating every day, and not all simultaneously. Most of the doves and some birds of the other species were trapped at sites outside the usual capture stations.

the Black-billed Wood dove, *Turtur abyssinicus*) and following them up to vanishing with 10×40 binoculars. Vanishing times and vanishing bearings were recorded according to the standard procedures used in this kind of study (see KEETON 1971).

Statistical methods. Directional data have been treated by vector analysis as in earlier papers on homing pigeons. The distributions of the vanishing bearings were tested for randomness using the Rayleigh and the V test (BATSCHLET 1981).

RESULTS

Experiments on site fidelity and home range

Table 1 gives a list of the species of the birds ringed, together with data on recaptures of displaced and non-displaced birds. The numbers of unbanded birds trapped in the winter (January and February) are given too; these numbers can be

Table 1.
List of ringed birds and recapture data. We only give details for the species in which a minimum of seven individuals were banded, or a minimum of two recaptures were recorded.

	Species	N	Rh	Hh	%	Rd	Hd	%	n
1	<i>Ploceus melanocephalus</i> (Linnaeus 1758)	195	104	11	10.6	91	3	3.3	450
2 *	<i>Sylvia borin</i> (Boddaert 1783)	50	30	2	6.7	20	0	0.0	0
3	<i>Streptopelia senegalensis</i> (Linnaeus 1776)	45	2	0	0.0	45 ⁺	4	8.9	7
4	<i>Turtur abyssinicus</i> (Sharpe 1902)	38	2	0	0.0	36	4	11.1	9
5	<i>Prinia subflava</i> (Gmelin 1789)	32	19	8	42.1	13	0	0.0	17
6	<i>Uraeginthus bengalus</i> (Linnaeus 1776)	29	15	1	6.7	14	0	0.0	9
7 *	<i>Phoenicurus phoenicurus</i> (Linnaeus 1758)	21	21	4	19.0	0	—	—	7
8	<i>Cameroptera brachyura</i> (Vieillot 1820)	17	17	8	47.1	0	—	—	5
9	<i>Pycnonotus barbatus</i> (Desfontaine 1789)	17	17	5	29.4	0	—	—	10
10 *	<i>Sylvia atricapilla</i> (Linnaeus 1758)	15	15	0	0.0	0	—	—	0
11 *	<i>Hippolais pallida</i> (Hemprich & Ehrenberg 1833)	14	14	1	7.1	0	—	—	6
12	<i>Sevinus mozambicus</i> (Müller 1776)	13	13	0	0.0	0	—	—	0
13	<i>Ploceus cucullatus</i> (Müller 1776)	11	8	0	0.0	3	0	0.0	113
14 *	<i>Luscinia megarhynchos</i> (Brehm 1831)	10	10	3	30.0	0	—	—	0
15	<i>Lagonostica senegala</i> (Linnaeus 1776)	8	4	0	0.0	4	1	25.0	4
16	<i>Laniarius barbarus</i> (Linnaeus 1766)	8	8	1	12.5	0	—	—	5
17	<i>Pogoniulus chrysoconus</i> (Temminck 1832)	8	8	5	62.5	0	—	—	12
18 *	<i>Hippolais polyglotta</i> (Vieillot 1817)	7	7	0	0.0	0	—	—	0
19 *	<i>Phylloscopus collybita</i> (Vieillot 1817)	7	7	0	0.0	0	—	—	2
20	<i>Sylvietta brachyura</i> (Lafresnaye 1839)	2	2	2	100.0	0	—	—	3
	Total	547	323	51	15.8	226	12	5.3	659

N = number of ringed individual birds for each given species (palearctic migrants have been marked by asterisks); n = number of unbanded birds trapped in January and February; Rh and Rd = birds released at the capture site (home) and at 5 km from home, respectively; Hh (%) = recaptured birds among those released at home and related recapture percentage; Hd (%) = recaptured birds among those released 5 km from the home site and related recapture percentage. + Two L-doves (*Streptopelia senegalensis*) homed after release have been used again in subsequent release experiments; so, in this case, the total number of banded birds (45) is less than the total number of data (2 Rh + 45 Rd).




Fig. 3. — Recapture intervals in 12 species of birds, the same species and the same birds included in Fig. 2. The horizontal lines connect two short vertical marks indicating the capture and the recapture dates for each single bird during the whole capture season (November 14th, 1988, to February 24th, 1989). In the cases in which there were two subsequent recaptures of the same individual, the first recapture is indicated by a vertical mark at an intermediate point somewhere along the horizontal line at the appropriate date. In the right margin, the mean recapture interval is indicated for each species. The arrows indicate the dates on which mistnets were operating. The abbreviations in the left margin indicate the species according to the explanations given in Fig. 2.

capture site, the recapture distances are quite small. In fact, 35 out of 50 birds (70.0%) were recaptured within 50 m of the former capture site. As to the palearctic breeders wintering in, or migrating through, our study plot, we have only nine samples (see Fig. 2), but these too yielded figures consistent with the global result: 6 out of 9 birds (66.7%) were, in fact, recaptured within 50 m of the former capture site.

Data on the recapture interval [time elapsing between the capture and subsequent recapture(s)] are shown in Fig. 3. It turns out that the mean recapture interval is quite long, ranging from 30 to 72 days, the only exception being the Garden warbler (*Sylvia borin*). Thus, in most of the species considered in our study, it appears that individual birds have a tendency to stay in the area for quite a long time.

Homing experiments with Laughing doves and Black-billed Wood doves (Streptopelia senegalensis and Turtur abyssinicus)

Only 26 out of 45 released Laughing (L-) doves (57.8%) and 16 out of 36 Black-billed Wood (BbW-) doves (44.4%) produced useful vanishing bearings; the other




Fig. 4. — Initial orientation of Laughing doves (A and B) and Black-billed Wood doves (C and D). In these diagrams, releases performed on several days and at two different release sites have been pooled with the home direction set to 0° (A and C) and with the North set to 0° (B and D). Each symbol refers to the vanishing bearing of a single bird; open symbols: birds released at 13 km SSE; dotted symbols: birds released at 25 km NNW. The arrows represent the mean vectors, whose length can be read using the scale in the A (top left) diagram. Dashed arrows are used when the bearing distributions do not differ from random according to the Rayleigh test.

birds, in fact, landed soon after the release or were lost behind bushes or trees close to the experimenters. The doves that vanished tended to fly very close to ground level and, due to this behaviour of these birds and to their small body size, vanishing times longer than 1 min were quite rare.

On the whole, nine releases (four at 13 and five at 25 km from home; home direction 328° and 157°, respectively) were performed. The orientational data are presented in Table 2 and in Fig. 4. After pooling the vanishing bearings with the home direction set to 0°, the bearing distribution of both the L- and BbW-dove

Table 2.
 Orientational behaviour of doves: pooled results with the home direction and the North direction set to 0°.

Sp	n	N	Home vector				Compass vector	
			a	r	Dh	Hc	a	r
L	26	(45)	325°	0.16	—35°	0.127	187°	0.60 ***
BbW	16	(38)	354°	0.32	— 6°	0.316 *	232°	0.42

Sp = species of the released birds (L: *Streptopelia senegalensis*; BbW: *Turtur abyssinicus*); n and N = number of recorded bearings and of birds actually released. For each species, the direction (a) and the length (r) of the mean vector is given. Dh and Hc = deviation from the home direction and homeward component, respectively. Significance by the Rayleigh (r) and the V test (Hc) is indicated by asterisks: * = $P < 0.05$, *** = $P < 0.001$.

turned out not to differ from random (Rayleigh test: $P > 0.1$); however, the V test, which estimates not only the scatter but also the angular deviation of the vanishing bearings from the expected (home) direction, reveals that the bearing distribution of the BbW-dove (Fig. 4C), unlike the L-dove, shows a significant clustering around the home direction (V test: $P < 0.05$). After pooling the vanishing data with respect to North set to 0° (Fig. 4, B and D), a tendency to fly Southward (L-dove) or Southwestward (BbW-dove) is revealed (Rayleigh test: $P < 0.001$ and $0.05 < P < 0.1$, respectively).

Only 8 birds were recaptured at the ringing station after the releases: 4 L-doves (three released at 13 km and one at 25 km from home) and 4 BbW-doves (all released at 13 km).

DISCUSSION

Experiments on site fidelity and home range

Our experimental design aimed to investigate the home range, site fidelity and homing success of birds outside the breeding season; this topic has not, in fact, received as much attention as that devoted to the study of the breeding territory, on which much empirical information is available in the literature. The results of this study were also intended to provide the base for a comparative analysis drawing on the results of similar experiments performed on birds wintering in the Mediterranean area, i.e. under very different ecological conditions (BENVENUTI & IOALÈ 1980a 1980b, 1983; IOALÈ & BENVENUTI 1982, 1983).

The present contribution to this subject, however, is limited by several factors, including the difficulty of defining the limits between the breeding and non-breeding season in our area. The period in which our data were collected was outside the main breeding time of the species taken into consideration. However, we cannot exclude the possibility that some of the birds used in our tests were in breeding conditions; Laughing doves, in fact, breed all through the year in this region, and Black-billed

Wood doves may still be active around the nest in November (see URBAN et al. 1986). On the other hand, most small passerine birds — and 17 out of 20 species listed in Table 1 are passerines — are not involved in widespread breeding activity in the period in which our experiments were performed (most birds were recaptured in January or February; see Fig. 3). Of course, this problem only affects tropical species, certainly not the palearctic breeders wintering in tropical regions (7 out of 20 species; numbers 2, 7, 10, 11, 14, 18 and 19 in the list in Table 1). Another problem arises from the fact that we made no attempt to determine the age of the banded birds.

Bearing in mind these limits (including the fact that few data are available for many of the species listed in Table 1), we may try to analyze the relationship between the birds considered in our investigation and the study area. In our study site fidelity is testified and measured by the percentage of the banded birds which have subsequently been retrapped in the area. This recapture percentage varies greatly among the species. The small sample size certainly contributes to this variability, which may also reflect different degrees of site attachment in different species. If we consider the recapture rate of non-displaced birds, the species listed in Table 1 may be distributed in three categories which correspond to three different kinds of relationship between individual birds and their habitat.

(I) The first category includes species of birds attached to a home site of limited extent, as shown by the high recapture percentages, small recapture distances and long mean intervals between the first and last capture, indicating that the birds reside in the area permanently or for a long time. Several species of local birds (*Prinia subflava*, *Camaroptera brachyura*, *Pycnonotus barbatus*, *Pogoniulus chrysoconus*) and of palearctic breeders (*Phoenicurus phoenicurus* and *Luscinia megarhynchos*) fit this pattern of features. It is worth noting once again that the home site in the latter two species is the wintering site, which is temporally and spatially distant from the breeding territory; on the other hand, it is not so easy to determine whether the home site of local residents is both the breeding and non-breeding site, as they may be attached to the breeding territory even in non-breeding periods.

(II) We have included in a second category those species in which the members of the population are not attached to a small home site but range over a large area, which may be much wider than our study plot. In species adopting this strategy we expect long recapture intervals (the population is present in the area, at least during the winter) and low recapture percentages (individual birds range over a wide area and the probability of trapping them twice is rather small). The presence of the species in the area, in spite of the low recapture rates, should be confirmed by frequent captures of non-banded individuals during the trapping season. A local bird (*Ploceus melanocephalus*) and a palearctic breeder (*Hippolais pallida*) fit this set of features.

(III) The third category includes birds migrating through our study plot to reach their winter quarters in other areas. These birds are, therefore, only present for short periods during migration and we expect no recaptures during the winter (but recaptures may occur during the subsequent migratory trip in the opposite direction); recaptures can only occur within a time-span of a few days, giving short recapture intervals. Two species of palearctic breeders can be included in this category: *Sylvia borin* and *Hippolais polyglotta*. For the latter species, the small sample size calls for further investigation, but our data find support in MOREAU (1972), according to whom these birds winter at lower latitudes. The same author reports that the Blackcap (*Sylvia atricapilla*) winters at the latitude of our study plot, but we have no

recapture data for this species. This result, in conjunction with the fact that we have no captures of unbanded individuals during the winter, indicates our study plot is apparently not a favourable area for the Blackcap outside the migratory periods.

In addition, the field data for *Serinus mozambicus*, an African species for which we have no recapture records, fit this category of birds.

As regards the species which have not been included in any of the three categories, the small sample sizes connected with unclear results call for further collection of empirical data.

It is interesting to note that, unlike *Ploceus melanocephalus*, many other species show a global tendency to occupy a small home site, as witnessed by the fact that most of the recaptures were within 50 m of the former capture site (see Fig. 2).

As shown by BENVENUTI & IOALÈ (1980a) and by IOALÈ & BENVENUTI (1983), strong home fidelity is not necessarily related to good homing performances after passive displacement. In fact, several species of wintering passerine birds show high recapture percentages in non-displaced birds, i.e. they are faithful to their home site, but show quite poor homing success after a displacement of 4-8 km. Besides this, there are lower recapture percentages in non-displaced birds than in displaced ones in the data reported here; this aspect of our study, however, is at a very preliminary stage and cannot yet be discussed in detail.

Homing experiments with doves

Considering that homing pigeons use a map based on olfactory information (see PAPI 1986, WALLRAFF 1986a), other species of birds belonging to the same family (Columbidae) are interesting for comparative analysis. Laughing doves and Black-billed Wood doves appear to be particularly suitable for this kind of investigation because they are quite common as resident birds in our study area, can be trapped in good number and can be easily manipulated and subjected to homing experiments.

Compared to pigeons, the orientational behaviour of the doves was quite poor, and a barely significant homeward tendency is only revealed in BbW-doves. Longer mean vectors, however, especially in the case of L-doves, are obtained by pooling the orientational data with respect to North; in this case the birds show a tendency to fly toward the South-Southwestern sector of the horizon. The meaning of this orientational strategy is not clear: both wild birds and homing pigeons often reveal a stereotyped tendency to fly in a compass direction with no apparent relationship to the home direction («nonsense orientation» according to MATTHEWS 1968, or «preferred compass direction» according to WALLRAFF 1986b).

It is worth noting, however, that the doves used in our experiments were captured in an area along the sea, where the coast approximately runs from NNW to SSE. Thus, a compass orientation in directions included in this sector, such as that exhibited by our BbW-doves (but not by L-doves), is the best one for a bird to reach the coastal area from inland regions (the release sites were 2 and 10 km from the sea). A similar orientational strategy is shown by several littoral and river-bank inhabiting arthropods and other animals which exhibit stereotyped oriented movements toward the water direction when they are displaced to dry environments (references in WEHNER & WEHNER 1986); this mechanism, however, would be surprising in birds, which do not simply need to reach an area characterized by optimal ecological

conditions, but a specific small fraction of the proper habitat (i.e. their individual territory).

Firm conclusions cannot be drawn from our experiments, which are a preliminary study whose aim is to verify the feasibility of using L- and BbW-doves for homing experiments. Our tests have shown that these two species of birds can produce oriented vanishing bearings and that the BbW-dove appeared to show superior homeward orientation to that of the L-dove, which exhibited stereotyped southward orientation; further research, with a greater number of samples, is now needed to verify our results and to investigate the orientational strategy of displaced doves.

ACKNOWLEDGEMENTS

We are especially grateful to Dr G. Jarry who kindly supplied us with the rings of the Centre de Recherches sur la Biologie des populations d'Oiseaux (Muséum National d'Histoire Naturelle, Paris). We also thank Prof. F. Papi for his criticism of an earlier draft of the manuscript. This research was supported by the National Research Council (Consiglio Nazionale delle Ricerche).

REFERENCES

- ABLE K.P. 1980. Mechanisms of orientation, navigation and homing, pp. 283-373. In: Gauthreaux S.A. Jr, Edit. Animal migration, orientation and navigation. *New York: Academic Press*, XII+387 pp.
- BALDACCINI N.E., FRUGIS S., MONGINI E. & RAGIONIERI L. 1989. Influence of topographical features on the initial orientation of the sand martin (*Riparia riparia*). In: Orientation and navigation: birds, humans and other animals. *London: Royal Institut of Navigation*.
- BATSCHULET E. 1981. Circular statistics in biology. *New York: Academic Press*, XV+371 pp.
- BENVENUTI S. & IOALÈ P. 1980a. Homing experiments on birds displaced from their wintering ground. *Journal für Ornithologie* 121: 281-286.
- BENVENUTI S. & IOALÈ P. 1980b. Fedeltà al luogo di svernamento, in anni successivi, in alcune specie di uccelli. *Avocetta* 4: 133-139.
- BENVENUTI S. & IOALÈ P. 1983. Age differences in the dispersal behaviour of robins *Erithacus rubecula* studied by counting wing spots. *Ringing and Migration* 4: 237-242.
- IOALÈ P. & BENVENUTI S. 1982. Seasonal and diurnal variation of weight in four passeriformes in autumn and winter. *Avocetta* 6: 63-74.
- IOALÈ P. & BENVENUTI S. 1983. Site attachment and homing ability in passerine birds. *Monitore Zoologico Italiano (Nuova Serie)* 17: 279-294.
- IOALÈ P., BENVENUTI S., GAGLIARDO A., GWINNER E. & SCHWABL H. 1988. Indagine sui fattori che determinano la cessazione della migrazione post-nuziale nella passera scopaiola (*Prunella modularis*). *Naturalista Siciliano (Supplemento)* IV (XII): 241-244.
- KEETON W.T. K. 1971. Magnets interfere with pigeon homing. *Proceedings of the National Academy of Sciences* 68 (1): 102-106.
- MATTHEWS G.V.T. 1968. Bird navigation. *Cambridge: University Press*, X + 197 pp.
- MOREAU R.E. 1972. The Palearctic-African bird migration system. *London: Academic Press*, XV+384 pp.
- PAPI F. 1986. Pigeon navigation: solved problems and open questions. *Monitore Zoologico Italiano (Nuova Serie)* 20: 471-517.
- RALPH C.J. & MEWALDT L.R. 1975. Timing of site fixation upon the wintering grounds in sparrows. *Auk* 92: 698-705.
- RALPH C.J. & MEWALDT L.R. 1976. Homing success in wintering sparrows. *Auk* 93: 1-14.

- URBAN E.K., FRY C.H. & KEITH S. 1986. The birds of Africa (II). *London: Academic Press*, XVI + 552 pp.
- WALLRAFF H.G. 1986a. Relevance of olfaction and atmospheric odours to pigeon homing, pp. 71-80. In: Beugnon G., Edit. *Orientation in space. Toulouse: Privat, IEC*, 127 pp.
- WALLRAFF H.G. 1986b. Directional components derived from initial-orientation data of inexperienced homing pigeons. *Journal of Comparative Physiology (A)* 159: 143-159.
- WEHNER R. & WEHNER S. 1986. Path integration in desert ants. Approaching a long-standing puzzle in insect navigation. *Monitore Zoologico Italiano (Nuova Serie)* 20: 309-331.
- WILTSCHKO W. & WILTSCHKO R. 1978. A theoretical model for migratory orientation and homing in birds. *Oikos* 30: 177-187.