

INTRODUCTION AND SUMMARY OF METHODS AND RESULTS
FROM THE TROPICAL STOCK ASSESSMENT WORKSHOP

Jeffrey J. Polovina, Ronald A. Benco, Albert H. Carlot, Esperance Cillaurren, Paul Dalzell, Ned Howard, Donald Kobayashi, Tevita F. Latu, Paul Lokani, Gianandar Nath, Helen Pitiale, Apesai Sesewa, Richard Shomura, Taniela Sua, Gideon Tiroba, and Sosaia Tulua

A 3-week workshop was convened to assist fishery scientists from selected Pacific island countries in analyzing their research and commercial catch and effort data on deepwater snappers. Workshop participants came from the Cook Islands, Fiji, Kingdom of Tonga, Papua New Guinea, Solomon Islands, Tuvalu, Vanuatu, and Western Samoa. The ultimate objectives of the analyses were to estimate the maximum sustainable yield (MSY) and the fishing effort that achieves the MSY for deepwater snappers in each country. As a simplifying step, all analyses treated this multispecies resource as a single group by combining all species.

Since the deepwater snapper fisheries in all of the Pacific islands represented at the workshop are undeveloped or very recently developed, no long-term time series of catch and effort data are available to apply to production models. An alternate approach estimates unexploited exploitable standing stock (N_0) from the depletion of an area, typically a small seamount or island, and then estimates MSY as a fraction of the unexploited exploitable biomass (B_0 ; Polovina and Ralston 1986). To estimate N_0 from a depletion or intensive fishing study, two analytical models are used. If the depletion has occurred during a short time period so that the number of fish added by recruitment or removed by natural mortality is a minimal part of the population, then the Leslie model (Ricker 1975) is used. This model estimates catchability (q) and N_0 by regressing catch (expressed in numbers of fish) per unit effort (CPUE), against the cumulative number of fish caught. However, if the period covered by the depletion is long enough so that recruitment and natural mortality may have a significant impact on the population, then the Allen model (Sainsbury 1984) is used to estimate catchability (q) and N_0 . The Allen model requires an estimate of natural mortality (M); it estimates q and mean recruitment (R) by regressing catch on two variables: effort and the product of effort with adjusted cumulative catch. Generally when the period covered by the depletion study is less than 9 months, then the Leslie model is used; when the period exceeds 9 months, then the Allen model is used. For the depletion and MSY analyses, whenever an estimate of M is required, two values ($M = 0.25/\text{year}$ and $0.50/\text{year}$) are used to represent a range from the published literature on deepwater snappers (Ralston 1987).

Once estimates of N_0 are obtained from either the Leslie or Allen model, they are adjusted for the habitat area of the depletion site by dividing N_0 by the length (in nautical miles) of the 200 m isobath. The estimates of N_0 are converted to B_0 by multiplying N_0 by mean fish weight.

Estimates of MSY are computed with three approaches: the Gulland method, which estimates MSY as $0.5MB_0$; the Pauly method, which estimates MSY as $2.3(w)^{-0.26}$, where w is the mean (in grams) of the asymptotic weight and the weight at the onset of sexual maturity (Pauly 1983); and a method based on the Beverton-Holt equation in which MSY is estimated as a fraction of B_0 with the fraction as a function of parameters, including M/K , where K is the von Bertalanffy growth constant, and age at recruitment to the fishery (t_r) (Beddington and Cooke 1983). The parameter M/K is estimated at 2.0, based on estimates for M and K for deepwater snappers from the literature (Ralston 1987). The estimated age at recruitment to the fishery is 3 years for Vanuatu and 4 years for all of the other study areas. For each country, an MSY estimate is computed as the product of the MSY range per nautical mile of 200 m isobath and the total length of the 200 m isobath.

Fishing mortality (F) is estimated from catchability, which is estimated from the depletion analyses for each country. The value of q from each depletion analysis is multiplied by the length of 200 m isobath to obtain a standardized q . The mean of all standardized q values from within a country is multiplied by the total fishing effort and then divided by the total length of the 200 m isobath to estimate annual F , which can be interpreted: When F equals M , then fishing effort is optimal to produce MSY; when F exceeds M , then fishing effort is excessive.

The results of 18 depletion analyses from the workshop are presented in Table 1. Table 2 presents a summary of the estimated unexploited exploitable biomass per nautical mile of 200 m isobath from the depletion analyses by seamount and island grouping. Except for a few outliers, most estimates of B_0 cluster around the median values of 2.7 metric tons (t)/nmi for seamounts and 0.7 t/nmi for islands. Since the seamounts are smaller than the islands and have a steeper slope, the difference in the estimates of B_0 /nmi between seamounts and islands may, to some extent, be due to the concentration of biomass over the smaller habitat areas available at seamounts.

The ratios of MSY to B_0 by the Beverton and Holt, Gulland, and Pauly models have a range of about 10 to 30% (Table 3). For an M of 0.25, based on the Gulland estimate, a lower bound of MSY estimated at 9% B_0 is obtained. Conversely, using the Beverton and Holt equation and an M of 0.5 or using the Pauly method, which depends only on fish weight, estimates MSY at about 30% B_0 . Thus, the range of 10 to 30% B_0 is estimated to include the true value of MSY.

Table 4 presents the MSY estimate for each country. For Tonga, current landings from the seamounts are almost twice the upper bound of the estimated MSY, and F is about equal to M . This result suggests that fishing effort is at the level which produces MSY and, once the fishery at the seamounts reaches its equilibrium level, the catches and CPUE will be substantially lower. However, if new seamounts are found, the higher catches may be sustained for a longer time. In Western Samoa, the

Table 1. Summary of results of depletion analyses

| COUNTRY | Area | Type | Principal Species | Nautical Miles of 200 meter isobaths | (Method) | Number of Fish / rmi | Fish Mean Wgt (Kg.) | Unexploited Biomass / rmi (Tonnes) | Recruitment R/rmi/year | |
|------------------|-----------------------|----------|--|--------------------------------------|----------|----------------------|---------------------|------------------------------------|--------------------------------|--------------------------|
| MARIANAS | Pathfinder Reef | Seamount | <i>Pristipomoides zonatus</i> <i>P. auricilla</i> <i>P. flavipinnis</i> <i>Etelis carbunculus</i> | 3.00 | (Leslie) | 1354 | 1.5 | (Leslie) 2.0 | | |
| TONGA | | | <i>P. filamentosus</i> <i>P. flavipinnis</i> <i>Etelis coruscans</i> <i>E. carbunculus</i> | | (Allen) | M=.25 M=.50 | (Allen) | (Allen) | (Number of fish) | |
| | 801 | Seamount | | 5.00 | | 616 | 554 | 4.4 | M=.25 M=.50 2.7 2.4 | M=.25 M=.50 197 344 |
| | 901 | Seamount | | 6.80 | | 614 | 351 | 4.4 | 2.7 1.5 | 137 202 |
| | 903 | Seamount | | 35.00 | | 547 | 608 | 4.4 | 2.4 2.7 | 118 265 |
| | 1004 | Seamount | | 1.20 | | 1927 | 1233 | 4.4 | 8.5 5.4 | 493 762 |
| FIJI | SEAMOUNTS | | <i>Etelis coruscans</i> <i>E. carbunculus</i> | | (Leslie) | | | (Leslie) | | |
| | Taveuni | Seamount | | 3.10 | | 333 | | 7.5 | 2.5 | |
| | Napuka | Seamount | | 4.20 | | 507 | | 7.5 | 3.8 | |
| | Kadavu | Seamount | | 18.00 | | 200 | | 7.5 | 1.5 | |
| | Savusavu | Seamount | | 7.90 | | 189 | | 6.5 | 1.2 | |
| | GRID 147 | Seamount | | 13.10 | | 931 | | 7.5 | 7.0 | |
| FIJI | SHELF | | | | | | | | | |
| | Nasilai to Ovalau | Island | | 50.00 | | 94 | | 7.5 | 0.7 | |
| | Ovalau to MoonReef | Island | | 42.00 | | 95 | | 7.5 | 0.7 | |
| | MoonReef to Ellington | Island | | 40.00 | | 75 | | 7.5 | 0.6 | |
| | GRID 482 | Island | | 40.00 | | 73 | | 7.5 | 0.5 | |
| VANUATU | | | <i>Etelis carbunculus</i> <i>E. coruscans</i> <i>E. radiosus</i> <i>Lutjanus malabaricus</i> | | (Allen) | M=.25 M=.50 | (Allen) | (Allen) | Recruitment R/rmi/year - tonne | |
| | Paama | Island | | 28.00 | | 1277 | 1100 | 1.8 | M=.25 M=.50 2.3 1.1 | M=.25 M=.50 0.30 0.37 |
| | Ambae | Island | | 62.00 | | 389 | 167 | 1.8 | 0.7 0.3 | 0.10 0.13 |
| PAPUA NEW GUINEA | | | <i>P. multidentis</i> <i>P. filamentosus</i> <i>E. carbunculus</i> <i>Wattala mosambitus</i> <i>E. malabaricus</i> <i>Paracaesia stonei</i> | | (Leslie) | | | (Leslie) | | |
| | Schoutten Islands | Island | | 33.00 | | 121 | | 2 | 0.2 | |
| | Kavieng | Seamount | | 0.25 | | 416 | | 2 | 3.3 | |

Table 2. Length of 200m isobaths and Bo/nmi of 200m isobath for all depletion analyses

| | Nautical Miles | Unexploited Biomass / nmi. |
|-----------------------|----------------|----------------------------|
| SEAMOUNTS | | |
| Kavieng | 0.25 | 3.30 |
| 1004 | 1.20 | 8.50 |
| Pathfinder | 3.00 | 2.00 |
| Tavenui | 3.10 | 2.50 |
| Napuka | 4.20 | 3.80 |
| 801 | 5.00 | 2.70 |
| 901 | 6.80 | 2.70 |
| Savusavu | 7.90 | 1.40 |
| GRID 147 | 13.10 | 7.00 |
| Kadavu | 18.00 | 1.50 |
| 903 | 35.00 | 2.40 |
| | | 2.70 = Median |
| ISLAND | | |
| Paama | 28.00 | 2.30 |
| Schoutten Islands | 33.00 | 0.20 |
| MoonReef to Ellington | 40.00 | 0.60 |
| GRID 482 | 40.00 | 0.55 |
| Ovalau to MoonReef | 42.00 | 0.70 |
| Nasilaf to Ovalau | 50.00 | 0.70 |
| Ambae | 62.00 | 0.70 |
| | | 0.70 = Median |

Table 3. Ratios of MSY to B₀ for the three methods

| | MSY/B M=.25 | MSY/B M=.5 |
|-------------------------------|----------------|---------------|
| Gulland (1/2m) | 0.13 | 0.25 |
| Beverton-Holt when m/k = 2 | | |
| age at recruitment = 3 years | 0.06 | 0.18 |
| age at recruitment = 4 years | 0.09 | 0.28 |
| Pauly (2.3w) ^{-0.26} | | |
| Marianas | 0.34 | |
| Papua New Guinea | 0.28 | |
| Western Samoa | 0.27 | |
| Vanuatu | 0.32 | |
| Tonga | 0.28 | |

Table 4. Estimates of MSY, 1988 Landings, and F for deepwater snappers at selected Pacific Island countries

| COUNTRY | Estimated MSY (Tons/yr) | 1988 Landings (Tons/yr) | 1988 Estimated F (fishing mortality) per year |
|---|-------------------------|-------------------------|---|
| TONGA | 77 - 222 | 391 | 0.38 |
| WESTERN SAMOA | 17 - 50 | 25 | (*) |
| FIJI current fishing area of 500 nmi of 200m habitat now being fished | 70 - 200 | 300 | 0.12 |
| FIJI potential fishing area of 3000 nmi | 426 - 1280 | 0 | 0 |
| PAPUA NEW GUINEA | 170 - 270 | 0 | 0 |
| VANUATU | 113 - 190 | 40 | .07 |

(*) No estimate of q available to estimate F

deepwater snapper resource is already close to MSY, but some increase is possible. The fishery in Fiji is currently fishing only about 20% of the estimated 3,000 nmi of 200 m isobath. Thus, Fiji has the potential to increase catches if this unfished habitat has populations of deepwater snappers. However, if the fishery remains only within the habitat currently fished, catches will decline perhaps by 50%. The level of fishing effort may be a little lower than the level which produces MSY. Papua New Guinea has no appreciable landings of deepwater snappers, but a potential exists with MSY in the range of 170 to 270 t/year. Vanuatu's landings represent only one-third to one-fifth of the potential MSY, as indicated by the current landings and level of F ; therefore, the potential exists for greater landings.

A point worth emphasizing is that the results from this workshop represent preliminary estimates of future MSY based on estimates of unexploited biomass and certain mathematical models that project the response of the populations to exploitation. The MSY estimates represent a sustainable equilibrium level, which is lower than the catches that can be obtained in the early stages of the fishery. As the fishery in each country develops, these preliminary estimates of MSY can be improved from a time series of catch and fishing effort data recorded by island.

This workshop also examined the utility of length-frequency data to estimate relative fishing mortality but found that size-frequency samples from the commercial catches do not provide very accurate indicators of fishing mortality, perhaps because of fish size and depth relationships and size-specific targeting by fishermen.

Finally, this workshop focused on the dynamics of the resource and did not consider the dynamics of the fishing operations. In some countries, market conditions and economics of other fisheries may limit fishing for deepwater snappers so that the MSY level is not achieved. In other instances, the distances from port to some of the deepwater snapper grounds may also limit fishing effort below the level resulting in MSY.

CITATIONS

- Beddington, J. R., and J. G. Cooke.
1983. The potential yield of fish stocks. FAO Fish. Tech. Pap. 242, 47 p.
- Pauly, D.
1983. Some simple methods for the assessment of tropical fish stocks. FAO Fish. Tech. Pap. 234, 52 p.
- Polovina, J. J., and S. Ralston.
1986. An approach to yield assessment for unexploited resources with application to the deep slope fishes of the Marianas. Fish. Bull., U.S. 84:759-770.

Ralston, S.

1987. Mortality rates of snappers and groupers. Pages 375-404 in J. J. Polovina and S. Ralston (editors), Tropical snappers and groupers: biology and fishery management. Westview Press, Boulder.

Ricker, W. E.

1975. Computation and interpretation of biological statistics of fish populations. Bull. Fish. Res. Board Can. 191, 382 p.

Sainsbury, K. J.

1984. Optimal mesh size for tropical multispecies trawl fisheries. J. Cons. int. Explor. Mer 41:129-139.

3

NOAA Technical Memorandum NMFS

This TM series is used for documentation and timely communication of preliminary results, interim reports, or special purpose information; and have not received complete formal review, editorial control, or detailed editing.



SEPTEMBER 1990

UNITED STATES AGENCY FOR INTERNATIONAL DEVELOPMENT AND NATIONAL MARINE FISHERIES SERVICE WORKSHOP ON TROPICAL FISH STOCK ASSESSMENT, 5-26 JULY 1989, HONOLULU, HAWAII

Edited by

Jeffrey J. Polovina
Southwest Fisheries Science Center Honolulu Laboratory
National Marine Fisheries Service, NOAA
Honolulu, Hawaii 96822-2396

and

Richard S. Shomura
University of Hawaii
School of Ocean and Earth Science and Technology
Hawaii Institute of Marine Biology
Honolulu, Hawaii 96822

NOAA-TM-NMFS-SWFSC-148

U.S. DEPARTMENT OF COMMERCE
Robert A. Mosbacher, Secretary
National Oceanic and Atmospheric Administration
John A. Knauss, Under Secretary for Oceans and Atmosphere
National Marine Fisheries Service
William W. Fox, Jr., Assistant Administrator for Fisheries

8 JUL. 1992

ORSTOM Fonds Documentaire

N° : 35600, ex 1

Cote : B M P5 IX

NOAA Technical Memorandum NMFS

The National Oceanic and Atmospheric Administration (NOAA), organized in 1970, has evolved into an agency which establishes national policies and manages and conserves our oceanic, coastal, and atmospheric resources. An organizational element within NOAA, the Office of Fisheries is responsible for fisheries policy and the direction of the National Marine Fisheries Service (NMFS).

In addition to its formal publications, the NMFS uses the NOAA Technical Memorandum series to issue informal scientific and technical publications when complete formal review and editorial processing are not appropriate or feasible. Documents within this series, however, reflect sound professional work and may be referenced in the formal scientific and technical literature.