

AGU 1991 FALL MEETING

American Geophysical Union Fall Meeting December 9-13, 1991 San Francisco, California

The AGU 1991 Fall Meeting Abstract Volume is published as a supplement to *Eos, Transactions, American Geophysical Union*, by the American Geophysical Union, 2000 Florida Avenue, N.W., Washington, DC 20009, USA. Second-class postage paid at Washington, D.C., and at additional mailing offices. POSTMASTER: Send address changes to *Eos*, American Geophysical Union, 2000 Florida Avenue, N.W., Washington, DC 20009, USA. Subscription price to regular members is included in annual dues (\$20 per year).

Copyright 1991 by the American Geophysical Union. Material in this publication may be photocopied by individual scientists for research or classroom use. Permission is also granted to use short quotes, figures, and tables for publication in scientific books and journals. For permission for any other uses, contact the AGU Publications Office.

Eos Editor-in-Chief: A. F. Spilhaus, Jr.
Meetings Staff: Group Director of Research and Meetings, L. H. Meredith
Head of Meetings: Brenda L. Weaver
Meetings Manager: Christine Hooke
Meetings Assistant: Kristen Hansen
Meetings Secretary: Steven Bell
Assistant Production Manager: Ronald Scott
Production Coordinator: Judy Castagna
Artists: Don Hendrickson, Dae Sung Kim, Elizabeth Caesar, Nancy Sims, and Renée Winfield



042A-2 1330h POSTER

Interstitial Waters in Atolls and Barrier Reefs : Evidences for Endo-Upwelling Circulation

F Rougerie (ORSTOM, BP 529, Papeete, Tahiti, French Polynesia ; 689-43-98-87 ; Fax 689-42-95-55)
C Andrić (ORSTOM/LODYC, UPMC, case 100, Tour 14, 2° ét, 4 place Jussieu, 75252, Paris Cedex 05, France ; 33-1-44-27-70-77 ; Fax 33-1-44-27-38-05)

Measurements of chemical properties of interstitial waters from shallow boreholes in atoll (35 m), lagoonal pinnacle (17 m) and barrier reef (50 m) in french Polynesia indicate positive anomalous concentrations of dissolved organic nutrients and CO₂ compared to adjacent oceanic and lagoonal waters. Passive tracers such as helium-3 and chlorofluoromethane (F12), characterized by opposite boundary conditions, are used as tools to determine the origin of the nutrient rich interstitial waters. Both tracers data indicate that interstitial water properties result from a mixing of two water masses : Tropical Surface Water (0-100 m) and Antarctic Intermediate Water (500-1500 m). They confirm the penetration and upward circulation inside the reef structure of oceanic water with a recharge depth from 700-800 m. These data give new evidences to the geothermal endo-upwelling model in which Antarctic Intermediate Water enters the porous reef framework, is then driven upward by local geothermal gradient and emerges at the reef crest to provide nutrients to the flourishing algal-coral ecosystem.

4 JUL 1992

ORSTOM Fonds Documentaire

N° : 35705, ex 1

Cote : B

M P21