

These results affect the practice of the drop by drop distribution technique in two ways. First, the use of very dense suspension, in the range of 10,000 to 15,000 nematodes per ml is the most convenient : the variability between drop content is small (Fig. 2) and the time required for collecting the samples is shorter. Secondly, the efficiency of the method used to obtain the mixing of the nematode suspension must be checked for each species.

REFERENCES

- CASTRO, G. A. & FAIRBAIRN, D. (1969). Carbohydrates and lipids in *Trichinella spiralis* larvae and their utilization *in vitro*. *J. Parasitol.*, 55 : 51-58.
- LELLOUCH, M. J. (1964). *Numérations bactériennes, titrage des virus*. Inst. Stat. Univ. Paris. Multigraph. 104 p.
- LUC, M. & TAYLOR, D. P. (1977). *Heterodera oryzae*. *C.I.H. Descr. Pl. Parasit. Nem.* Set 7, n° 91, 3 p.
- MEANS, R. E. & PARCHER, J. V. (1964). *Physical properties of soils*. Constable and Co, Ltd, London, 464 p.
- MORIARTY, F. (1963). Variance of counts from larval suspension of eelworms. *Pl. Path.*, 12 : 177-179.
- PETERS, B. G. (1941). Dilution egg counts and the Poisson distribution series. *J. Helminth.*, 19 : 59-62.
- PETERS, B. G. (1952). Toxicity tests with vinegar eelworm. 1. Counting and culturing. *J. Helminth.*, 26 : 97-110.
- REVERSAT, G. (1976). Étude de la composition biochimique globale des juvéniles des nématodes *Meloidogyne javanica* et *Heterodera oryzae*. *Cah. ORSTOM, Sér. Biol.*, 11 : 225-234.
- SOUTHEY, J. F. (1970). *Laboratory methods for work with plant and soil nematodes*. Technical Bulletin N° 2, London, Ministry of Agriculture, Fisheries and Food.
- WILSON, P. A. G. (1976). The ageing process in infective larvae of the roundworms *Haemonchus contortus* and *Nippostrongylus brasiliensis* : carbohydrate content. *Z. ParasitKde*, 49 : 243-252.

Accepté pour publication le 19 octobre 1979.