

COUPLING A GROWTH MODEL TO A PEANUT RUST MODEL

S. SAVARY
Wageningen Agricultural University

Introduction

Peanut rust and leaf spot are two major foliar diseases of peanut in western Africa. A first attempt was made to build a simulation model for peanut rust which is a newcomer in this part of the world. This work was undertaken as part of a cooperative project between ORSTOM (Office de la Recherche Scientifique et Technique Outre-Mer), and the Wageningen Agricultural University (Savary et al. 1990).

Structure of the Model

Two rust damage components are considered: a reduction of photosynthetically active leaf area, and a flow of carbohydrates used for spore production, which is directly derived from the net photosynthetic rate before any partitioning to the growing organs. Three leaf spot damage components are considered: a reduction of photosynthetically active leaf area, defoliation, and compensation for defoliation.

Crop growth affects the rust epidemic in three ways: the trapping surface of the canopy varies with its LAI, canopy growth makes new sites available for infection, and defoliation reduces the amount of available and occupied sites.

Multiplication and radial growth of *Cercospora* lesions leads to destruction of occupied sites, which is represented by additional terms for the rates of rust lesion mortality in their various stages. This effect is superimposed on that of defoliation.

Model Performances

Evaluation of the model involves verification runs, and comparisons of model outputs with observed data (Teng 1985).

Variations of the daily multiplication factor (Zadoks 1971) and of the pustule spore content with varied weather conditions were found to correspond to the rules introduced in the model. Both variables are in balance, within a range of values which fits estimated values in the field. Various combinations for the values of the varietal coefficients produced a hierarchy of components of resistance which corresponds to observed results from a range of peanut cultivars, and agrees with previous simulation results using similar system designs for the pathogen unit (Zadoks 1971; Teng et al. 1977).

Simulated outputs were compared to actual data representing three separate epidemics, using the living leaf area index and rust severity as reference variables. The similarity between simulated and observed curves, with respect to the timing of the peaks, the day of epidemic upsurge, and the slope of the curves, is encouraging.

In view of its relative simplicity, the performance of this peanut rust simulation model may be considered to comply with the requirements which should be expected from a preliminary model. Several improvements, especially in the crop growth unit, may be considered to increase the explanatory value of the model, and allow its further development.

REFERENCES

- Savary, S., P.D. De Jong, R. Rabbinge, and J.C. Zadoks. 1990. Dynamic simulation of groundnut rust: A preliminary model. *Agricultural Systems* (in press).
- Teng, P.S. 1985. A comparison of simulation approaches to epidemic modeling. *Annu. Rev. Phytopathol.* 23:351-379.
- Teng, P.S., M.J. Blackie, and R.C. Close. 1977. A simulation analysis of crop yield loss due to rust disease. *Agricultural Systems* 2:189-198.
- Van Keulen, H., F.W.T. Penning de Vries, and E.M. Drees. 1982. A summary model for crop growth. p. 87-97. *In* F.W.T. Penning de Vries and H.H. van Laar (eds.) *Simulation of plant growth and crop production*. PUDOC, Wageningen, Netherlands.
- Zadoks, J.C. 1971. Systems analysis and the dynamic of epidemics. *Phytopathology* 61:600-610.
- Zadoks, J.C. and R. Rabbinge. 1985. Modeling to a purpose. p. 231-244. *In* C.A. Gilligan (ed.) *Advances in Plant Pathology*. Vol 3. Academic Press, London.

CONTACT PERSON:

S. SAVARY
ORSTOM, Dept. of Plant Pathology
IIRSDA, B.P. V51 Abidjan
Ivory Coast

PRESENT ADDRESS:

c/o Dept. of Phytopathology
Wageningen Agricultural University
P.O. Box 8025, 6700 EE Wageningen
The Netherlands

WORKSHOP ON