

# *In vitro* Activity and Mutagenicity of Bisbenzylisoquinolines and Quinones Against *Trypanosoma cruzi* Trypomastigotes

A. Rojas de Arias\*, A. Inchausti, M. Acurrat, N. Fleitas, and E. Rodriguez

Instituto de Investigaciones en Ciencias de la Salud, Rio de la Plata y La Gerenza, C. P. 2511, Asunción, Paraguay

A. Fournet

Institut Français de Recherche Scientifique pour le Développement en Coopération (ORSTOM), Département Santé, 213, rue La Fayette, 75480 Paris, Cedex 10, France

The accidental transmission of Chagas' disease by donor blood is recognized as a serious problem in the Latin America. This paper describes the screening of natural products as possible new chemoprophylactic additives in blood banks. Ten plant-derived alkaloids, three terpenes, three quinones and 14 crude plant extracts were tested against bloodstream forms of *Trypanosoma cruzi* Y strain *in vitro* at 4 °C at a concentration of 250 µg/mL, using gentian violet as the baseline drug. The bisbenzylisoquinoline alkaloids, cocsuline, daphnandrine, daphnoline, isochondodendrine, gyrocarpine, limacine and pheanthine and the naphthoquinone, plumbagin completely lysed the trypomastigote forms of *T. cruzi* at a concentration of 250 µg/mL, this activity was verified by the subculture of the treated medium during 4 months. The active alkaloids, pheanthine, daphnoline and limacine were evaluated for mutagenicity by the sister chromatid exchange assay (SCE) in peripheral lymphocytes. Daphnoline and pheanthine elicited no significant increase of the SCE up to 50 µg/mL, while limacine significantly increased the SCE values at a concentration of 25 µg/mL, and 50 µg/mL.

**Keywords:** *Trypanosoma cruzi*; trypomastigote; bisbenzylisoquinoline alkaloids; mutagenicity; naphthoquinones; terpenes.

## INTRODUCTION

Chagas' disease is a widespread infection in Central and South America. It is caused by the protozoan *Trypanosoma cruzi* and is naturally transmitted by Reduviidae bugs (Garcia-Zapata and Marsden, 1986). Blood transfusion is the second most important mechanism of transmission in both endemic and non-endemic areas. This fact is of epidemiological importance due to the migration of infected individuals from endemic rural environments to urban areas. It has been estimated that about 20 000 blood samples could be transmitting the disease annually by blood transfusion in Latin America (Schmuñis, 1985). Another situation relates to hospitals and small health centres in endemic regions, which lack facilities to store blood, thus practising immediate transfusion (Croft *et al.*, 1988; Docampo *et al.*, 1988). The *in vitro* trypanocidal effect of the triphenylmethane dye gentian violet (crystal violet) was first reported by Nussenzweig *et al.* in 1953. Gentian violet is added to blood samples at a concentration of 250 µg/mL and treated blood is stored 24 h before use. The main disadvantages of this treatment are the time required for drug action before transfusion and the colouring of the blood (Docampo *et al.*, 1988). Gentian violet shows no serious toxic effects at recommended concentrations (Rezende *et al.*, 1965). However, a potential carcinogenicity in rodents (Littelfield *et al.*, 1985) and a low mutagenicity have been reported (Docampo and Moreno, 1985; Thomas and

MacPhee, 1984). Furthermore, the use of gentian violet is not easily accepted by patients and medical doctors (Letelier *et al.*, 1990).

Several plant derived and synthetic compounds as well as crude plant extracts have been evaluated as trypanocidal agents against epimastigote and trypomastigote forms of *Trypanosoma cruzi* *in vitro* (Aldunate *et al.*, 1986; Cavin *et al.*, 1987; Croft *et al.*, 1988; Fournet *et al.*, 1988b; Chiari *et al.*, 1991). In the present work, the trypanocidal effect of crude plant extracts and plant-derived compounds, comprising alkaloids, terpenes and naphthoquinones was assessed on the bloodstream forms of *T. cruzi*. The active products were also evaluated for mutagenicity by induction of sister chromatid exchange (SCE) in peripheral blood lymphocyte culture.

## MATERIAL AND METHODS

**Plant material.** The plants were collected in Bolivia between 1985 and 1988. Herbarium specimens were submitted for identification by botanists in the Herbariums in Bolivia, United States and Europe. Voucher specimens were deposited in the National Herbarium of Bolivia in La Paz. Crude extracts were prepared from 12 plants belonging to the botanical families Annonaceae, Araliaceae, Asteraceae, Euphorbiaceae, Lauraceae, Sapindaceae and Ulmaceae. Pure compounds were isolated from natural sources and identified by spectral data (NMR, mass, IR, UV).

\* Author to whom correspondence should be addressed.

27 MARS 1995

**Parasites.** *Trypanosoma cruzi* Y strain infected albino mice were used 7 days after infection. Blood was obtained by cardiac puncture using 3.8% sodium citrate as anticoagulant in a 7:3 blood:anticoagulant ratio. The parasitaemia ranged from  $1 \times 10^5$  to  $5 \times 10^5$  parasites per mL. Plant extracts were dissolved in cold DMSO (dimethyl sulphoxide) to a final concentration of 250 µg/mL. Aliquots of 10 µL were mixed with 100 µL of infected blood, using a microtitre plate in triplicate. DMSO plus infected blood and infected blood alone, and as baseline activity, infected blood with gentian violet at 250 µg/mL were used as controls. Different concentrations of parasites ( $1 \times 10^5$  and  $1 \times 10^6$  parasites per mL) and extracts (4, 20, 40, 100 and 250 µg/mL) were used. The plates were shaken for 10 min at room temperature and kept at 4 °C for 24 h. Each solution was microscopically observed at 400×, placing a 5 µL sample on a slide and covering it with a 22×22 mm overglass for parasite counting (Schempler, 1978). Samples negative to *T. cruzi* by microscopy were cultivated in liver infusion tryptose medium (LIT) and observed for 4 months to detect epimastigote forms (Chiari and Camargo, 1984). Lethal concentrations (LC<sub>50</sub> and LC<sub>90</sub>) were calculated for each extract with a log-probit and means. Variance and standard deviation were calculated for each extract concentration.

**Induction of sister chromatid exchange (SCE).** Human lymphocytes were obtained from healthy donors and cultures were prepared using Hungerford's modified method (Rossner *et al.*, 1987). One half ml of heparinized blood was incubated in 5 mL of RPMI 1640 medium supplemented with 10% bovine serum, 0.1 mL phytohaemagglutinin and 7.5% sodium bicarbonate at

**Table 1. Activity of crude plant extracts towards trypomastigote forms of *Trypanosoma cruzi* in vitro**

Voucher <sup>a</sup>	Botanical name	Part <sup>b</sup> and extract <sup>c</sup>	Lysis at 250 µg/mL (%)
	ANNONACEAE		
A.F. 481	<i>Oxandra espintana</i> (Spruce) Baill.	SB, PE	21
	ARALIACEAE		
A.F. 760	<i>Oreopanax</i> sp.	L, Ch	29
	ASTERACEAE		
A.F. 631	<i>Jungia polita</i> Griseb.	AP, Ch	31
A.F. 616	<i>Munnozia founetii</i> H. Robinson	L, EtOH	16
A.F. 434	<i>Munnozia maronii</i> André	L, PE	29
A.F. 768	<i>Ophryosporus piquerioides</i> (DC) Benth	AP, EA	29
A.F. 675	<i>Senecio hieronymii</i> Griseb.	L, EA	16
	EUPHORBIACEAE		
A.F. 846	<i>Pera benensis</i> Rusby	SB, Ch	31
	LAURACEAE		
A.F. 856	<i>Aniba</i> sp.	SB, Ch	29
		SB, PE	29
A.F. 786	<i>Aniba canelilla</i> H. B. K.	SB, PE	22
		RB, PE	35
	SAPINDACEAE		
A.F. 780	<i>Serjania tenuifolia</i> Radlk.	L, EtOH	17
	ULMACEAE		
A.F. 884	<i>Ampelocera edentula</i> Kuhlm.	SB, Qui.	24
	Gentian violet		100

<sup>a</sup> Voucher: Voucher number refers to A. Fournet's collection.

<sup>b</sup> Part: AP, aerial part; B, bark; L, leaf; S, stem.

<sup>c</sup> Extract: Ch, chloroform; EA, ethyl acetate; EtOH, ethanol; PE, petroleum ether.

**Table 2. Effect of selected plant-derived alkaloids, quinones and terpenes on the trypomastigote forms of *Trypanosoma cruzi* in vitro**

Compounds	Lysis at 250 µg/mL (%)	Reference
Gentian violet	100	
Alkaloids		
Coccoline	100	Lavault <i>et al.</i> 1987
Cycleanine	54	Hocquemiller <i>et al.</i> , 1984
Daphnandrine	100	Lavault <i>et al.</i> , 1987
Daphnoline	100	Lavault <i>et al.</i> , 1987
Gyrocarpine	100	Chalandre <i>et al.</i> , 1986
Isochondodendrine	100	Lavault <i>et al.</i> , 1985
Limacine	100	Lavault <i>et al.</i> , 1986
Obaberine	37	Weber <i>et al.</i> , 1988
Pheanthine	100	Chalandre <i>et al.</i> , 1986
Quinones		
Plumbagin	100	Fournet <i>et al.</i> , 1992
3,3'-biplumbagin	41	Fournet <i>et al.</i> , 1992
8,8'-biplumbagin	39	Fournet <i>et al.</i> , 1992
Terpenes		
Dehydrozaluzanin C	31	Fournet <i>et al.</i> , 1993
Espintanol	2	Hocquemiller <i>et al.</i> , 1991
Isoespintanol	8	Hocquemiller <i>et al.</i> , 1991

37 °C for 72 h. Tested materials were dissolved in DMSO and added to the culture. 5-bromo-deoxyuridine (BrdU) was added to each culture to a final concentration of 5 µg/mL. Incubation was performed in the dark to avoid SCE induced by photolysis of BrdU-substituted DNA. After two rounds of replication, demecolcine (0.02 µg/mL) was added to each culture 2 h before harvesting. Cells were treated with a hypotonic solution, fixed with methanol/acetic acid and dropped on clean chilled glass slides. Differential sister chromatid staining was performed as described by Perry and Wolff (1974). In each sample, 20 metaphases showing differences in chromatid staining were observed and photographed. Terminal SCE were scored as 2. SCE in the centromeric region were not scored because they were indistinguishable from the twisting of sister chromatids.

## RESULTS

From the crude extracts, *Oreopanax* sp., *Jungia polita*, *Munnozia maronii*, *Ophryosporus piquerioides*, *Pera benensis* and *Aniba* sp. (Table 1) showed a third of the activity of the standard trypanocide gentian violet and are worthy of further studies. The naphthoquinone plumbagin showed 100% lysis while the dimeric derivative, 3,3'-biplumbagin and 8,8'-biplumbagin were half as active as the corresponding monomer. The terpenoids dehydrozaluzanin C, espintanol and isoespintanol proved to be weakly active in our assay conditions.

**Table 3. Determination of lethal concentration of compounds tested in vitro against trypomastigotes of *T. cruzi***

Compound	LC <sub>50</sub> (µg/mL)	LC <sub>90</sub> (µg/mL)
Pheanthine	40	175
Limacine	35	180
Daphnoline	8	50
Coccoline	—	4

**Table 4. Frequency of sister chromatid exchanges (SCE) in peripheral lymphocytes induced by active compounds at different concentrations**

Compound	Concentration ( $\mu\text{g/mL}$ )					DMSO 1%
	100	50	25	10	5	
Pheanthine	11.3 $\pm$ 2.6 <sup>a</sup>	8.1 $\pm$ 2.5 <sup>a</sup>	6.4 $\pm$ 1.9	5.3 $\pm$ 1.7	5.3 $\pm$ 1.7	5.9 $\pm$ 1.4
Limacine	ND	23.6 $\pm$ 4.7 <sup>a</sup>	19.5 $\pm$ 1.8 <sup>a</sup>	8.8 $\pm$ 2.3 <sup>a</sup>	8.8 $\pm$ 2.3 <sup>a</sup>	5.9 $\pm$ 1.5
Daphnoline		9.1 $\pm$ 2.5 <sup>a</sup>	7.3 $\pm$ 2.4 <sup>a</sup>	7.1 $\pm$ 2.8	7.1 $\pm$ 2.8	5.8 $\pm$ 2.3

Values are presented as mean $\pm$ SD

<sup>a</sup>Significantly different from control ( $p < 0.001$ )

ND: not done

The bisbenzylisoquinoline alkaloids, coccoline, daphnandrine, daphnoline, isochondodendrine, gyrocarpine, limacine and pheanthine completely lysed the bloodstream forms of *T. cruzi* (Table 2) at a concentration of 250  $\mu\text{g/mL}$  thus being comparable in activity to gentian violet. The means and lytic concentrations ( $\text{LC}_{50}$  and  $\text{LC}_{90}$ ) were determined for four of the ten alkaloids (Table 3).

Daphnoline and coccoline showed an interesting activity when compared with gentian violet. Furthermore, daphnoline was not cytotoxic at concentrations below 10  $\mu\text{g/mL}$ . When it was tested at a concentration at 250  $\mu\text{g/mL}$  under low parasitaemia, no parasites were detected in our conditions. Pheanthine and limacine showed  $\text{LC}_{50}$  and  $\text{LC}_{90}$  fivefold higher than daphnoline. The mutagenicity of pheanthine, limacine and daphnoline was also assessed (Table 4). Pheanthine and daphnoline elicited no significant increase of the SCE up to 50  $\mu\text{g/mL}$ , while limacine significantly increased the SCE values at 25 and 50  $\mu\text{g/mL}$ . The spontaneous frequency of aberrant cells in negative controls was in the range of 0%–5.9%.

## DISCUSSION

All the crude extracts and the pure compounds presented an inhibitory effect on the promastigote forms of *Leishmania* and the epimastigote forms of *Trypanosoma cruzi* *in vitro* (Fournet, 1991). In this study no crude extract was found active against the blood forms of *Trypanosoma cruzi* *in vitro*. Seven bisbenzylisoquinoline alkaloids of nine tested showed an activity on the trypomastigote forms of *T. cruzi*, particularly coccoline and daphnoline which showed no mutagenic effects. The same compounds previously presented an inhibitory activity *in vitro* against different strains of promastigote forms of *Leishmania* spp.

and epimastigote forms of *T. cruzi* at a concentration of 10  $\mu\text{g/mL}$  (Fournet *et al.*, 1988a, 1988b). Recent publications have demonstrated the antiprotozoal properties of the bisbenzylisoquinonines, *in vitro* against *Trypanosoma brucei brucei*, *Plasmodium falciparum* and *P. berghei berghei* (Schiff, 1991). In our study, no structure–activity relationship with the bisbenzylisoquinoline alkaloids was observed.

The only active naphthoquinone, plumbagin is an antiprotozoal compound with activity against *L. amazonensis* and *L. donovani* *in vitro* and *in vivo* (Croft *et al.*, 1985; Fournet *et al.*, 1992) and antibacterial and antifungal activities (Gujar, 1990). Furthermore, by topical application it also shows a significant suppression of skin lesions (Fournet, 1991; Wright and Phillipson, 1990). The result demonstrates again that compounds known to generate free radicals, for example plumbagin, are effective trypanocidal compounds *in vitro*.

In conclusion, it would be interesting to complete this study by testing these alkaloids and other bisbenzylisoquinoline alkaloids against other strains of *T. cruzi*, and also their effectiveness at 37 °C or 25 °C, the same conditions to those in the small health centres of rural Latin America. Furthermore, the compounds tested have the disadvantage of there being no product licences for use in human medicine and it is not easy to isolate the bisbenzylisoquinoline alkaloids by natural or synthetic route.

## Acknowledgements

We are grateful to G. Schmeda-Hirschmann, E. Ferro and A. Figueredo for technical advice. This study was partially supported by Deutsche Gesellschaft für Technische Zusammenarbeit (GTZ). The authors thank Professor J. Bruneton and Dr P. Richomme (Laboratoire de Pharmacognosie, Faculté de Pharmacie d'Angers, France) for supplying the samples of some bisbenzylisoquinolines (coccoline, daphnandrine, daphnoline, gyrocarpine and pheanthine).

## REFERENCES

- Aldunate, J., Ferreira, J., Letelier, M. E., Repetto, Y., and Morello, A. (1986). t-Butyl-4-hydroxyanisole, a novel respiratory chain inhibitor. *FEBS Lett.* **195**, 295–297.
- Cavin, J. C., Krassner, S. M., and Rodriguez, E. (1987). Plant-derived alkaloids active against *Trypanosoma cruzi*. *J. Ethnopharmacol.* **19**, 89–94.
- Chalandre, M. C., Bruneton, J., Cabalion, P. and Guinaudeau, H. (1986). Alcaloïdes de *Gyrocarpus americanus*. *J. Nat. Prod.* **49**, 101–105.
- Chiari, E., and Camargo, E. R. (1984). *Culturing Antigens of Parasites. A Laboratory Manual*. Editorial Fio Cruz, Rio de Janeiro.
- Chiari, E., De Oliveira, A. B., Raslam, D. S., Mesquita, A. A. L., and Tavares, K. G. (1991). Screening *in vitro* of natural products against blood forms of *Trypanosoma cruzi*. *Trans. R. Soc. Trop. Med. Hyg.* **85**, 372–374.
- Croft, S. L., Evans, A. T., and Neal, R. A. (1985). The activity of plumbagin and other electron carriers against *Leishmania donovani* and *Leishmania mexicana amazonensis*. *Ann. Trop. Med. Parasitol.* **79**, 651–653.
- Croft, S. L., Walker, J. J., and Gutteridge, W. E. (1988). Screening of drugs for rapid activity against *Trypanosoma cruzi* trypomastigotes *in vitro*. *Trop. Med. Parasitol.* **39**, 145–148.
- Docampo, R., and Moreno, S. N. J. (1985). Biochemical toxico-

- logy of antiparasitic compounds used in the chemotherapy and chemoprophylaxis of American trypanosomiasis (Chagas' disease). In *Reviews in Biochemical Toxicology* ed. by E. Hodgson, J. R. Bend and R. M. Phipot, pp. 159–204. Elsevier, New York.
- Docampo, R., Moreno, S. N. J., Gadelha, F. R., De Souza, W., and Cruz, F. S. (1988). Prevention of Chagas' disease resulting from blood transfusion by treatment of blood: toxicity and mode of action of gentian violet. *Biomed. Environ. Sci.* **1**, 406–413.
- Fournet, A. (1991). *Plantes Médicinales Boliviennes Antiparasitaires (leishmaniose et maladie de Chagas): Galipea longiflora Krause (Rutaceae), Pera benensis Rusby et Ampelocera edentula Kuhl (Ulmaceae)*. Thèse de Doctorat de l'Université de Paris-Sud, Mention Sciences Pharmaceutiques.
- Fournet, A., Angelo Barrios, A., Muñoz, V., Roblot, F., Hocquemiller, R., and Cavé, A. (1992). Biological and chemical studies of *Pera benensis*, a Bolivian plant used in folk medicine as a treatment of cutaneous leishmaniasis. *J. Ethnopharmacol.* **37**, 159–164.
- Fournet, A., Muñoz, V., Angelo, A., and Aguilar, M. (1990). Plantes médicinales boliviennes antiparasitaires. *Bull. Soc. Fr. Parasitol.* **8**, S9 A 29.
- Fournet, A., Muñoz, V., Manjon, A. M. et al. (1988a). Activité antiparasitaire d'alkaloïdes bisbenzylisoquinoléiques. I. Activité *in vitro* sur des promastigotes de trois souches de *Leishmania*. *J. Ethnopharmacol.* **24**, 327–335.
- Fournet, A., Manjon, A. M., Muñoz, V. et al. (1988b). Activité antiparasitaire d'alkaloïdes bisbenzylisoquinoléiques. II. Activité *in vitro* sur des épimasstigotes de trois souches typifiées de *Trypanosoma cruzi*. *J. Ethnopharmacol.* **24**, 337–343.
- Fournet, A., Muñoz, V., Roblot, F., Hocquemiller, R., Gantier, J. C. and Cavé, A. Antiprotozoal activity of dehydrozaluzanin C a sesquiterpene lactone isolated from *Munnozia maronii* (Asteraceae). *Phytotherapy Res.* **7**, 111–115.
- Garcia-Zapata, M. T. A., and Marsden, P. D. (1986). Chagas' disease. *Clin. Trop. Med. Comm. Dis.* **1**, 557–585.
- Gujar, G. T. (1990). Plumbagin, a naturally occurring naphthoquinone. Its pharmacological and pesticidal activity. *Fitoterapia* **59**, 387–393.
- Hocquemiller, R., Cabalion, P., Fournet, A., and Cavé, A. (1984). Alkaloïdes des Annonacées. XLIX. Alkaloïdes d'*Isolona hexaloba*, *I. zenkeri* et *I. pilosa*. *Planta Med.* **23**–25.
- Hocquemiller, R., Cortes, D., Arango, G. J. et al. (1991). Isolement et synthèse de l'espintanol, nouveau monoterpène antiparasitaire. *J. Nat. Prod.* **54**, 445–452.
- Lavault, M., Bruneton, J., Cavé, A. et al. (1987). Alkaloïdes bisbenzylisoquinoléiques de *Albertisia* cf. *A. papuana* Can. *J. Chem.* **65**, 343–347.
- Lavault, M., Fournet, A., Guinaudeau, H., and Bruneton, J. (1985). Bisbenzylisoquinoline N-oxide from *Curarea candidans*. *J. Chem. Res. (M)* 2786–2793.
- Lavault, M., Fournet, A., Guinaudeau, H., and Bruneton, J. (1986). Alkaloïdes de *Caryomene olivascens*. Nouvelles structures bisbenzylisoquinoléiques. *Chem. Pharm. Bull.* **34**, 1148–1152.
- Letelier, M. E., Rodriguez, E., Wallace, A., Lorca, M., Repetto, Y., Morello, A., and Aldunate, J. (1990). *Trypanosoma cruzi*: a possible control of transfusion induced Chagas' disease by phenolic antioxidants. *Exp. Parasitol.* **71**, 1–7.
- Littlefield, N. A., Blackwell, B. N., Hewitt, C. C., and Gaylor, D. (1985). Chronic toxicity and carcinogenicity studies of gentian violet in mice. *Fundament. Appl. Toxicol.* **5**, 902–912.
- Nussenzweig, V. R., Sonntag, R., Biancalana, A., Freitas, J. L. P., Amata Neto, V., and Kloetzel, J. (1953). Acao de corantes tri-fenil metânicos sobre *Trypanosoma cruzi* *in vitro*. Emprego da violeta de gentiana na profilaxia de transmissao da molestia de Chagas por transfusao de sangue. *Hospital de Rio de Janeiro* **44**, 731–744.
- Perry, P., and Wolff, S. (1974). New Giemsa method for the differential staining of sister chromatid. *Nature* **251**, 156–158.
- Rezende, J. M., Zupelli, W., and Bafutto, M. G. (1965). O problema da transmissao da doenca de Chagas por transfusao de sangue. Emprego da violeta de gentiana como medida profilactiva. *Rev. Goiana Med.* **11**, 35–47.
- Rossner, P., Cerna, M., Lepsi, P., and Pohlova, H. (1987). Mutagenicity of some substituted 1-phenyl-3,3-dimethyltrazene. II Chromosomal aberrations in human peripheral lymphocytes *in vitro*. *Mutat. Res.* **190**, 183–186.
- Schempler, B. R. (1978). Estudos experimentais de quimioprofilaxia da transmissao da doenca de Chagas por transfusao sanguinea. *Rev. Pat. Trop.* **7**, 55–111.
- Schiff Jr, P. L. (1991). Bisbenzylisoquinoline alkaloids. *J. Nat. Prod.* **54**, 645–749.
- Schmuñis, G. A. (1985). Chagas' disease and blood transfusion. In *Immunity and Blood Transfusion*, ed. by R. Y. Dodd and L. F. Baker, pp. 127–145. Alan R. Liss, New York.
- Thomas, S. N., and MacPhee, D. G. (1984). Crystal violet: a direct-acting frameshift mutagen whose mutagenicity is enhanced by mammalian metabolism. *Mutat. Res.* **140**, 165–167.
- Weber, J. F., Fournet, A., and Bruneton, J. (1988). Alkaloidal content of four *Berberis* species structure of berberilaurine, a new bisbenzyltetrahydroisoquinoline. *J. Nat. Prod.* **52**, 81–84.
- Wright, C. W., and Phillipson, J. D. (1990). Natural products and development of selective antiprotozoal drugs. *Phytother. Res.* **4**, 127–139.