

interpretation that CD1b and CD1b-like molecules were deleted during murine evolution. The conclusion that CD1b-like molecules are not a recent innovation is significant as it suggests that cell-mediated immunity directed at non-peptide antigens may have occurred early in the development of the immune system, perhaps even preceding the recognition of peptide antigens by the polymorphic, classical MHC class I and II molecules.

Steven Balk

Division of Hematology-Oncology,
Beth Israel Hospital,
Boston,
Massachusetts 02215, USA

1. Parham, P. *Nature* **372**, 615 (1994).
2. Beckman, E.M. et al. *Nature* **372**, 691-694 (1994).
3. Wang, C.-R. et al. *Molec. Immun.* **25**, 945-952 (1988).
4. Calabi, F. et al. *Immunogenetics* **30**, 370-377 (1989).
5. Bujdosó, R. et al. *J. exp. Med.* **170**, 1285-1318 (1989).
6. Moseley, W.S., Watson, M.L., Kingsmore, S.F. & Seldin, M.F. *Immunogenetics* **30**, 378-382 (1989).

'Functional' haplodiploidy

SIR — The coffee berry borer (*Hypothenemus hampei*) is the main insect pest of coffee worldwide. The recent evolution and rapid spread of high levels of resistance to endosulfan (a cyclodiene-type insecticide) in the South Pacific island of New Caledonia therefore constitutes a major threat to the international coffee industry¹. Like many other members of the scolytid bark beetle, this species shows a highly female-biased sex ratio and incestuous inbreeding²; thus, a single mated female enters the coffee berry and the numerous female progeny are obliged to mate with their few dwarf and flightless male sibs. In some bark beetle species this type of life history (inbreeding polygyny and spanandry) is associated with haplodiploidy².

We have previously shown that cyclodiene resistance in *H. hampei* is associated with a single point mutation in the GABA (γ -aminobutyric acid) receptor subunit gene *Resistance to dieldrin*, or *Rdl*, and that this same mutation is present in several strains from New Caledonia³. Previous insecticide bioassays have also shown that resistance is apparently sex-linked⁴. Thus, consistent with true haplodiploid inheritance, paternally derived resistance is not expressed in males or transmitted through males to progeny.

To test this possibility, we used two PCR-based diagnostic tests, polymerase chain reaction amplification of specific alleles (PASA)³ and single-stranded conformational polymorphism analysis (SSCP)⁵, to determine the genotype of the progeny of reciprocal crosses of F₁ males and females of known parentage. Interestingly, PASA with either the resistant or susceptible specific PCR primer (PASA-R or PASA-S) shows that both resistant and




FIG. 1 SSCP and PASA analysis of males and females homozygous susceptible SS, homozygous resistant RR or heterozygous RS for resistance. Note that in females both maternal and paternal alleles (in parentheses) can be detected by either technique, whereas in male progeny both techniques fail to amplify paternally derived alleles. Thus, banding patterns associated with the predicted genotypes S(S) and S(R) are identical, as are the patterns for R(R) and R(S).



FIG. 2 Cytology of male spermatocyte. The arrow indicates the degenerating paternally derived set of chromosomes and the presumptive maternally derived set are numbered 1-7.

susceptible alleles fail to amplify in males when paternally derived. These data are confirmed by the simultaneous use of SSCP on the same amplification products: SSCP banding patterns for the S(S) and S(R), and the R(R) and R(S) each show the same pattern, because the paternally derived alleles (in parentheses) are not amplified in males (Fig. 1). This again appears consistent with haplodiploidy, in which males would be haploid and females diploid.

Cytological examination, however, reveals that although somatic tissues of females are 2n = 14 as previously reported⁶, in the male soma one of the sets of chromosomes (presumably the paternal

the observed rapid spread of resistance in New Caledonia^{8,9}: first, because recessive or semi-dominant resistance genes are exposed directly to selection in effectively hemizygous males; second, as there is no outbreeding, resistance will be perpetuated within individual, completely isolated, inbreeding lines; and third, within these lines, the continuous maternal backcrossing will promote the rate at which resistance genes achieve homozygosity. Thus, once a resistance-associated mutation has arisen it will be rapidly perpetuated within, and spread via, a single inbreeding line.

Luc O. Brun
Philippe Borsa
Veronique Gaudichon

Institut Français de Recherche Scientifique
pour le Développement en Coopération
(ORSTOM),

Noumea, New Caledonia
Jeffrey J. Stuart
Department of Entomology,
Purdue University,
West Lafayette,
Indiana 47907, USA

Kate Aronstein
Christine Coustau
Richard H. ffrench-Constant
Department of Entomology,
University of Wisconsin,
Madison, Wisconsin 53706, USA

1. Brun, L. O., Marcillaud, C., Gaudichon, V. & Suckling, D. M. *J. econ. Ent.* **82**, 1311-1316 (1989).
2. Kirkendall, L. R. in *Evolution and Diversity of Sex Ratio in Insects and Mites* (eds Wrensch, D. L. & Ebbert, M. A.) 235-345 (Chapman & Hall, New York, 1993).
3. ffrench-Constant, R. H., Steichen, J. C. & Brun, L. O. *Bull. ent. Res.* **84**, 11-16 (1994).
4. Brun, L. O. et al. *J. econ. Ent.* (in the press)
5. Coustau, C. & ffrench-Constant, R. H. *Pestic. Sci.* (in the press).
6. Bergamin, J. & Kerr, W. E. *Clenc. Cult.* **3**, 117-121 (1951).
7. Bull, J. J. in *Evolution of Sex Determining Mechanisms* 145-191 (Benjamin/Cummings, Menlo Park, California, 1983).
8. Brun, L. O., Marcillaud, C., Gaudichon, V. & Suckling, D. M. *Bull. ent. Res.* **80**, 129-135 (1990).
9. Brun, L. O. & Suckling, D. M. *J. econ. Ent.* **85**, 325-334 (1992).

ORSTOM Fonds Documentaire

N° : 41.760 ex 1

Cote : B