

Estimating Surplus Production and Maximum Sustainable Yield from Biomass Data when Catch and Effort Time Series are not Available

S. GARCIA, P. SPARRE and J. CSIRKE

*Fisheries Department, Food and Agriculture Organization of the United Nations,
via delle Terme Caracalla, 00100 Rome (Italy)*

(Accepted for publication 30 January 1989)

ABSTRACT

Garcia, S., Sparre, P. and Csirke, J., 1989. Estimating surplus production and maximum sustainable yield from biomass data when catch and effort time series are not available. *Fish. Res.*, 8: 13-23.

In order to describe a simple method of estimating maximum sustainable yield (MSY), we first demonstrate that the parameters of the two well-known surplus production models of Schaefer and Fox can be expressed in terms of fishing mortality that yields maximum sustainable yield (F_{MSY}), annual yield and mean biomass. If F_{MSY} is known, or, alternatively, is assumed to be some specified function of a known natural mortality rate, it thereby becomes possible to estimate MSY and describe the surplus production function even when the data are limited to one year of estimates for catch and mean biomass. The method is compared to other simple methods for estimating MSY and its potential application is discussed.

INTRODUCTION

In spite of the advances made in the field of fishery sciences and the establishment by most fishery research institutions of data bases for stock assessment purposes, there are still situations where no catch and effort time series exist. In such cases estimation of potential yield is usually based on rough estimates of standing stocks (e.g. from resource surveys) and the general knowledge of some biological characteristics of the species.

Two equations have been proposed in the past to estimate the potential yield of unexploited (Gulland, 1971, p. 2) and exploited (Cadima, in Troadec, 1977) fish stocks for which only estimates of overall biomass and natural mortality are available. Although these were proposed for data-limited situations where the results may be neither very accurate nor precise, the simplicity and ease of application contributed to their extended use. In this paper we discuss the

requirements and sources of bias of these two equations and show the limitations of Cadima's estimator for exploited stocks. Furthermore, we propose an alternative set of simple formulae with which to estimate maximum sustainable yield (MSY) of exploited stocks and draw the surplus production curve using estimates of total biomass and catch during one year as input. The alternatives are based on the most widely used surplus production models of Schaefer (1954) and Fox (1970).

GULLAND'S FORMULA

Gulland (1971) proposed the following estimator of the MSY of a virgin stock when estimates of the natural mortality rate (M) and the biomass of the virgin stock (B_0) are available:

$$MSY = 0.5 MB_0 \quad (1)$$

This estimator has been extensively used with variable success. It is based on the observation that in the Schaefer (1954) production model, the biomass at MSY is equal to half the biomass in the virgin state ($B_{MSY} = 0.5B_0$). It is also based on the assumption already made by Tiurin (1962) and Alverson and Pereyra (1969) that the fishing mortality (F_{MSY}) at MSY is roughly equal to M . It follows that $MSY = F_{MSY}B_{MSY}$ can be replaced by $MSY = M0.5B_0$.

Considering that the ratio between B_0 and B_{MSY} and between M and F_{MSY} could be different for different species groups, Gulland proposed the following generalized form of eqn. (1):

$$MSY = xMB_0 \quad (2)$$

and suggested that x could be estimated from the Beverton and Holt yield tables.

Gulland's equation has been criticized by Francis (1974), Deriso (1982) and Beddington and Cooke (1983), among others, and it is now generally recognized that F_{MSY} is often lower than M and that eqn. (1) overestimates MSY. Therefore eqn. (2), where x is usually smaller than 0.5, should be preferred. For instance, Garcia and LeReste (1981) noted that for tropical penaeids the value of x was in the range 0.32–0.44. Beddington and Cooke (1983) give a collection of graphs from which approximate values of x can be read for various combinations of growth parameters, age at first capture and natural mortality. It is worth noting, however, that these two authors and Gulland assume a den-

CADIMA'S FORMULA

A generalized version of Gulland's estimator was proposed by Cadima (in Troadec, 1977) for exploited fish stocks, for which only limited stock assessment data are available. Cadima's estimator has the form

$$MSY = 0.5ZB_c \quad (3)$$

where B_c is the current average (annual) biomass and Z is the total mortality rate.

Since $Z = F + M$ and $Y_c = FB_c$, the author suggested that in the absence of data on Z the equation could be rewritten as

$$MSY = 0.5(Y_c + MB_c) \quad (4)$$

where Y_c is the total current catch and B_c is the estimated current average biomass. As most stocks are now already exploited, this equation is more frequently used in developing (and some developed) fisheries where catch and effort time series are not yet available, but biomass estimates are occasionally obtained from, for instance, trawl or acoustic surveys.

It should be noted that the validity of eqns. (3) and (4) depends on the current level of exploitation. Equation (3) implies that ZB or $0.5(Y + MB)$ remains constant at all levels of Z . However, by definition (ZB), the total annual production (i.e. the total biomass dying from natural causes and from fishing) varies with Z , as implied by the fundamentals of the production model itself (see Allen, 1971; Csirke and Caddy, 1983). Cadima's estimator is therefore not consistent. Assuming the underlying model (Gulland's formula and the assumption that $F_{MSY} = M$) is correct, Cadima's estimator gives unbiased estimates only when:

- (a) the stock is virgin (then $Z = M, B_c = B_0$ and $Y_c = 0$, and eqns. (3) and (4) become strictly equivalent to Gulland's estimator for virgin stocks, with the same underlying assumptions);
- (b) the stock examined happens to be fished at MSY at the time of the survey for biomass estimates. Then $Z = Z_{MSY} = F_{MSY} + M = 2F_{MSY}$, $Y_c = MSY$, and $B_c = B_{MSY}$, and eqns. (3) and (4), respectively, simplify to $MSY = 0.5Z_{MSY}B_{MSY} = 0.5 \times 2F_{MSY}B_{MSY} = F_{MSY}B_{MSY} = MSY$ and $MSY = 0.5(MSY + MB_{MSY}) = 0.5(MSY + F_{MSY}B_{MSY}) = MSY$

At any other level of exploitation, Cadima's estimator will be biased if we assume the stock to follow a surplus production model. As illustrated later, the use of Cadima's estimator will either overestimate or underestimate MSY depending on whether the stock is exploited below or above the MSY level.

ALTERNATIVES TO CADIMA'S FORMULA

Since there is still a need for gross estimators such as the above to be used in data-limited situations, we propose two general equations, derived directly

from the production models, to estimate the potential yield and the yield curves of exploited fish stocks. These have basically the same foundations and field applications as the Gulland and Cadima estimators, but are more consistent with the underlying models.

The two proposals have been derived based on the two best known and commonly used surplus production models, the Schaefer (1954) model and the Fox (1970) model. Both methods assume that the observations B_c and Y_c are available for one year only. They also assume that M is known and they assume a relationship between M and F_{MSY} of the form

$$M = XF_{MSY}$$

where X is a constant depending on stock parameters.

The two surplus production models may be written as

$$\text{Schaefer: } Y = aF - bF^2 \text{ or } Y/F = a - bF$$

$$\text{Fox: } Y = F \exp(c - dF) \text{ or } \ln(Y/F) = c - dF$$

where a, b, c and d are constants.

As $F = Y/B$ and $Y/F = B$, we can write these surplus production models as follows:

$$\text{Schaefer: } B = a - b(Y/B)$$

$$\text{Fox: } \ln B = c - d(Y/B)$$

If one observation on B_c and Y_c is available and we have a "guesstimate" of F_{MSY} (or X and M), we can write

$$\text{Schaefer: } B_c = a - b(Y_c/B_c) \text{ and } F_{MSY} = XM = -a/2b$$

$$\text{Fox: } \ln B_c = c - d(Y_c/B_c) \text{ and } F_{MSY} = XM = -1/d$$

These two sets of equations can be solved for a and b in the Schaefer model and c and d in the Fox model to obtain

$$\text{Schaefer: } a = \frac{2F_{MSY}B_c^2}{2F_{MSY}B_c - Y_c}; \quad b = \frac{B_c^2}{2F_{MSY}B_c - Y_c} \quad (5a)$$

$$\text{Fox: } c = \ln B_c + Y_c/(B_c F_{MSY}); \quad d = -1/F_{MSY} \quad (5b)$$

Once we have (a, b) or (c, d) we can draw the yield curves and estimate MSY by

$$\text{Schaefer: } MSY = -a^2/4b \quad (6a)$$

$$\text{Fox: } MSY = -(1/d) \exp(c-1) \quad (6b)$$

Thus, one pair of observations (B_c, Y_c) and assumptions on M and the relationship between M and F_{MSY} ($F_{MSY} = XM$) are sufficient information to get a first rough estimate of the yield curve and of MSY.

Schaefer model

Let us assume that the stock in question responds to the Schaefer production model. By replacing a and b in eqn. (6a) by their value in eqn. (5a) we get

$$MSY = \frac{(F_{MSY} B_c)^2}{2F_{MSY} B_c - Y_c} \quad (7a)$$

In the special case that $F_{MSY} = M$, the estimator becomes

$$MSY = \frac{(M B_c)^2}{2M B_c - Y_c} \quad (7b)$$

If the stock is unfished (i.e. when $F=0$, $Y_c=0$ and $B_c=B_o$), eqn. (7b) becomes Gulland's original equation. It should also be noted that eqn. (7b) and eqn. (4) are equivalent only when $Y_c = M B_c$ and, therefore, when $F=M$. This shows that Cadima's estimator is consistent with Gulland's equation and unbiased at MSY, but gives erroneous values at any other level of exploitation.

It is also worth noting that if F approaches $2F_{MSY}$ and Y and B_c approach zero estimators (7a) and (7b) become highly unstable, particularly consider-

PERFORMANCE OF THE ESTIMATORS

The performance of the rough estimators for MSY depends on the accuracy of the data used (B_c , Y_c) and on the appropriateness of the underlying assumptions of the selected estimator. Can the surplus production of the stock be represented as a function of a Schaefer- or a Fox-type model? Would it not be better approached by a simple Beverton and Holt yield per recruit function, i.e. assuming that recruitment is independent of stock size? Was the stock really in equilibrium when the biomass was measured? It is, of course, not possible to answer these questions in the absence of long data series, but it is useful to look at the sort of effects one should expect if the assumptions underlying the rough estimator do not correspond to the properties of the stock in question.

To illustrate these questions, three sets of artificial yield and biomass data were simulated under the assumption that the stock behaved as predicted by the Beverton and Holt model, the Schaefer model or the Fox model. Using these simulated observations as input to Cadima's estimator and to the two




Fig. 1. Comparison between the new MSY estimators based on the Schaefer (eqn. 7b) and Fox (eqn. 8b) models, and Cadima's MSY estimator when the stock follows a Beverton and Holt yield model.

$$Y = F \exp \left[-M(t_c - t_r) \right] RW_{\infty} \left[\frac{1}{Z} - \frac{3S}{Z+K} + \frac{3S^2}{Z+2K} - \frac{S^3}{Z+3K} \right]$$

where $S = \exp -K(t_c - t_o)$. In this model $F_{MSY} = 0.207$ and $MSY = 526$ with the following parameters: $K = 0.3$; $t_o = 0$; $t_c = t_r = 0$; $RW_{\infty} = 10\ 000$.

alternative estimators proposed in this paper, we can study their behaviour under various assumptions about the dynamics of the stock.

The simulated yield and biomass curves are shown in Figs. 1–3, which also give the values of the parameters used. To facilitate comparisons, the (arbitrary) parameters behind the simulated data sets are selected so that the yield when $F=M$ ($=0.3$) takes the same value (the arbitrary value of 500) in all three cases.

When the real stock production function is the Beverton and Holt model (1957), Fig. 1 shows that for $F=M$ ($\simeq F_{\text{MSY}}$) Cadima's estimator and eqn. (7b) or (8b) give the same results, and these are very close to the "true" MSY as defined by the Beverton and Holt model (Gulland's estimator would give the same result as Cadima's at $F=0$). All estimators seriously overestimate the MSY when used at levels of effort below M . Beyond $F=M$ the estimators diverge greatly, under or overestimating MSY. If a 20% error margin is accepted, however, it can be seen that, for instance, the Fox-model-based estimator (eqn. 8b) gives acceptable results within a wider range of F than Cadima's estimator

$0 \leq F \leq F_{MSY}$, considering the other uncertainties in the process, but the performance of Cadima's estimator drops drastically beyond F_{MSY} .

When the real production function is a Fox model, Fig. 3 shows that Cadima's estimator (as well as Gulland's) overestimates MSY when $F=0$. At MSY the former answer is correct. For $F > F_{MSY}$ MSY is largely underestimated, while eqn. (8b) finds the correct MSY at all levels of fishing.

Validation of the estimates

By definition, a model is a simplified representation of a complex phenomenon used to illustrate one or more processes (in the latter the evolution of total catches as a function of fishing). However, one must bear in mind that even if the model was conceptually correct, using a punctual estimate (such as the stock size $= B_c$ in a given year) to determine the potential yield of a stock and its trajectory under various fishing levels is an adventurous, although sometimes necessary, exercise. Due caution should therefore be taken in selecting an estimator, in validating the estimates and in expressing management or development advice on the basis of such estimators.

When working with the above equations, fishery scientists should check whether the basic assumption of the models are fulfilled. For example, the biomass (B_c) is meant to be the exploited average biomass, and both the catch and the biomass referred to should have the same age (or size) structure. For instance, the biomass estimate should not include small sizes which are not available to the fishery (as could be the case when using acoustic estimates). This biomass is the annual biomass value and seasonal oscillations caused by




Fig. 3. Comparison between Cadima's MSY estimator and a new MSY estimator (eqn. 8b) where the stocks follow a Fox model with the following parameters: $M=0.3$; $d=(1/M)=3.3333$; $MSY=500$. Note that $c=1+\ln(MSY/M)$.

changes in growth, mortality or recruitment, which are likely to be more important in short-living species such as shrimps, squids, anchovies, etc., should be taken into account and as far as possible levelled off to obtain an appropriate annual average of the total biomass. If seasonal oscillations of biomass are caused by migrations, then the peak biomass representing the real stock size should be used. If another country is exploiting the same stock at another season of the year in another area, the total catches should be included in the calculations and the MSY estimate should refer to the whole unit stock.

The estimated potential yield could also be validated by comparison with other stocks for which information might be available. Some of the questions to be asked are: how does the estimate of MSY/km^2 stand with respect to the estimates for other stocks of the same species in ecologically similar areas, exploited under similar fishing regimes?; does the size structure of the catch provide support for assessments implying that the stock is heavily overfished (e.g. predominance of juveniles) or underfished (e.g. predominance of large, old fish)? It should be noted that a closer look at the length-frequency composition of the catches would give some guidance on relative levels of exploitation, and this should be among the first data to be collected in any developing fishery in order to allow some estimations of total mortality rate for cross-

models from which they were derived. These have been widely discussed by several authors (see e.g. Larkin, 1977; Sissenwine, 1978; Garcia and LeReste, 1981) and the main assumptions are that the biological processes involved are deterministic, the fishery is on a single stock with stable age/size characteristics, the catchability is not density-dependent, and there are no time lags. In addition, the proposed estimators use biomass estimates (as well as F_{MSY} and M estimates) that are usually poorly estimated or based on educated guesses. This may be particularly critical if no sound auxiliary research data are available to constrain the estimates or assess their reliability.

The comparison of estimators' performance has not exhausted the possible combinations of errors which can be made. The amount and direction of the error depend on the quality of the input data, the level of fishing when the estimate is made, and the degree of coherence between the assumptions underlying the estimator used and the reality of the stock. These examples show that non-consistent estimators, like eqns. (3) and (4), can generate errors, especially beyond F_{MSY} . They also show that estimators which are internally coherent can lead to error if the resource is best described by a different model from the one on which the estimator is based. These differences and their consequences on management decisions could be very important, even for an approximate first estimate. It is therefore advisable to use as many models as

possible (Y/R , Fox and Schaefer) in order to have some idea of the degree of uncertainty attached to the MSY estimate and to the shape of the production curve (especially beyond F_{MSY}).

Fishery scientists are often asked to express their results in terms usable by managers, and in data-poor conditions this may mean relying, to a great extent, on formulations like the ones proposed here until proper data files are built up

assessment methods. It must be pointed out, however, that most of these estimators become highly unreliable at high levels of effort and therefore should not be used under such conditions.

The proposed methods should not be considered as a substitute for the classical catch and effort analysis, but rather as a more robust substitute for the Cadima estimator. It is our opinion that the new proposals, which provide an estimation of MSY as well as a first approximation to the surplus production curve, may offer a better basis for a preliminary management-oriented analysis.

ACKNOWLEDGEMENT

The authors are grateful to Dr. Ian F. Somers, Division of Fisheries Research, CSIRO, Australia, for his useful and constructive comments on an earlier draft of this paper.

REFERENCES

- Allen, K.R., 1971. Relation between production and biomass. *J. Fish. Res. Board Can.*, 28: 1573-1581.
- Alverson, D.L. and Pereyra, W.T., 1969. Demersal fish exploitations in the Northeastern Pacific Ocean. An evaluation of exploratory fishing methods and analytical approaches to stock sizes and yield forecasts. *J. Fish. Res. Board Can.*, 26: 1985-2001.
- Beddington, J. and Cooke, J.G., 1983. The potential yield of fish stocks. *FAO Fish. Tech. Pap.*, 242: 47 pp.
- Beverton, R.J.H. and Holt, S.J., 1957. On the dynamics of exploited fish populations. *Fish. Invest. Minist. Agric. Fish. Food (G.B.) Ser. II Salmon Freshwater Fish.*
- Caddy, J.F. and Csirke, J., 1983. Approximations to sustainable yield for exploited and unexploited stocks. *Oceanogr. Trop.*, 18(1): 3-15.
- Csirke, J. and Caddy, J.F., 1983. Production modeling using mortality estimates. *Can. J. Fish. Aquat. Sci.*, 40: 43-51.
- Deriso, R.P., 1982. Relationship of fishing mortality and growth and the level of maximum sustainable yield. *Can. J. Fish. Aquat. Sci.*, 39: 1054-1058.
- F. W. L. 1979. A general model for the yield of a fish population.