

6. Oceanapins A-F, Unique Branched Ceramides Isolated from the Haplosclerid Sponge *Oceanapia cf. tenuis* of the Coral Sea

by Ines Mancini^{a)}, Graziano Guella^{a)}, Cécile Debitus^{b)}, and Francesco Pietra^{a)*}

^{a)} Istituto di Chimica, Università degli Studi di Trento, I-38050 Povo-Trento


^{b)} ORSTOM, Centre de Nouméa, B. P. A5, Nouméa Cédex, Nouvelle Calédonie

Dedicated to the memory of *Margherita Ratti*

(17.V.93)

A series of ceramides, called oceanapins A-F (2-7), which are unique for branching at both the sphingosine and fatty-acid chains, have been isolated as pure compounds from the haplosclerid sponge *Oceanapia cf. tenuis* of the Coral Sea. Following acid hydrolysis, both the fatty-acid and the sphingosine portions were obtained separately, which allowed their unequivocal structural definition. The absolute configuration was secured *via* protection of C(1')-OH and Mosher's esterification at C(3')-OH of the oceanapins.

1. Introduction. - Sphingosine (1) [1] is long known from hydrolysates of brain tissues [2] which, like all animal sphingolipids [3], also contain sphingosine amides of long-chain fatty acids (ceramides) and their glycosyl derivatives (cerebrosides).




Sphingosine derivatives were later also isolated from seaweeds, both green [4] [5] and red [6], symbiotic dinoflagellates [7], and sea anemones [8]. All these sphingosine derivatives are built on linear chains. Isopropyl [9] [10] or ante-iso termini [11] have only been found in the sphingosine portion of ceramides [9] [10] and cerebrosides [11] of a few other marine invertebrates.

With few exceptions [7] [8], sphingosine derivatives were found to occur as mixtures of homologous or isomeric compounds, the separation of which has never been achieved [4-6] [9-12], though eagerly solicited [13]. We report here on the isolation from the tropical sponge *Oceanapia cf. tenuis* (Haplosclerida) of pure homologous or isomeric ceramides that are unique for branching on both the sphingosine and the fatty-acid chains. This allowed us also to assign their absolute configuration.

2. Results and Discussion. - 2.1. *Chromatographic Separation.* A combination of FC, CN- and RP-HPLC proved to be an excellent method for separating oceanapins 2-7



(Scheme 1). As expected, the observed retention times correlate with the length of the lipophilic chain. Accordingly, oceanapins with short and/or branched side chains are eluted first (Table).



a) 1.2M H₂SO₄ in 85% MeOH, reflux, 6 h.

Table. HPLC Retention Times vs. Chain Length and Relative Abundance for Oceanapins 2-7

Oceanapins	No. of C-atoms			Relative abundance ^{a)}	<i>t_R</i> ^{b)}
	total	fatty-acid chain	sphingosine chain		
2	39	23	16	3	13.2
3	40	23	17	1	14.1
4	40	23	17	2.5	15.2
5	41	23	18	7	16.2
6	41	22	19	3	17.4
7	42	23	19	10.7	18.5

^{a)} From area integration of reversed-phase HPLC peaks with MeOH, 1 ml/min, λ 215 nm.

^{b)} HPLC retention times under conditions defined in ^{a)}.


2.2. *Connectivity and Relative Configuration.* For the abundant oceanapin F (7) C-atoms were assigned *via* ¹H, ¹³C correlations, suggesting the presence of two OH substituents from signals for a deshielded CH₂ (δ (H) 3.70, 3.96) and CH group (δ (H) 4.29), correlated to the δ (C) 62.52 (*t*) and the δ (C) 74.72 (*d*) signals, respectively. An amide functionality (CONH) was also suggested by the signals at δ (C) 173.85 (*s*) and δ (H) 6.30 (*br. d*) signals, and by correlation of the signal at δ (H) 3.89 (*dt*) with δ (C) 54.45 (*d, CHNH*), in agreement with IR absorption bands at 3360 and 1640 cm⁻¹. An (*E*)-olefinic bond was indicated by signals at δ (H) 5.51 and 5.76 (*J* = 15.3 Hz) correlated to 2 *d*'s at δ (C) 128.80 and 134.30, respectively. At this point, differential decoupling irradiations and ¹H, ¹H and long-range ¹H, ¹³C experiments allowed us to establish the C(1')-C(5') portion (*Exper. Part*).

Long aliphatic chains were suggested by a δ (H) 1.24 (*br. s*) and a series of *t* between δ (H) 29.1 and 30.0 that also occur with the other oceanapins. The chains proved to be either unbranched (δ (H) 0.87 (*t*), correlated to δ (C) 14.09 (*q*), for 2, 4, and 6), or branched with either *i*-Pr terminus (δ (H) 0.85 (*d*), correlated to δ (C) 22.64 (*q*) and 27.95 (*d*), for 2-5 and 7) or ante-iso terminus (δ (H) 0.84 (*t*) and 0.85 (*d*), correlated to δ (C) 11.39 (*q*), and 19.21 (*q*), and 34.38 (*d*), for 6 and 7). Branching of the ante-iso type for oceanapins 6 and 7 was also supported by calculations through the empirical Lindeman-Adams rule [14].

Mass spectra afforded further structural insight by showing fragment ions for loss of H₂O from the molecular ion, besides diagnostic fragment ions at either m/z 378 for 2-5, and 7 or m/z 364 for 6¹) of compositions C₂₄H₄₆NO and C₂₅H₄₈NO (HR-EI-MS) for 6 and 7, respectively, which must derive from the cleavage of the C(2)-C(3) bond. EI-MS data (*Exper. Part*) indicated a C₂₃ amidic chain for oceanapins B (3) and D (5), which have the same chain terminus.

The whole C-C connectivity was assigned from the sphingosine and fatty-acid moieties obtained as pure compounds from acid hydrolysis of the oceanapins (sphingosines 8a-d and methyl esters 9a-c; *Scheme 1*).

The *erythro*-configuration for acetamide 12, prepared from oceanapin F (7; *Scheme 2*), was deduced from the identity of the NMR spectra (δ (H-C(2')) 3.83 ppm, $J(2',1') = 5.2$ and 9.1, $J(2',3') = 9.6$) with acetamide 11, prepared from the *N*-acyl derivative 10 of commercial D-*erythro*-sphingosine (1). This established *erythro*-configuration also for oceanapin F (7). This conclusion was also supported by molecular-mechanics calculations for acetamide 11 in both *threo*- and *erythro*-configuration. Thus, calculations for the preferred conformation of 11 (*erythro*) gave coupling constants in agreement with the experiment: $J(2',1') = 5.2$ and 11.0 (for torsional angle 2 H-C(1')-C(2')-H 55° and 172°), and $J(2',3') = 9.9$ (for torsional angle H-C(2')-C(3')-H 177°). In contrast, the corresponding J for the preferred conformation of 11 (*threo*) were found to be different from 12: $J(2',1') = 2.4$ and 1.7, corresponding to torsional angle 2 H-C(1')-C(2')-H of 51° and 67°.




a) Ac₂O, EtOH, 0°, 1 h. b) Acetone, CuSO₄, reflux, 6 h.

2.3. Absolute Configuration. Previous assignments of the absolute configuration of acylsphingosines were either based on CD measurements on mixtures, such as for caulerpicin (2'*S*,3'*R*) [4c] and the ceramide portion of cerebroside [15], or on total synthesis, such as for symbioramide [7] (= (2*S*,3*R*,2'*R*,3'*E*)-*N*-(2'-hydroxyoctadec-3'-enyl)-dihydrosphingosine) [16] and the sphingosine derivative of *Anemonia sulcata* [10], (2*R*,3*S*) [17].

For the oceanapins we have decided to prepare both Mosher's esters [18] (+)-14a and (-)-14b of the silyl ether (+)-13 [19] derived from 5. The NMR data, expressed as $\Delta\delta$ ($\delta_S - \delta_R$) in *Scheme 3*, show that the ¹H-NMR signals for 2 H-C(1'), H-C(2'), 2 H-C(2'') and N-H of ester (+)-14a are observed upfield from the corresponding protons of ester (-)-14b. The opposite trend is observed for H-C(4'), H-C(5'), and 2 H-C(6'). This result can be consistently explained by the diamagnetic effect of the Ph ring, thus indicating the (*R*)-configuration at C(3'). Since *erythro*-configuration has been estab-

¹) Reversed-phase HPLC (*Table*) gave two other homologous oceanapins, albeit in too small amounts (ca. 0.2 mg each) for complete structural study. The one eluted at t_R 11.6 min gave two significant EI-MS fragment ions, m/z 561 ($[M - H_2O]^+$), indicating a total of 37 C-atoms, and m/z 336, suggesting a C₂₀ amidic chain. For the oceanapin eluted at t_R 12.4 min, fragment ions m/z 364 and 589 ($[M - H_2O]^+$) indicated a C₂₂ amidic chain or a total of 39 C-atoms, respectively. For the latter, comparison of t_R with oceanapin A (2: *Table*) suggested branching at both chains.



a) t BuPh₂SiCl, 1*H*-imidazole, DMF, 60°, 12 h.

b) (*S*)-MTPA, pyridine, CCl₄, r.t., 24 h.

c) (*R*)-MTPA, pyridine, CCl₄, r.t., 24 h.

^{a)} The $\Delta\delta = \delta_S - \delta_R$ represent the difference, in Hz, of the corresponding ¹H-NMR resonances between the (*S*)-MTPA ester (–)-14b and the (*R*)-MTPA ester (+)-14a.

lished between C(2') and C(3'), the absolute configuration of oceanapin D (5) is thus (2'*S*,3'*R*).

We thank Dr. *Jane Fromont*, Sir Fisher Marine Biology Institute, University James Cook, Townsville Qld., for the sponge identification, Mrs. *M. Rossi* and *A. Sterni* for skilled technical aid, and *MURST* (Progetti 40%) and *CNR* (Roma) for financial support. This work has been carried out within the collaborative program *ORSTOM-CNRS* on 'Marine Substances of Biological Interest'.

Experimental Part

1. *General*. All evaporations were carried out at reduced pressure. Yields are given on reacted compounds. M.p.: *Kofler* hot-stage microscope. Pyridine was freshly distilled from BaO and DMF (stored on flamed 4 Å molecular sieves) were used. Flash-chromatography (FC): *Merck Si-60*, 15–25 μm. TLC: *Merck 'Kieselgel' 60 PF₂₅₄* plates. HPLC: *Merck LiChrosorb CN*; reversed-phase (RP) HPLC: *Merck LiChrosorb RP18*; 7 μm, 25 × 1-cm columns; solvent flux 5 ml/min; UV monitoring λ 215 nm, if not otherwise stated. Polarimetric data: *JASCO-DIP-181* polarimeter. IR: *Perkin-Elmer-337* spectrometer; ν_{\max} in cm⁻¹. NMR (CDCl₃): δ in ppm rel. to internal SiMe₄ (= 0 ppm), *J* in Hz; *Varian-XL-300* spectrometer; ¹H at 299.94 MHz (the coupling pattern of many protons has been clarified by differential decoupling irradiations [20] and ¹H,¹H COSY [21]); ¹³C at 75.43 MHz, multiplicities from DEPT experiments [22]; H–C assignments from one-bond [23a] and long-range ¹H,¹³C COSY experiments [23b]. For natural oceanapins A–E (2–6), only the ¹³C- and ¹H-NMR signals that differ from those of oceanapin F (7) are reported; all other signals for oceanapins 2–6 proved to be superimposable. In analogy, for sphingosines 8b–d, only NMR signals that differ from those of 8a are reported. EI-MS (*m/z*, (%)): *Kratos-MS80* mass spectrometer with home-built data system and equipped with a *Vacumetrics-DIP* gun for FAB spectra. Molecular mechanics calculations: MMX, PCMOD.4, *Serena Software*, Bloomington, Indiana.

2. *Collection and Isolation*. The sponge was collected by scuba diving in Woodin Channel, New Caledonia, in August 1985 and was identified by Dr. *Jane Fromont*. The sponge was immediately frozen, freeze-dried (600 g), and CH₂Cl₂ extracted. The solvent was evaporated and the residue (5.04 g) subjected to FC (hexane/AcOEt gradient

elution) collecting 20 fractions of 100 ml each. *Fr. 13* was further subjected to FC (Et₂O) and then to HPLC (CN, hexane/*i*-PrOH 96:4), giving a mixture of oceanapins 2–7 (62 mg, 0.01%; *t_R* 8.7 min) which could be cleanly separated by reversed-phase HPLC (MeOH): 2 (6.0 mg), 3 (2.0 mg), 4 (4.9 mg), 5 (14.3 mg), 6 (6.2 mg), and 7 (21.5 mg); *t_R* values in the *Table*.

3. *Oceanapin A* (= N-[*(2S,3R,4E)*-1,3-Dihydroxyhexadec-4-en-2-yl]-21-methyl-docosanamide; 2). White powder. M.p. (hexane) 65–75°. ¹H-NMR: 1.24 (br. s, CH₂(8') to CH₂(15'), CH₂(4) to CH₂(20)); 0.87 (*t*, J(16', 15') = 6.6, 3 H-C(16')). ¹³C-NMR: 29.1–30.0 (series of *t*, C(7')–C(13'), C(4)–C(18)); 31.90 (*t*, C(14')); 22.67 (*t*, C(15')); 14.09 (*q*, C(16')). EI-MS: 589 (1, [M – H₂O]⁺), 572 (1), 558 (1), 379 (11), 378 (9), 337 (4), 111 (6), 43 (100).

4. *Oceanapin B* (= N-[*(2S,3R,4E)*-1,3-Dihydroxy-15-methylhexadec-4-en-2-yl]-21-methyl-docosanamide; 3). White powder. M.p. (hexane) 63–68°. ¹H-NMR: 1.24 (br. s, CH₂(8') to CH₂(14'), CH₂(4) to CH₂(20)); 1.49 (*m*, H-C(15'), H-C(21)); 0.85 (*d*, J = 6.6, 2 Me-C(15'), 2 Me-C(21)). ¹³C-NMR: 29.1–30.0 (series of *t*, C(7')–C(12'), C(4)–C(18)); 27.41 (*t*, C(13'), C(19)); 39.05 (*d*, C(14'), C(20)); 27.96 (*d*, C(15'), C(21)); 22.65 (*q*, C(16'), Me-C(15'), Me-C(22), Me-C(21)). EI-MS: 603 (1, [M – H₂O]⁺), 589 (1), 586 (0.7), 573 (1), 379 (17), 378 (14), 111 (7), 83 (26), 57 (63), 43 (100).

5. *Oceanapin C* (= N-[*(2S,3R,4E)*-1,3-Dihydroxy-15-methylhexadec-4-en-2-yl]tricosanamide; 4). White powder. M.p. (hexane) 65–70°. ¹H-NMR: 1.24 (br. s, CH₂(8') to CH₂(14'), CH₂(4) to CH₂(22)); 1.49 (*m*, H-C(15')), 0.85 (*d*, J = 6.6, 2 Me-C(15')), 0.87 (*t*, J(23,22) = 6.6, 3 H-C(23)). ¹³C-NMR: 29.1–30.0 (series of *t*, C(7')–C(12'), C(4)–C(20)); 27.41 (*t*, C(13')); 39.05 (*t*, C(14')); 27.96 (*d*, C(15')); 22.65 (*q*, C(16'), Me-C(15')); 31.90 (*t*, C(21)); 22.67 (*t*, C(22)); 14.09 (*q*, C(23)). EI-MS: 603 (0.5, [M – H₂O]⁺), 572 (1), 379 (3), 378 (3), 111 (6), 83 (24), 57 (60), 43 (100).

6. *Oceanapin D* (= N-[*(2S,3R,4E)*-1,3-Dihydroxy-16-methylheptadec-4-en-2-yl]-21-methyl-docosanamide; 5). White powder. M.p. (hexane) 70–72°. ¹H-NMR: 1.24 (br. s, CH₂(8') to CH₂(15'), CH₂(4) to CH₂(20)); 1.49 (*m*, H-C(16'), H-C(21)); 0.85 (*d*, J = 6.6, 2 Me-C(16'), 2 Me-C(21)). ¹³C-NMR: 29.1–30.0 (series of *t*, C(7')–C(13'), C(4)–C(18)); 27.40 (*t*, C(14'), C(19)); 39.05 (*t*, C(15'), C(20)); 27.95 (*d*, C(16'), C(21)); 22.65 (*q*, C(17'), Me-C(16'), C(22), Me-C(21)). EI-MS: 617 (0.8, [M – H₂O]⁺), 586 (1), 379 (21), 378 (18), 337 (2), 83 (27), 82 (20), 57 (62), 43 (100). FAB-MS (3-nitrobenzyl alcohol matrix): 658 (0.5, [M + Na]⁺), 618 (1.1, [M + H – H₂O]⁺).

7. *Oceanapin E* (= N-[*(2S,3R,4E)*-1,3-Dihydroxy-16-methyloctadec-4-en-2-yl]docosanamide; 6). White powder. M.p. (hexane) 65–68°. ¹H-NMR: 1.24 (br. s, CH₂(8') to CH₂(15'), CH₂(17'), CH₂(4) to CH₂(21)); 0.87 (*t*, J(22,21) = 6.6, 3 H-C(22)). ¹³C-NMR: 29.1–30.0 (series of *t*, C(7')–C(13'), C(17'), C(4)–C(19)); 31.90 (*t*, C(20)); 22.68 (*t*, C(21)), 14.09 (*q*, C(22)). EI-MS: 617 (2, [M – H₂O]⁺), 586 (3), 366 (22), 364 (31), 252 (4), 111 (13), 83 (48), 82 (45), 57 (94), 43 (100). HR-EI-MS: 364.3576 ± 0.0050 (C₂₄H₄₆NO, calc. 364.3579).

8. *Oceanapin F* (= N-[*(2S,3R,4E)*-1,3-Dihydroxy-16-methyloctadec-4-en-2-yl]-21-methyl-docosanamide; 7). White powder. M.p. (hexane) 70–75°. [α]_D²⁰ = –2.1, [α]_D²⁰₃₆₅ = –8.0 (c = 0.35, CHCl₃)². IR (nujol): 3360 (br.), 2920vs, 2850vs, 1640vs, 1470s, 1360m, 1178m, 1068s, 988m. ¹H-NMR: 3.70 (br. *dd*, J_{gem} = 11.0, 3.7), 3.96 (*dt*, J_{gem} = 11.0, 3.7, 2 H-C(1')), 3.89 (*ddt*, J = 6.6, 5.6, 3.7, H-C(2')), 4.29 (br. *ddd*, J(3',2') = 5.6, J(3',4') = 6.6, J(OH,3') = 4.7, H-C(3')), 2.62 (*d*, J(OH,3') = 4.7, OH-C(3')), 5.51 (*ddt*, J(4',5') = 15.3, J(4',3') = 6.6, J(4',6') = 1.2, H-C(4')), 5.76 (*ddd*, J(5',4') = 15.3, J(5',6') = 6.9, J(5',3') = 1.2, H-C(5')), 2.02 (*q*, J(6',5') ≈ J(6',7') = 6.9, CH₂(6')), 1.30 (*m*, CH₂(7'), H-C(16')), 1.24 (br. s, CH₂(8') to CH₂(15'), CH₂(17'), CH₂(4) to CH₂(20)); 0.84 (*t*, J(18',17') = 6.6, 3 H-C(18')), 6.30 (br. *d*, J(NH,2') = 6.6, NH-C(1)), 2.22 (*t*, J(2,3) = 7.5, CH₂(2)); 1.60 (*m*, CH₂(3)); 1.49 (*m*, H-C(21)); 0.85 (*d*, J = 6.6, 2 Me-C(21), Me-C(16')). ¹³C-NMR: 62.52 (*t*, C(1')), 54.45 (*d*, C(2')), 74.72 (*d*, C(3')), 128.80 (*d*, C(4')), 134.30 (*d*, C(5')), 32.25 (*t*, C(6')), 29.1–30.0 (series of *t*, C(7')–C(13'), C(17'), C(4)–C(18)); 27.10 (*t*, C(14')), 36.63 (*t*, C(15')), 34.38 (*d*, C(16')), 11.39 (*q*, C(18')); 19.21 (*q*, Me-C(16')); 173.85 (*s*, C(1)); 36.83 (*t*, C(2)); 25.74 (*t*, C(3)); 27.40 (*t*, C(19)); 39.05 (*t*, C(20)); 27.95 (*d*, C(21)); 22.64 (*q*, C(22), Me-C(21)). EI-MS: 631 (1, [M – H₂O]⁺), 614 (1), 602 (2), 379 (27), 378 (16), 350 (4), 337 (3), 252 (2), 111 (8), 83 (30), 82 (18), 57 (69), 43 (100). HR-EI-MS: 378.3735 ± 0.0040 (C₂₅H₄₈NO, calc. 378.3736).

9. *Acid Hydrolysis of Oceanapins A–F* (2–7). Each single, pure oceanapin (2–10 mg) was dissolved in 3 ml of 1.2M H₂SO₄ in 85% MeOH and heated at reflux for 6 h. The mixture was then cooled, treated with H₂O (1 ml), and then extracted with hexane (3 × 3 ml). The org. phase was percolated through a *Whatman* phase-separation filter and evaporated: methyl ester 9. The aq. layer was treated with 2M NaOH, then extracted with AcOEt (3 × 3 ml), percolated through a *Whatman* phase-separation filter, and finally evaporated: sphingosine 8.

²) Also for the other oceanapins, we obtained low [α]_D values, which, coupled to the scarce availability of these compounds, resulted in large uncertainties in [α]_D.

(4E)-2-Aminohexadec-4-ene-1,3-diol (8a): $^1\text{H-NMR}$: 3.67, 3.61 (2dd, $J_{\text{gem}} = 11.0$, $J(1,2) = 5.7$, $\text{CH}_2(1)$); 2.83 ('q', $J(2,1) \approx J(2,3) = 5.4$, $\text{H-C}(2)$); 4.03 (ddd, $J(3,4) = 6.6$, $J(3,2) = 5.4$, $J(3,5) = 1.2$, $\text{H-C}(3)$); 5.45 (ddt, $J(4,5) = 15.3$, $J(4,3) = 6.6$, $J(4,6) = 1.2$, $\text{H-C}(4)$); 5.73 (dtd, $J(5,4) = 15.3$, $J(5,6) = 6.6$, $J(5,3) = 1.2$, $\text{H-C}(5)$); 2.03 (q, $J(6,5) = J(6,7) = 6.6$, $\text{CH}_2(6)$); 1.35 (quint., $J(7,6) \approx J(7,8) = 6.6$, $\text{CH}_2(7)$); 1.24 (m, $\text{CH}_2(8)$ to $\text{CH}_2(15)$); 0.86 (t, $J(16,15) = 6.6$, 3 $\text{H-C}(16)$); 2.45 (br. s, OH). $^{13}\text{C-NMR}$: 64.04 (t, C(1)); 56.15 (d, C(2)); 75.39 (d, C(3)); 129.30 (d, C(4)); 134.68 (d, C(5)); 32.33 (t, C(6)); 29.7-29.2 (series of t, C(7)-C(13)); 31.90 (t, C(14)); 22.67 (t, C(15)); 14.09 (q, C(16)).

(4E)-2-Amino-15-methylhexadec-4-ene-1,3-diol (8b): $^1\text{H-NMR}$: 1.24 (m, $\text{CH}_2(8)$ to $\text{CH}_2(14)$); 1.50 (m, $\text{H-C}(15)$); 0.85 (d, $J = 6.6$, 2 $\text{Me-C}(15)$). FAB-MS (3-nitrobenzyl alcohol matrix): 286 (5, $[M + \text{H}]^+$), 268 (9, $[M + \text{H} - \text{H}_2\text{O}]^+$).

(4E)-2-Amino-16-methylheptadec-4-ene-1,3-diol (8c): $^1\text{H-NMR}$: 1.50 (m, $\text{H-C}(16)$); 0.85 (d, $J = 6.6$, 2 $\text{Me-C}(16)$). $^{13}\text{C-NMR}$: 27.41 (t, C(14)); 39.05 (t, C(15)); 27.96 (d, C(16)); 22.66 (q, C(17), $\text{Me-C}(16)$). FAB-MS (3-nitrobenzyl alcohol matrix): 300 (8, $[M + \text{H}]^+$), 282 (12, $[M + \text{H} - \text{H}_2\text{O}]^+$).

(4E)-2-Amino-16-methyloctadec-4-ene-1,3-diol (8d): $^1\text{H-NMR}$: 1.24 (br. s, $\text{CH}_2(8)$ to $\text{CH}_2(15)$, $\text{CH}_2(17)$); 1.40 (m, $\text{H-C}(16)$); 0.84 (t, $J(18,17) = 6.6$, 3 $\text{H-C}(18)$); 0.85 (d, $J(\text{Me-C}(16),16) = 6.6$, $\text{Me-C}(16)$). $^{13}\text{C-NMR}$: 29.1-29.9 (series of t, C(7)-C(13), C(17)); 27.41 (t, C(14)); 36.33 (t, C(15)); 34.39 (d, C(16)); 11.40 (q, C(18)); 19.21 (q, $\text{Me-C}(16)$). FAB-MS (3-nitrobenzyl alcohol matrix): 314 (4, $[M + \text{H}]^+$), 296 (6, $[M + \text{H} - \text{H}_2\text{O}]^+$).

Methyl 21-Methyldocosanoate (9a): $^1\text{H-NMR}$: 2.29 (t, $J(2,3) = 7.2$, $\text{CH}_2(2)$); 1.24 (br. s, $\text{CH}_2(3)$ to $\text{CH}_2(20)$); 1.50 (m, $\text{H-C}(21)$); 0.85 (d, $J = 6.6$, 2 $\text{Me-C}(21)$); 3.65 (s, MeO). $^{13}\text{C-NMR}$: 174.37 (s, C(1)); 34.12 (t, C(2)); 24.96 (t, C(3)); 31.92 (t, C(4)); 29.2-29.7 (series of t, C(5)-C(18)); 27.42 (t, C(19)); 39.05 (t, C(20)); 27.96 (d, C(21)); 22.66 (q, 2 $\text{Me-C}(21)$); 51.44 (q, MeO). EI-MS: 368 (39, M^+), 337 (2), 325 (9), 87 (70), 74 (100).

Methyl Tricosanoate (9b): $^1\text{H-NMR}$: 2.29 (t, $J(2,3) = 7.2$, $\text{CH}_2(2)$); 1.24 (m, $\text{CH}_2(3)$ to $\text{CH}_2(22)$); 0.87 (t, $J = 6.9$, 3 $\text{H-C}(23)$). $^{13}\text{C-NMR}$: 174.4 (s, C(1)); 34.12 (t, C(2)); 24.9 (t, C(3)); 31.91 (t, C(4)); 29.2-29.7 (series of t, C(5)-C(20)); 31.92 (t, C(21)); 22.69 (t, C(22)); 14.10 (q, C(23)); 51.44 (q, MeO). EI-MS: 368 (10, M^+), 337 (1), 325 (9), 87 (70), 74 (100).

Methyl Docosanoate (9c): $^1\text{H-NMR}$: 2.29 (t, $J(2,3) = 7.2$, $\text{CH}_2(2)$); 1.24 (br. s, $\text{CH}_2(3)$ to $\text{CH}_2(21)$); 0.87 (t, $J = 6.9$, 3 $\text{H-C}(22)$). $^{13}\text{C-NMR}$: 174.37 (s, C(1)); 34.12 (t, C(2)); 24.96 (t, C(3)); 31.91 (t, C(4)); 29.1-29.7 (series of t, C(5)-C(19)); 31.92 (t, C(20)); 22.69 (t, C(21)); 14.11 (q, C(22)); 51.44 (q, MeO). EI-MS: 354 (8, M^+), 87 (70), 74 (100).

10. N-*f*(2*S*,3*R*,4*E*)-1,3-(Isopropylidenedioxy)octadec-4-enyl]acetamide (11). To D-erythro-sphingosine (= (2*S*,3*R*,4*E*)-2-aminooctadec-4-ene-1,3-diol; 1; *Sigma*; 3.0 mg, 0.010 mmol) in EtOH (0.5 ml) was added an excess of Ac_2O and stirred for 1 h at r.t. The mixture was evaporated to give monoacetate 10 (3.3 mg, 97%) which was then dissolved in dry acetone (1 ml). Excess dry CuSO_4 was added and the mixture heated at reflux with stirring for 6 h, then cooled, filtered, and evaporated: 11 (3.2 mg, 87%).

N-Acetoxy-sphingosine (= N-*f*(2*S*,3*R*,4*E*)-1,3-Dihydroxyoctadec-4-en-2-yl]acetamide; 10): $^1\text{H-NMR}$: 3.95, 3.70 (2 br. d, $J_{\text{gem}} = 10.8$, $\text{CH}_2(1)$); 3.89 (m, $\text{H-C}(2)$); 4.31 (m, $\text{H-C}(3)$); 5.52 (dd, $J(4,5) = 15.3$, $J(4,3) = 6.3$, $\text{H-C}(4)$); 5.78 (dt, $J(5,4) = 15.3$, $J(5,6) = 6.9$, $\text{H-C}(5)$); 2.04 (m, $\text{CH}_2(6)$); 1.35 (m, $\text{CH}_2(7)$); 1.24 (br. s, $\text{CH}_2(8)$ to $\text{CH}_2(17)$); 0.87 (t, $J(18,17) = 6.9$, 3 $\text{H-C}(18)$); 2.03 (s, NHCOMe). $^{13}\text{C-NMR}$: 62.39 (t, C(1)); 54.37 (d, C(2)); 74.35 (d, C(3)); 128.70 (d, C(4)); 134.29 (d, C(5)); 32.26 (t, C(6)); 29.1-29.7 (series of t, C(7)-C(15)); 31.91 (t, C(16)); 22.68 (t, C(17)); 14.10 (q, C(18)); 170.74 (s, NHCOMe); 23.40 (q, MeCO). EI-MS: 324 (1), 323 (0.5, $[M - \text{H}_2\text{O}]^+$), 310 (1), 292 (1), 102 (69), 85 (98), 57 (24), 43 (100).

Data of 11: $^1\text{H-NMR}$: 4.00, 3.61 (2dd, $J_{\text{gem}} = 11.3$, $J(1,2) = 5.2$, 9.1, $\text{CH}_2(1)$); 3.83 (dddd, $J(2,1) = 5.2$, 9.1, $J(2,3) = 9.6$, $J(2,\text{NH}) = 8.3$, $\text{H-C}(2)$); 4.04 (dd, $J(3,2) = 9.6$, $J(3,4) = 7.5$, $\text{H-C}(3)$); 5.41 (ddt, $J(4,5) = 15.4$, $J(4,3) = 7.5$, $J(4,6) = 1.4$, $\text{H-C}(4)$); 5.74 (dt, $J(5,4) = 15.3$, $J(5,6) = 6.8$, $\text{H-C}(5)$); 2.02 (q, $J = 6.8$, $\text{CH}_2(6)$); 1.35 (m, $\text{CH}_2(7)$); 1.24 (br. s, $\text{CH}_2(8)$ to $\text{CH}_2(17)$); 0.87 (t, $J(18,17) = 6.9$, 3 $\text{H-C}(18)$); 2.04 (s, NHCOMe); 1.47, 1.41 (2s, Me_2C); 5.19 (br. d, $J = 8.4$, NH). $^{13}\text{C-NMR}$: 62.70 (t, C(1)); 48.28 (d, C(2)); 74.12 (d, C(3)); 127.25 (d, C(4)); 136.53 (d, C(5)); 32.30 (t, C(6)); 29.1-29.7 (series of t, C(7)-C(15)); 31.90 (t, C(16)); 22.67 (t, C(17)); 14.10 (q, C(18)); 169.75 (s, NHCOMe); 23.38 (q, MeCO); 98.87 (s, Me_2C); 28.40, 28.96 (2q, Me_2C). EI-MS: 366 (1.6, $[M - 15]^+$), 324 (2), 323 (1), 102 (10), 85 (65), 82 (23), 57 (49), 43 (100).

11. N-*f*(2*S*,3*R*,4*E*)-1,3-(Isopropylidenedioxy)-16-methyloctadec-4-en-2-yl]-21-methyldocosanamide (12). As described for 11, 7 (3.9 mg, 0.06 mmol) was transformed into 12 (4.0 mg, 97%). $^1\text{H-NMR}$: 3.99, 3.64 (2dd, $J_{\text{gem}} = 11.3$, $J(1',2') = 5.1$ or 9.3, $\text{CH}_2(1')$); 3.82 (dddd, $J(2',1') = 9.3$, 5.3, $J(2',3') = 9.6$, $J(2',\text{NH}) = 8.1$, $\text{H-C}(2')$); 4.07 (dd, $J(3',2') = 9.6$, $J(3',4') = 7.8$, $\text{H-C}(3')$); 3.82 (dddd, $J(2',1'a) = 5.1$, $J(2',1'b) = 9.3$, $J(2',3') = 9.6$, $J(2',\text{NH}) = 8.1$, $\text{H-C}(2')$); 5.39 (ddt, $J(4',5') = 15.3$, $J(4',3') = 7.8$, $J(4',6') = 1.2$, $\text{H-C}(4')$); 5.13 (d, $J(\text{NH},2') = 8.1$, NH); 1.48, 1.41 (2s, Me_2C). $^{13}\text{C-NMR}$: 62.72 (t, C(1')); 48.11 (d, C(2')); 74.10 (d, C(3')); 127.28 (d,

C(4''); 136.51 (*d*, C(5'')); 32.33 (*t*, C(6'')); 29.2-29.7 (series of *t*, C(7'')-C(13''), C(17''), C(4)-C(18)); 27.11 (*t*, C(14'')); 36.64 (*t*, C(15'')); 34.39 (*d*, C(16'')); 19.21 (*q*, *Me*-C(16'')); 11.40 (*q*, C(18'')); 172.88 (*s*, C(1)); 36.89 (*t*, C(2)); 25.69 (*t*, C(3)); 27.41 (*t*, C(19)); 39.05 (*t*, C(20)); 27.95 (*d*, C(21)); 22.65 (*q*, 2 *Me*-C(21)); 98.83 (*s*, *Me*₂C); 28.99, 28.47 (2*s*, *Me*₂C). EI-MS: 674 (2, [*M* - 15]⁺), 632 (1.5), 379 (63), 378 (12), 278 (18), 85 (25), 82 (23), 57 (60), 43 (100).

12. Mosher's Esters. A soln. of 5 (9.0 mg, 0.014 mmol), 1 H-imidazole (2.5 mg, 0.036 mmol), and (*tert*-butyl)diphenylsilyl chloride (Aldrich; 5.0 mg, 0.018 mol) in dry DMF (1 ml) was stirred at 60° under N₂ for 12 h. The mixture was cooled, treated with H₂O (2 ml), and extracted with AcOEt (3 × 3 ml). The org. phase was washed with sat. aq. NaCl soln., dried (Na₂SO₄), and evaporated to give a residue that was subjected to prep. TLC (hexane/AcOEt 7:3): pure (+)-13 (10.8 mg, 89%), *R*_f 0.7. To a soln. of (+)-13 (5.3 mg, 0.006 mmol) in dry pyridine (0.2 ml) and CCl₄ (0.2 ml) were added 3 mol-equiv. of (+)-(*S*)-MTPACI (Aldrich) and stirred at r.t. for 24 h. Sat. aq. CuSO₄ soln. (3 ml) was added and the mixture percolated through a Whatman phase-separation filter. The filtrate was evaporated and the residue subjected to HPLC (CN, hexane/*i*-PrOH 98:2, λ = 225 nm): pure (+)-14a (*t*_R 5.3 min; 4.8 mg, 90%) besides unreacted (+)-13 (*t*_R 9.3 min, 1 mg).

Following the same procedure with (-)-(*R*)-MTPACI and (+)-13 (4.0 mg), pure (-)-14b (*t*_R 5.5 min; 4.4 mg, 88%) was obtained.

N-[(2*S*,3*R*,4*E*)-1-[(*tert*-Butyl)diphenylsilyloxy]-3-hydroxy-16-methylheptadec-4-en-2-yl]-21-methyl-docosanamide ((+)-13): [α]_D²⁰ = +5, [α]_D²⁰ = +37 (*c* = 0.4, CHCl₃). ¹H-NMR: 3.95, 3.75 (2*m*, CH₂(1'')); 4.20 (*m*, H-C(3'')); 5.46 (*ddt*, *J*(4',5') = 15.3, *J*(4',3') = 6.0, *J*(4',6') = 1.2, H-C(4'')); 5.76 (*dd*, *J*(5',4') = 15.3, *J*(5',6') = 6.9, *J*(5',3') = 1.2, H-C(5'')); 2.02 (*q*', *J*(6',7') ≈ *J*(6',5') = 6.9, CH₂(6'')); 1.30-1.20 (*m*, CH₂(7') to CH₂(15'')), CH₂(4) to CH₂(20)); 1.50 (*m*, H-C(16'')), H-C(21)); 0.85 (*d*, *J* = 6.6, 2 *Me*-C(16'), 2 *Me*-C(21)); 6.10 (*d*, *J*(NH,2') = 8.1, NH); 2.18 (*t*, *J*(2,3) = 7.5, CH₂(2)); 1.60 (*m*, CH₂(3)); 3.56 (*br. d*, *J* = 7.2, OH); 1.06 (*s*, *Me*₃C); 7.61, 7.39 (2*m*, Ph). ¹³C-NMR: 63.98 (*t*, C(1'')); 53.92 (*d*, C(2'')); 74.28 (*d*, C(3'')); 128.93 (*d*, C(4'')); 133.38 (*d*, C(5'')); 32.32 (*t*, C(6'')); 29.2-29.9 (series of *t*, C(7'')-C(13''), C(4)-C(18)); 27.41 (*t*, C(14'')), C(19)); 36.82 (*t*, C(15'')); 27.95 (*t*, C(16'')), C(21)); 22.65 (*q*, C(17'')), *Me*-C(16'')), C(22), *Me*-C(21)); 173.33 (*s*, C(1)); 36.82 (*t*, C(2)); 25.76 (*t*, C(3)); 26.84 (*q*, *Me*₃C); 19.14 (*s*, *Me*₃C); 132.37 (*s*, 2 C, Ph); 130.07 (*d*, 2 C, Ph); 127.89 (*d*, 4 C, Ph); 135.49 (*d*, 4 C, Ph). EI-MS: 855 (0.5, [*M* - H₂O]⁺), 799 (45), 798 (68), 264 (9), 199 (73), 57 (50), 43 (47).

(2*S*,3*R*,4*E*)-1-[(*tert*-Butyl)diphenylsilyloxy]-16-methyl-2-(21-methyl-docosanamido)hexadec-4-en-3-yl (*R*)-3,3,3-Trifluoro-2-methoxy-2-phenylpropanoate ((+)-14a): [α]_D²⁰ = +8.0 (*c* = 0.2, CHCl₃). ¹H-NMR: 3.78, 3.66 (*AB* of *ABX*, *J*(*AB*) = 10.8, *J*(*AX*) = 3.9, *J*(*BX*) = 4.2, CH₂(1'')); 4.29 (*m*, H-C(2'')); 5.63 (*t*, *J*(2',3') ≈ *J*(3',4') = 7.5, H-C(3'')); 5.27 (*ddt*, *J*(4',5') = 15.3, *J*(4',3') = 7.2, *J*(4',6') = 0.9, H-C(4'')); 5.80 (*dt*, *J*(5',4') = 15.3, *J*(5',6') = 6.9, H-C(5'')); 1.95 (*m*, CH₂(6'')); 1.20-1.30 (*m*, CH₂(7') to CH₂(15'')); 1.50 (*m*, H-C(16'')), H-C(21'')); 0.85 (*d*, *J* = 6.6, 2 *Me*-C(16'), 2 *Me*-C(21'')); 1.97 (*m*, CH₂(2'')); 1.60 (*m*, CH₂(3'')); 5.44 (*br. d*, *J*(NH,2') = 9.0, NH); 1.06 (*s*, *Me*₃C); 7.59, 7.40 (2*m*, Ph); 3.37 (*q*, ³*J*(H,F) = 0.9, MeO). ¹³C-NMR: 63.55 (*t*, C(1'')); 51.71 (*d*, C(2'')); 75.77 (*t*, C(3'')); 32.26 (*t*, C(6'')); 29.2-30.0 (series of *t*, C(7'')-C(13''), C(4'')-C(18'')); 27.42 (*t*, C(14'')), C(19'')); 36.82 (*t*, C(15'')); 27.96 (*d*, C(16'')), C(21'')); 22.65 (*q*, 2 *Me*-C(16'), 2 *Me*-C(21'')); 172.40 (*s*, C(1'')); 36.82 (*t*, C(2'')); 25.63 (*t*, C(3'')); 26.88 (*q*, *Me*₃C); 19.27 (*s*, *Me*₃C); 55.40 (*q*, MeO); 138.99 (*d*); 135.54 (*d*); 130.02 (*d*); 129.51 (*d*); 128.27 (*d*); 127.90 (*d*); 127.42 (*d*); 123.60 (*d*). EI-MS: 856 (1, [*M* - MTPAO]⁺), 799 (100), 462 (27), 199 (23), 57 (86), 43 (62).

(2*S*,3*R*,4*E*)-1-[(*tert*-Butyl)diphenylsilyloxy]-16-methyl-2-(21-methyl-docosanamido)hexadec-4-en-3-yl (*S*)-3,3,3-Trifluoro-2-methoxy-2-phenylpropanoate ((-)-14b): [α]_D²⁰ = -10.0 (*c* = 0.2, CHCl₃). ¹H-NMR (only signals that differ from (+)-14a are reported): 3.65, 3.60 (*AB* of *ABX*, *J*(*AB*) = 10.5, *J*(*AX*) ≈ *J*(*BX*) = 3.6, CH₂(1'')); 4.25 (*m*, H-C(2'')); 5.65 (*t*, *J*(2',3') ≈ *J*(3',4') = 7.5, H-C(3'')); 5.43 (*ddt*, *J*(4',5') = 15.3, *J*(4',3') = 7.3, *J*(4',6') = 1.3, H-C(4'')); 5.91 (*dt*, *J*(5',4') = 15.3, *J*(5',6') = 6.9, H-C(5'')); 1.99 (*m*, CH₂(6'')); 1.93 (*m*, CH₂(2'')); 5.41 (*br. d*, *J*(NH,2') = 9.0, NH); 3.48 (*q*, ³*J*(H,F) = 0.9, MeO). ¹³C-NMR (only signals are reported that differ from those of (+)-14a): 76.03 (*t*, C(3'')); 172.50 (*s*, C(1'')); 36.75 (*t*, C(2'')); 55.45 (*q*, MeO); 139.78 (*d*); 135.53 (*d*); 129.99 (*d*); 128.69 (*d*); 128.34 (*d*); 127.51 (*d*); 127.26 (*d*); 124.00 (*d*). EI-MS: practically superimposable to that of (+)-14a.

REFERENCES

- [1] H. E. Carter, F. J. Glick, W. P. Norris, G. E. Phillips, *J. Biol. Chem.* **1947**, *170*, 285.
- [2] J. L. W. Thudicum, 'Die Konstitution des Gehirns des Menschen und der Tiere'. Tübingen, 1901.
- [3] A. R. Johnson, J. B. Davenort, Eds., 'Biochemistry and Methodology of Lipids'. Wiley-Interscience, New York, 1971.
- [4] a) G. Aguilar-Santos, M. S. Doty, in 'Drugs from the Sea', Ed. H. D. Freudenthal. Marine Technology Society, Washington, 1967; b) M. Mahendran, S. Somasundaram, R. H. Thomson, *Phytochemistry* **1979**, *18*, 1885; c) P. G. Nielsen, J. S. Carlié, C. Christophersen, *Phytochemistry* **1982**, *21*, 1643.

- [5] H. S. Garg, M. Sharma, D. S. Bhakuni, B. N. Pramanik, A. K. Bose, *Tetrahedron Lett.* **1992**, *33*, 1641.
- [6] J. H. Cardellina, R. E. Moore, *Phytochemistry* **1978**, *17*, 554.
- [7] J. Kobayashi, M. Ishibashi, H. Nakamura, Y. Hirata, T. Yamasu, T. Sasaki, Y. Ohizumi, *Experientia* **1988**, *44*, 800.
- [8] K. Chebaane, M. Guyot, *Tetrahedron Lett.* **1986**, *27*, 1495.
- [9] F. Cafieri, E. Fattorusso, *Liebigs Ann. Chem.* **1990**, 1441.
- [10] M. Endo, M. Nagakawa, Y. Hamamoto, *Pure Appl. Chem.* **1986**, *58*, 387.
- [11] S. G. Batrakov, V. B. Muratov, O. G. Sakandelidze, A. V. Sulima, B. V. Rozynov, *Soviet J. Biorg. Chem.* **1983**, *9*, 292.
- [12] S. Hirsch, Y. Kashman, *Tetrahedron* **1989**, *45*, 3897.
- [13] J. D. Faulkner, *Nat. Prod. Rept.* **1984**, *1*, 251.
- [14] F. W. Wehrli, A. P. Marchand, S. Wehrli, 'Interpretation of Carbon-13 NMR Spectra', 2nd edn., Wiley, 1988, p. 56-57.
- [15] K. Munesata, M. Yuasa, T. Suga, *J. Chem. Soc., Perkin Trans. 1* **1991**, 189.
- [16] M. Nakagawa, J. Yoshida, T. Hino, *Chem. Lett.* **1990**, 1407.
- [17] M. Nakagawa, A. Tsuruoka, J. Yoshida, T. Hino, *J. Chem. Soc., Chem. Commun.* **1990**, 603.
- [18] J. A. Dale, D. L. Dull, H. S. Mosher, *J. Org. Chem.* **1969**, *34*, 2543; J. A. Dale, H. S. Mosher, *J. Am. Chem. Soc.* **1973**, *95*, 512; G. R. Sullivan, J. A. Dale, H. S. Mosher, *J. Org. Chem.* **1973**, *38*, 2143; I. Ohtani, T. Kusumi, Y. Kashman, H. Kakisawa, *J. Am. Chem. Soc.* **1991**, *113*, 4092.
- [19] S. Hanessian, P. Lavalley, *Can. J. Chem.* **1975**, *53*, 2975.
- [20] J. K. M. Sanders, J. D. Mersch, *Prog. N. M. R. Spectrosc.* **1982**, *15*, 353.
- [21] A. Bax, R. Freeman, G. Morris, *J. Magn. Reson.* **1981**, *42*, 164.
- [22] D. H. Doddrell, D. T. Pegg, H. R. Bendall, *J. Magn. Reson.* **1982**, *48*, 323; D. T. Pegg, D. M. Doddrell, M. R. Bendall, *J. Chem. Phys.* **1982**, *77*, 2745.
- [23] a) A. Bax, *J. Magn. Reson.* **1983**, *43*, 517; b) H. Kessler, C. Criesinger, J. Zarbock, H. R. Loosli, *ibid.* **1984**, *57*, 331.